

Palaniappan Kannappan; Pushpa Narayan Rathie  
On a generalized directed-divergence function

*Czechoslovak Mathematical Journal*, Vol. 24 (1974), No. 1, 5–14

Persistent URL: <http://dml.cz/dmlcz/101212>

## Terms of use:

© Institute of Mathematics AS CR, 1974

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A GENERALIZED DIRECTED-DIVERGENCE FUNCTION

PL. KANNAPPAN and P. N. RATHIE, Waterloo

(Received March 15, 1971)

1. INTRODUCTION AND DEFINITIONS

Consider three finite discrete probability distributions  $P = (p_1, \dots, p_n)$ ,  $p_i \geq 0$ ,  $\sum_{i=1}^n p_i = 1$ ,  $Q = (q_1, \dots, q_n)$ ,  $q_i \geq 0$ ,  $\sum_{i=1}^n q_i = 1$  and  $R = (r_1, \dots, r_n)$ ,  $r_i \geq 0$ ,  $\sum_{i=1}^n r_i = 1$ . For these probability distributions, we defined in an earlier paper [7] the following generalized directed-divergence of type  $\beta$  ( $\beta \neq 1$ ):

$$(1) \quad I_n^\beta \begin{pmatrix} p_1, \dots, p_n \\ q_1, \dots, q_n \\ r_1, \dots, r_n \end{pmatrix} = (2^{\beta-1} - 1) \left[ \sum_{i=1}^n p_i q_i^{\beta-1} r_i^{1-\beta} - 1 \right], \quad \beta \neq 1.$$

For  $\beta \rightarrow 1$ , (1) gives

$$(2) \quad I_n^1 \begin{pmatrix} p_1, \dots, p_n \\ q_1, \dots, q_n \\ r_1, \dots, r_n \end{pmatrix} = \sum_{i=1}^n p_i \log (q_i/r_i).$$

For  $P \equiv Q$ , (2) reduces to the well-known directed-divergence [3]

$$(3) \quad I_n^1 \begin{pmatrix} p_1, \dots, p_n \\ p_1, \dots, p_n \\ r_1, \dots, r_n \end{pmatrix} = \sum_{i=1}^n p_i \log (p_i/r_i).$$

The object of this paper is to generalize further the quantities defined by (1) and (2) by introducing the concept of generalized directed-divergence and to characterize this by means of a functional equation. The generalizations are given in the following definitions:

**Definition 1.** A real valued function  $f$  on  $I \times I \times I$  where  $I = [0, 1]$ , is called a generalized directed-divergence function of order  $\alpha (\neq 1)$  and type  $\beta (\beta \neq 1)$  if  $f$  is a solution of the functional equation

$$(4) \quad f(x, y, z) + (1-x)^\alpha (1-y)^{\beta-1} (1-z)^{1-\beta} f\left(\frac{u}{1-x}, \frac{v}{1-y}, \frac{w}{1-z}\right) = \\ = f(u, v, w) + (1-u)^\alpha (1-v)^{\beta-1} (1-w)^{1-\beta} f\left(\frac{x}{1-u}, \frac{y}{1-v}, \frac{z}{1-w}\right),$$

for  $x, y, z, u, v, w \in [0, 1)$  with  $x+u, y+v, z+w \in I$ , satisfying further

$$(5) \quad f(0, 0, 0) = f(1, 1, 1),$$

and

$$(6) \quad f\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) = (2^{1-\alpha} - 1)(2^{\beta-1} - 1)^{-1}.$$

**Definition 2.** If  $f$  is a generalized directed-divergence function of order  $\alpha (\neq 1)$  and type  $\beta (\neq 1)$  as defined above, then the generalized directed-divergence of order  $\alpha (\neq 1)$  and type  $\beta (\neq 1)$  is defined by the expression:

$$(7) \quad I_{\alpha, n}^\beta \begin{pmatrix} p_1, \dots, p_n \\ q_1, \dots, q_n \\ r_1, \dots, r_n \end{pmatrix} = \sum_{i=2}^n P_i^\alpha Q_i^{\beta-1} R_i^{1-\beta} f\left(\frac{p_i}{P_i}, \frac{q_i}{Q_i}, \frac{r_i}{R_i}\right)$$

where  $P_i = p_1 + \dots + p_i$ ,  $Q_i = q_1 + \dots + q_i$ ,  $R_i = r_1 + \dots + r_i$ , for  $i = 1, 2, \dots, n$  with  $P_n = Q_n = R_n = 1$ .

In § 2 the solution of (4) and an expression for (7) are given. In § 3, some interesting special cases are pointed out.

The results (4) and (7) unify and generalize all the existing results in this direction. For example (7) reduces to (i) Shannon's entropy when  $y = z$  and  $\alpha = \beta \rightarrow 1$  in (4); (ii) informatic function of typed introduced in [2] for  $x = y = z$  and  $\beta = 1$  in (4); (iii) directed divergence [1], [5].

## 2. THIS SECTION DEALS WITH TWO THEOREMS WHICH GIVE $f$ AND $I_{\alpha, n}^\beta$ RESPECTIVELY

**Theorem 1.** If  $f$  is a solution of the functional equation (4) satisfying the additional conditions (5) and (6), then  $f$  is given by

$$(8) \quad f(x, y, z) = [x^\alpha y^{\beta-1} z^{1-\beta} + (1-x)^\alpha (1-y)^{\beta-1} (1-z)^{1-\beta} - 1](2^{\beta-1} - 1)^{-1},$$

for all  $x, y, z \in I$  and conversely.

**Remark 1.** The solution  $f$  of (4) has the form (8), provided we use the notation  $0^\gamma = 0$  (for  $\gamma \neq 0$ ). Only to put  $f$  in the form given by (8), this notation is used, but nowhere in the proof of the theorem,  $0^\gamma = 0$  is used.

Proof. Putting  $x = 0$ ,  $y = 0$  and  $z = 0$  in (5), we get

$$[1 - (1 - u)^\alpha (1 - v)^{\beta-1} (1 - w)^{1-\beta}] f(0, 0, 0) = 0, \quad \text{for } u, v, w \in [0, 1],$$

which in turn gives  $f(0, 0, 0) = 0$ . Hence from (4), we have

$$(9) \quad f(1, 1, 1) = f(0, 0, 0) = 0.$$

Replacing  $u, v$  and  $w$  in (4) by  $1 - x, 1 - y$  and  $1 - z$  respectively and using (9), we obtain

$$(10) \quad f(x, y, z) = f(1 - x, 1 - y, 1 - z), \quad \text{for all } x, y, z \in (0, 1).$$

Setting

$$p = \frac{u}{1 - x}, \quad q = \frac{v}{1 - y}, \quad r = \frac{w}{1 - z}, \quad \xi = 1 - x, \quad \eta = 1 - y \quad \text{and} \quad \zeta = 1 - z$$

in (4), (4) can be rewritten as

$$(11) \quad f(1 - \xi, 1 - \eta, 1 - \zeta) + \xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} f(p, q, r) = f(p\xi, q\eta, r\zeta) + \\ + (1 - p\xi)^\alpha (1 - q\eta)^{\beta-1} (1 - r\zeta)^{1-\beta} f\left(\frac{1 - \xi}{1 - p\xi}, \frac{1 - \eta}{1 - q\eta}, \frac{1 - \zeta}{1 - r\zeta}\right),$$

for all  $p, q, r \in I$ ,  $\xi, \eta, \zeta \in (0, 1]$ , such that  $p\xi \neq 1$ ,  $q\eta \neq 1$  and  $r\zeta \neq 1$ .

From (10) and (11), we get

$$(12) \quad f(\xi, \eta, \zeta) + \xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} f(p, q, r) = f(p\xi, q\eta, r\zeta) + \\ + (1 - p\xi)^\alpha (1 - q\eta)^{\beta-1} (1 - r\zeta)^{1-\beta} f\left(\frac{1 - \xi}{1 - p\xi}, \frac{1 - \eta}{1 - q\eta}, \frac{1 - \zeta}{1 - r\zeta}\right),$$

holding for  $p, q, r \in I$  and  $\xi, \eta, \zeta \in (0, 1)$ .

Define

$$(13) \quad F(p, q, r, \xi, \eta, \zeta) = \\ = f(\xi, \eta, \zeta) + [\xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} + (1 - \xi)^\alpha (1 - \eta)^{\beta-1} (1 - \zeta)^{1-\beta}] f(p, q, r),$$

where  $p, q, r, \xi, \eta, \zeta \in (0, 1)$ .

We shall prove that

$$(14) \quad F(p, q, r, \xi, \eta, \zeta) = F(\xi, \eta, \zeta, p, q, r).$$

Now (12) and (13) yield

$$(15) \quad F(p, q, r, \xi, \eta, \zeta) = f(p\xi, q\eta, r\zeta) + (1 - p\xi)^\alpha (1 - q\eta)^{\beta-1} (1 - r\zeta)^{1-\beta} \cdot \left\{ f\left(\frac{1-\xi}{1-p\xi}, \frac{1-\eta}{1-q\eta}, \frac{1-\zeta}{1-r\zeta}\right) + \left(\frac{1-\xi}{1-p\xi}\right)^\alpha \left(\frac{1-\eta}{1-q\eta}\right)^{\beta-1} \left(\frac{1-\zeta}{1-r\zeta}\right)^{1-\beta} f(p, q, r) \right\},$$

for all  $p, q, r, \xi, \eta, \zeta \in (0,1)$ . Allowing

$$A = \frac{1-\xi}{1-p\xi}, \quad B = \frac{1-\eta}{1-q\eta} \quad \text{and} \quad C = \frac{1-\zeta}{1-r\zeta}$$

and making use of (10) and (12), we have

$$(16) \quad \begin{aligned} f\left(\frac{1-\xi}{1-p\xi}, \frac{1-\eta}{1-q\eta}, \frac{1-\zeta}{1-r\zeta}\right) + \left(\frac{1-\xi}{1-p\xi}\right)^\alpha \left(\frac{1-\eta}{1-q\eta}\right)^{\beta-1} \left(\frac{1-\zeta}{1-r\zeta}\right)^{1-\beta} f(p, q, r) &= \\ &= f(A, B, C) + A^\alpha B^{\beta-1} C^{1-\beta} f(p, q, r) = f(pA, qB, rC) + \\ &+ (1-pA)^\alpha (1-qB)^{\beta-1} (1-rC)^{1-\beta} f\left(\frac{1-A}{1-pA}, \frac{1-B}{1-qB}, \frac{1-C}{1-rC}\right) = \\ &= f(1-pA, 1-qB, 1-rC) + (1-pA)^\alpha (1-qB)^{\beta-1} (1-rC)^{1-\beta} \cdot \\ &\cdot f\left(\frac{1-A}{1-pA}, \frac{1-B}{1-qB}, \frac{1-C}{1-rC}\right) = f\left(\frac{1-p}{1-p\xi}, \frac{1-q}{1-q\eta}, \frac{1-r}{1-r\zeta}\right) + \\ &+ \left(\frac{1-p}{1-p\xi}\right)^\alpha \left(\frac{1-q}{1-q\eta}\right)^{\beta-1} \left(\frac{1-r}{1-r\zeta}\right)^{1-\beta} f(\xi, \eta, \zeta). \end{aligned}$$

The equation (15) and (16) indeed go to show that (14) is true for all  $p, q, r, \xi, \eta, \zeta \in (0, 1)$ . Thus from (13) and (14) result

$$(17) \quad \begin{aligned} f(p, q, r) + [p^\alpha q^{\beta-1} r^{1-\beta} + (1-p)^\alpha (1-q)^{\beta-1} (1-r)^{1-\beta}] f(\xi, \eta, \zeta) &= \\ = f(\xi, \eta, \zeta) + [\xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} + (1-\xi)^\alpha (1-\eta)^{\beta-1} (1-\zeta)^{1-\beta}] f(p, q, r), \end{aligned}$$

where  $\xi, \eta, \zeta, p, q, r \in (0, 1)$ .

Now (17) with  $\xi = \frac{1}{2}, \eta = \frac{1}{2}$  and  $\zeta = \frac{1}{2}$  and (6) give

$$(18) \quad \begin{aligned} f(p, q, r) &= \\ &= [p^\alpha q^{\beta-1} r^{1-\beta} + (1-p)^\alpha (1-q)^{\beta-1} (1-r)^{1-\beta} - 1] (2^{\beta-1} - 1)^{-1}, \end{aligned}$$

for all  $p, q, r \in (0, 1)$ .

Next, we will show that (18) remains true for all  $p, q, r \in I$ , that is, (8) is valid.

The equation (11) with  $\xi = 1, \eta = 1$  and  $\zeta = \frac{1}{2}$  gives,

$$(19) \quad \begin{aligned} & f(0, 0, \frac{1}{2}) + 2^{\beta-1} f(p, q, r) = \\ & = f\left(p, q, \frac{r}{2}\right) + (1-p)^\alpha (1-q)^{\beta-1} \left(1 - \frac{r}{2}\right)^{1-\beta} f\left(0, 0, \frac{1}{2-r}\right), \end{aligned}$$

for all  $p, q \in [0, 1)$  and  $r \in I$ .

For  $r = \frac{1}{2}$ , (19) becomes,

$$(20) \quad \begin{aligned} & f(0, 0, \frac{1}{2}) + 2^{\beta-1} f(p, q, \frac{1}{2}) = \\ & = f(p, q, \frac{1}{4}) + (1-p)^\alpha (1-q)^{\beta-1} \left(\frac{3}{4}\right)^{1-\beta} f(0, 0, \frac{2}{3}) \end{aligned}$$

where  $p, q \in [0, 1)$ .

Taking  $p = \frac{1}{2}$  and  $q = \frac{1}{2}$  in (20) and using (6), we have

$$(21) \quad \begin{aligned} & f(0, 0, \frac{1}{2}) + 2^{\beta-1} (2^{1-\alpha} - 1) (2^{\beta-1} - 1)^{-1} = \\ & = f(\frac{1}{2}, \frac{1}{2}, \frac{1}{4}) + 2^{1-\alpha-\beta} \left(\frac{3}{4}\right)^{1-\beta} f(0, 0, \frac{2}{3}). \end{aligned}$$

From (18), (20) for  $p, q \in (0, 1)$  and (21), result

$$(22) \quad f(0, 0, \frac{1}{2}) = 1.$$

Now (22) in (19) with  $r = 0$  gives,

$$(23) \quad f(p, q, 0) = [(1-p)^\alpha (1-q)^{\beta-1} - 1] (2^{\beta-1} - 1)^{-1}, \quad \text{for all } p, q \in [0, 1).$$

Taking  $\xi = 1, q = 0$  and  $r = 0$  in (11), (11) becomes,

$$(24) \quad \begin{aligned} & f(0, 1 - \eta, 1 - \zeta) + \eta^{\beta-1} \zeta^{1-\beta} f(p, 0, 0) = \\ & = f(p, 0, 0) + (1-p)^\alpha f(0, 1 - \eta, 1 - \zeta), \end{aligned}$$

where  $p \in [0, 1)$ , and  $\eta, \zeta \in (0, 1]$ .

In view of (23), (24) gives

$$(25) \quad f(0, 1 - \eta, 1 - \zeta) = (\eta^{\beta-1} \zeta^{1-\beta} - 1) (2^{\beta-1} - 1)^{-1}, \quad \text{for } \eta, \zeta \in (0, 1].$$

Letting  $p = 1, \eta = 1$  and  $\zeta = 1$  in (11), we get

$$(26) \quad \begin{aligned} & f(1 - \xi, 0, 0) + \xi^\alpha f(1, q, r) = \\ & = f(\xi, q, r) + (1 - \xi)^\alpha (1 - q)^{\beta-1} (1 - r)^{1-\beta} f(1, 0, 0), \end{aligned}$$

for  $\xi \in (0, 1), q, r \in [0, 1)$ .

By (23) and (26) with  $q = 0$  and  $r = 0$ , we get

$$(27) \quad f(1, 0, 0) = (1 - 2^{\beta-1})^{-1}.$$

Again from (26), using (18), (23) and (27), we obtain

$$(28) \quad f(1, q, r) = (q^{\beta-1}r^{1-\beta} - 1)(2^{\beta-1} - 1)^{-1}, \quad \text{for } q, r \in (0, 1).$$

$q = 0$ ,  $\xi = 1$  and  $\zeta = 1$  in (11) give,

$$(29) \quad \begin{aligned} f(0, 1 - \eta, 0) + \eta^{\beta-1} f(p, 0, r) &= \\ = f(p, 0, r) + (1 - p)^\alpha (1 - r)^{1-\beta} f(0, 1 - \eta, 0), \end{aligned}$$

where  $\eta \in (0, 1]$  and  $p, r \in [0, 1)$ .

With the help of (25) and (29), we have

$$(30) \quad f(p, 0, r) = [(1 - p)^\alpha (1 - r)^{1-\beta} - 1](2^{\beta-1} - 1)^{-1}, \quad \text{for } p, r \in [0, 1).$$

For  $q = 1$ ,  $\xi = 1$  and  $\zeta = 1$ , (11) gives

$$(31) \quad \begin{aligned} f(0, 1 - \eta, 0) + \eta^{\beta-1} f(p, 1, r) &= \\ = f(p, \eta, r) + (1 - p)^\alpha (1 - \eta)^{\beta-1} (1 - r)^{1-\beta} f(0, 1, 0), \end{aligned}$$

for  $\eta \in (0, 1)$  and  $p, r \in [0, 1)$ .

From (25) and (31) with  $p = 0$ ,  $r = 0$ , we get

$$(32) \quad f(0, 1, 0) = (1 - 2^{\beta-1})^{-1}.$$

Now (31), (32), (25) and (18) yield

$$(33) \quad f(p, 1, r) = (p^\alpha r^{1-\beta} - 1)(2^{\beta-1} - 1)^{-1}, \quad \text{for } p, r \in (0, 1).$$

With  $p = 1$ , (11) becomes

$$(34) \quad \begin{aligned} f(1 - \xi, 1 - \eta, 1 - \zeta) + \xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} f(1, q, r) &= \\ = f(\xi, q\eta, r\zeta) + (1 - \xi)^\alpha (1 - q\eta)^{\beta-1} (1 - r\zeta)^{1-\beta} f\left(1, \frac{1 - \eta}{1 - q\eta}, \frac{1 - \zeta}{1 - r\zeta}\right), \end{aligned}$$

where  $q, r \in I$ ,  $\xi \in (0, 1)$ ,  $\eta, \zeta \in (0, 1]$  such that  $q\eta \neq 1$  and  $r\zeta \neq 1$ .

Now  $r = 1$  and  $\eta = 1$  in (34) imply

$$(35) \quad \begin{aligned} f(1 - \xi, 0, 1 - \zeta) + \xi^\alpha \zeta^{1-\beta} f(1, q, 1) &= \\ = f(\xi, q, \zeta) + (1 - \xi)^\alpha (1 - q)^{\beta-1} (1 - \zeta)^{1-\beta} f(1, 0, 1), \end{aligned}$$

for  $q \in [0, 1)$ ,  $\xi, \zeta \in (0, 1)$ .

Consequently (30) and (35) with  $q = 0$ , give

$$(36) \quad f(1, 0, 1) = (1 - 2^{\beta-1})^{-1}.$$

Employing (30), (18), (35) and (36), we obtain

$$(37) \quad f(1, q, 1) = (q^{\beta-1} - 1)(2^{\beta-1} - 1)^{-1}, \quad \text{for } q \in [0, 1].$$

Taking  $\eta = 1$  and  $r = 0$ , (34) gives

$$(38) \quad \begin{aligned} f(1 - \xi, 0, 1 - \zeta) + \xi^\alpha \zeta^{1-\beta} f(1, q, 0) = \\ = f(\xi, q, 0) + (1 - \xi)^\alpha (1 - q)^{\beta-1} f(1, 0, 1 - \zeta), \end{aligned}$$

for  $\xi \in (0, 1)$ ,  $\zeta \in (0, 1]$  and  $q \in [0, 1)$ .

In view of (23), (27) and (38) with  $\zeta = 1$ , we get

$$(39) \quad f(1, q, 0) = (1 - 2^{\beta-1})^{-1}, \quad \text{for } q \in [0, 1).$$

Again from (23), (30), (38) and (39), we obtain

$$(40) \quad f(1, 0, 1 - \zeta) = (1 - 2^{\beta-1})^{-1}, \quad \text{for } \zeta \in (0, 1].$$

Putting  $q \approx 1$ , (34) gives

$$(41) \quad \begin{aligned} f(1 - \xi, 1 - \eta, 1 - \zeta) + \xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} f(1, 1, r) = \\ = f(\xi, \eta, r\zeta) + (1 - \xi)^\alpha (1 - \eta)^{\beta-1} (1 - r\zeta)^{1-\beta} f\left(1, 1, \frac{1 - \zeta}{1 - r\zeta}\right), \end{aligned}$$

where  $\xi, \eta \in (0, 1)$ ,  $\zeta \in (0, 1]$ ,  $r \in I$  with  $\zeta r \neq 1$ .

Combining (23) and (41) with  $\zeta = 1$  and  $r = 0$ , we have

$$(42) \quad f(1, 1, 0) = (1 - 2^{\beta-1})^{-1}.$$

Again from (23), (42), (18) and (41) with  $\zeta = 1$ , we get

$$(43) \quad f(1, 1, r) = (2^{\beta-1} - 1)^{-1} (r^{1-\beta} - 1), \quad \text{for } r \in (0, 1).$$

Taking  $r \approx 1$ , (11) becomes

$$(44) \quad \begin{aligned} f(1 - \xi, 1 - \eta, 1 - \zeta) + \xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} f(p, q, 1) = \\ = f(p\xi, q\eta, \zeta) + (1 - p\xi)^\alpha (1 - q\eta)^{\beta-1} (1 - \zeta)^{1-\beta} f\left(\frac{1 - \xi}{1 - p\xi}, \frac{1 - \eta}{1 - q\eta}, 1\right), \end{aligned}$$

for  $\xi, \eta \in (0, 1]$ ,  $\zeta \in (0, 1)$ ,  $p, q \in I$  such that  $p\xi \neq 1$  and  $q\eta \neq 1$ .



Letting  $\eta = 1$  and  $\zeta = 1$  in (44), we have

$$(45) \quad \begin{aligned} & f(0, 0, 1 - \zeta) + \zeta^{1-\beta} f(p, q, 1) = \\ & = f(p, q, \zeta) + (1 - p)^\alpha (1 - q)^{\beta-1} (1 - \zeta)^{1-\beta} f(0, 0, 1), \end{aligned}$$

where  $p, q \in [0, 1)$  and  $\zeta \in (0, 1)$ .

From (25) with  $\eta = 1$  and (45) with  $p = q = 0$ , we have

$$(46) \quad f(0, 0, 1) = (1 - 2^{\beta-1})^{-1}.$$

With the help of (22), (46) and (18), (19) with  $r = 1$  and  $p, q \in (0, 1)$  gives

$$(47) \quad f(p, q, 1) = (p^\alpha q^{\beta-1} - 1) (2^{\beta-1} - 1)^{-1}, \quad \text{where } p, q \in (0, 1).$$

Again from (45) with  $q = 0$ , (30) and (46), we get

$$(48) \quad f(p, 0, 1) = (1 - 2^{\beta-1})^{-1}, \quad \text{for } p \in (0, 1).$$

Setting  $p = 0$  and  $\eta = 1$  in (44), we have

$$\begin{aligned} & f(1 - \xi, 0, 1 - \zeta) + \xi^\alpha \zeta^{1-\beta} f(0, q, 1) = \\ & = f(0, q, \zeta) + (1 - q)^{\beta-1} (1 - \zeta)^{1-\beta} f(1 - \xi, 0, 1) \end{aligned}$$

for  $\xi \in (0, 1]$ ,  $\zeta \in (0, 1)$ ,  $q \in [0, 1)$ , which with the help of (30), (25) and (48) gives

$$(49) \quad f(0, q, 1) = (1 - 2^{\beta-1})^{-1}, \quad \text{for } q \in (0, 1).$$

For  $q = 1$  (44) becomes

$$(50) \quad \begin{aligned} & f(1 - \xi, 1 - \eta, 1 - \zeta) + \xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} f(p, 1, 1) = \\ & = f(p\xi, \eta, \zeta) + (1 - p\xi)^\alpha (1 - \eta)^{\beta-1} (1 - \zeta)^{1-\beta} f\left(\frac{1 - \xi}{1 - p\xi}, 1, 1\right), \end{aligned}$$

where  $\xi \in (0, 1]$ ,  $\eta, \zeta \in (0, 1)$  and  $p \in I$  such that  $p\xi \neq 1$ .

$\xi = 1$ ,  $p = 0$  in (50) together with (25) gives

$$(51) \quad f(0, 1, 1) = (1 - 2^{\beta-1})^{-1}.$$

Utilizing (46) with  $\xi = 1$ , (47), (14) and (25) we have

$$(52) \quad f(p, 1, 1) = (p^\alpha - 1) (2^{\beta-1} - 1)^{-1}, \quad \text{for } p \in (0, 1).$$

For  $q = 1$  (11) gives

$$(53) \quad f(1 - \xi, 1 - \eta, 1 - \zeta) + \xi^\alpha \eta^{\beta-1} \zeta^{1-\beta} f(p, 1, r) = \\ = f(p\xi, \eta, r\zeta) + (1 - p\xi)^\alpha (1 - \eta)^{\beta-1} (1 - r\zeta)^{1-\beta} f\left(\frac{1 - \xi}{1 - p\xi}, 1, \frac{1 - \zeta}{1 - r\zeta}\right),$$

for  $\xi, \zeta \in (0, 1]$ ,  $\eta \in (0, 1)$ ,  $p, r \in I$  such that  $p\xi \neq 1$  and  $r\zeta \neq 1$ .

Using (53) with  $\xi = 1$ ,  $\zeta = 1$  and  $r = 0$ , (32), (25) and (23), we get

$$(54) \quad f(p, 1, 0) = (1 - 2^{\beta-1})^{-1}, \quad \text{for } p \in (0, 1).$$

Finally from (53) with  $p = 0$ ,  $\xi = 1$  and  $\zeta = 1$ , (23), (25) and (32) we have

$$(55) \quad f(0, 1, r) = (1 - 2^{\beta-1})^{-1}, \quad \text{for } r \in (0, 1).$$

Combining (5), (23), (25), (27), (28), (30), (32), (33), (36), (37), (39), (40), (42), (43), (46), (47), (48), (49), (51), (52), (54) and (55), we conclude that  $f$  has the form given by (8) (with the notation  $0^\alpha = 0$ ,  $\alpha \neq 0$ ) and that (10) is true for all  $x, y, z \in I$ .

The converse can be easily verified by straight forward calculation.

**Theorem 2.** *The generalized directed-divergence of order  $\alpha (\neq 1)$  and type  $\beta (\neq 1)$  is given by*

$$(56) \quad I_{\alpha, n}^\beta \begin{pmatrix} p_1, \dots, p_n \\ q_1, \dots, q_n \\ r_1, \dots, r_n \end{pmatrix} = (2^{\beta-1} - 1)^{-1} \left[ \sum_{i=1}^n p_i^\alpha q_i^{\beta-1} r_i^{1-\beta} - 1 \right].$$

*Proof.* Equations (7) and (8) give,

$$I_{\alpha, n}^\beta \begin{pmatrix} p_1, \dots, p_n \\ q_1, \dots, q_n \\ r_1, \dots, r_n \end{pmatrix} = \\ = (2^{\beta-1} - 1)^{-1} \sum_{i=2}^n [p_i^\alpha q_i^{\beta-1} r_i^{1-\beta} + P_{i-1}^\alpha Q_{i-1}^{\beta-1} R_{i-1}^{1-\beta} - P_i^\alpha Q_i^{\beta-1} R_i^{1-\beta}] = \\ = (2^{\beta-1} - 1)^{-1} \left[ \sum_{i=2}^n p_i^\alpha q_i^{\beta-1} r_i^{1-\beta} + P_1^\alpha Q_1^{\beta-1} R_1^{1-\beta} - P_n^\alpha Q_n^{\beta-1} R_n^{1-\beta} \right] = \\ = (2^{\beta-1} - 1)^{-1} \left[ \sum_{i=1}^n p_i^\alpha q_i^{\beta-1} r_i^{1-\beta} - 1 \right].$$

This proves theorem 2.

### 3. SPECIAL CASES

An interesting special case of (56) is obtained for  $\alpha = \beta$  and  $Q \equiv R$ . In this case, (56) reduces to the entropy of type  $\beta$ ,

$$(58) \quad I(P) = (2^{\beta-1} - 1)^{-1} \left( \sum_{i=1}^n p_i^{\beta} - 1 \right), \quad \beta \neq 1.$$

which was discussed in [2], [3] and [8] respectively.

Another interesting special case of (56) is obtained by replacing  $\beta$  by  $2 - \beta$  and then taking  $\alpha = \beta$ ,  $P \equiv Q$ ; and is given by

$$(59) \quad I_{\beta,n}^{2-\beta} \left( \begin{matrix} p_1, \dots, p_n \\ p_1, \dots, p_n \\ r_1, \dots, r_n \end{matrix} \right) = (2^{1-\beta} - 1)^{-1} \left( \sum_{i=1}^n p_i r_i^{\beta-1} - 1 \right), \quad \beta \neq 1.$$

The quantity given in (59) is the inaccuracy of type  $\beta$  discussed earlier by us in [6].

#### References

- [1] *Aczel J. and Nath P.*: Axiomatic characterizations of some measures of divergence in information, *Z. Wahr. Verw. Greb.* 21 (1972), 215—224.
- [2] *Daróczy Z.*: Generalized information functions, *Information and Control*, 16 (1970), 36—51.
- [3] *Havrdá J. and Charvát F.*: Quantification method of classification processes, the concept of structural  $a$ -entropy, *Kybernetika*, 3 (1967), 30—35.
- [4] *Kullback S.*: Information theory and statistics, John Wiley and Sons, New York, 1959.
- [5] *Nath P.*: On the measures of errors in information, *J. Math. Sci., Delhi* 3, H6 (1970).
- [6] *Rathie P. N. and Kannappan Pl.*: An inaccuracy function of type  $\beta$ , *Ann. Inst. Statist. Math.*, 25 (1973), 205—214.
- [7] *Kannappan Pl. and Rathie P. N.*: An application of functional equation to information theory, *Ann. Polon. Math.* 26 (1972), 95—101.
- [8] *Vajda I.*: Axioms of  $a$ -entropy of generalized probability distribution, *Kybernetika*, 4 (1968), 105—110.

*Authors' address:* Faculty of Mathematics, University of Waterloo, Waterloo, Canada.