

Jarolím Bureš

Free constructions of 2-structures

Czechoslovak Mathematical Journal, Vol. 26 (1976), No. 1, 20–29

Persistent URL: <http://dml.cz/dmlcz/101369>

Terms of use:

© Institute of Mathematics AS CR, 1976

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

FREE CONSTRUCTIONS OF 2-STRUCTURES

JAROLÍM BUREŠ, Praha

(Received November 8, 1973)

Among the geometric structures generalizing affine and projective planes there are also 2-structures introduced by H. KARZEL in [1]. A detailed classification of these structures was presented by V. HAVEL in the mimeographed text [2]. Simultaneously, he rose the question if the 2-structures of all the types mentioned in [2] exist. This paper contains construction of 2-structures of all such types using free extensions of incidence structures.

My thanks go to V. Havel for his valuable advice and remarks.

1. PRELIMINARIES

Definition 1 ([1] p. 192). A regular incidence structure $(\mathcal{P}, \mathcal{L}, \in)$ is called a 2-structure if there is a decomposition $\mathcal{L} = \mathcal{L}_0 \cup \mathcal{L}_1 \cup \mathcal{L}_2$ of the set of lines \mathcal{L} into disjoint non-empty sets, such that the following conditions are satisfied:

- (1) $\forall A, B \in \mathcal{P}, A \neq B \exists! g \in \mathcal{L} (A \in g \wedge B \in g)$;
- (2) a) $\forall i \in \{1, 2\} \forall g, h \in \mathcal{L}_i (g = h \vee g \cap h = \emptyset)$;
- b) $\forall A \in \mathcal{P}, \forall i \in \{1, 2\} \exists! g_i \in \mathcal{L}_i (A \in g_i)$;
- (3) $\forall i \in \{1, 2\} \forall g \in \mathcal{L}_i, h \in \mathcal{L} \setminus \mathcal{L}_i (g \cap h \neq \emptyset)$.

Every affine plane is a 2-structure if we take for \mathcal{L}_1 and \mathcal{L}_2 two distinct classes of parallel lines and set $\mathcal{L}_0 = \mathcal{L} \setminus (\mathcal{L}_1 \cup \mathcal{L}_2)$. Hence we conclude.

Theorem 1 ([1] p. 193). *Every finite 2-structure is an affine plane.*

Definition 2. A 2-structure $(\mathcal{P}, \mathcal{L}, \in)$ is called a *weak affine plane* if there is an equivalence relation \parallel on \mathcal{L} called “parallelism” so that

- (4) $g \parallel h \Rightarrow g = h \vee g \cap h = \emptyset$;
- (5) $\forall P \in \mathcal{P} \forall g \in \mathcal{L} \exists! h \in \mathcal{L} (P \in h \wedge h \parallel g)$.

Definition 3 ([1] p. 198). Let M be a set, $\#M \geq 2$, G a subset of the group S_M of all permutations of M with the properties

$$(6) \forall a, a', b, b' \in M, a \neq b, a' \neq b' \exists! \gamma \in G (a^\gamma = a' \wedge b^\gamma = b');$$

$$(7) \text{Id} \in G.$$

The pair (M, G) is called a *sharply doubly transitive set of permutations* (with identity).

Denote a 2-structure $(\mathcal{P}, \mathcal{L}, \in)$ with the base line e by $(\mathcal{P}, \mathcal{L}, \in, e)$. Further, denote the line from \mathcal{L}_i going through the point A by $(A \rightarrow i)$ for $i = 1, 2$.

The relation between 2-structures with the base line and sharply doubly transitive sets of permutations is expressed in the following theorem.

Theorem 2 ([1] p. 198). a) If $(\mathcal{P}, \mathcal{L}, \in, e)$ is a 2-structure with the base line e , put $M = \mathcal{L}_2$ and $G = \{\tilde{g} \mid g \in \mathcal{L}_0\}$ the set of all maps of M onto M defined by

$$x^{\tilde{g}} = (((x \cap g) \rightarrow 1) \cap e) \rightarrow 2).$$

Then (M, G) is a *sharply doubly transitive set of permutations*.

b) If (M, G) is a *sharply doubly transitive set of permutations*, put $\mathcal{P} = M \times M$, $\hat{\gamma} = \{(x, x^\gamma) \mid x \in M$,

$$\langle a \rangle_1 = \{(a, y) \mid y \in M\}, \quad \langle a \rangle_2 = \{(x, a) \mid x \in M\},$$

$$\mathcal{L}_0 = \{\hat{\gamma} \mid \gamma \in G\}, \quad \mathcal{L}_1 = \{\langle a \rangle_1 \mid a \in M\}, \quad \mathcal{L}_2 = \{\langle a \rangle_2 \mid a \in M\}.$$

Then $(\mathcal{P}, \mathcal{L}, \in, \hat{\text{Id}})$ is a 2-structure with the base line $\hat{\text{Id}}$ (see Fig. 1).




Figure 1.

Theorem 3 ([3] p. 292). Let (M, G) be a *sharply doubly transitive set of permutations*. The corresponding 2-structure is an *affine plane* if and only if

$$(8) \forall a, b \in M \forall \gamma \in G a^\gamma \neq b \exists! \delta \in G [a^\delta = b \wedge (\forall x \in M x^\delta \neq x^\gamma)].$$

Definition 4. A permutation π of M is called *dispersive* if it satisfies

$$\forall x \in M \quad (x^\pi \neq x).$$

Denote by \mathcal{R}_M the set of all dispersive permutations from mS_M and for $G \subseteq S_M$ put $G' = G \cap \mathcal{R}_M$.

If M is finite, then (6) \Rightarrow (8) and Theorem 3 implies Theorem 1. For the proof see e.g. [2]. Generally (in the infinite case) the implication is not true.

A counter-example (by M. Hall). Denote by N the set of all non-negative integers, and put $N_k = \{0, 1, \dots, k\}$ for $k \in N$. A partial permutation of N is defined either as a permutation of N or as a bijection π from a finite subset $D_\pi \subseteq N$ onto $R_\pi \subseteq N$. We say that a partial permutation π has a height k if

$$N_k \subseteq D_\pi \cap R_\pi.$$

A finite set of partial permutations $S = \{\gamma_1, \gamma_2, \dots, \gamma_l\}$ is admissible of order k if

(9) there exists at most one $\gamma \in S$ such that $a^\gamma = c$, $b^\gamma = d$ for every $a, b, c, d \in N$, $a \neq b$, $c \neq d$.

(10) $\forall a, b, c, d \in N$, $a \neq b$, $c \neq d$, $a + b + c + d \leq k$ $\exists!$ $\gamma \in S$ ($a^\gamma = c \wedge b^\gamma = d$).

(11) Every $\gamma \in S$ has the height k .

It is obvious that for fixed k we can construct many admissible sets of partial permutations of order k .

To every admissible set $S = \{\gamma_1, \dots, \gamma_l\}$ of order k we can construct an admissible set \tilde{S} of order $k + 1$ in two steps:

I. Denote by T the set of all elements $(a, b, c, d) \in N \times N \times N \times N$ with the properties

(i) $a \neq b$, $c \neq d$, $a + b + c + d = k + 1$.

(ii) There exists no $\gamma \in S$ such that $a^\gamma = c$ and $b^\gamma = d$. Arrange the elements of T into the finite sequence:

$$(a_1, b_1, c_1, d_1), (a_2, b_2, c_2, d_2), \dots, (a_m, b_m, c_m, d_m)$$

and define partial permutations $\gamma_{l+1}, \dots, \gamma_{l+m}$ by

$$D_{\gamma_{l+i}} = \{a_i, b_i\}, \quad R_{\gamma_{l+i}} = \{c_i, d_i\}, \quad a_i^{\gamma_{l+i}} = c_i, \quad b_i^{\gamma_{l+i}} = d_i.$$

We obtain the set of partial permutations $S' = \{\gamma_1, \dots, \gamma_l, \gamma_{l+1}, \dots, \gamma_{l+m}\}$ which satisfies the conditions (9) and (10) but not (11).

II. Assign every partial permutation $\gamma_i \in S'$ a permutation $\tilde{\gamma}_i \in \tilde{S}$ which extends γ_i . Now we construct successively the permutations $\tilde{\gamma}_1, \dots, \tilde{\gamma}_{m+l}$.

Construction of $\tilde{\gamma}_1$: Let $(N \setminus D_{\gamma_1}) \cap N_{k+1} = \{t_1, \dots, t_r\}$, $t_1 < t_2 < \dots < t_r$ and $(N \setminus R_{\gamma_1}) \cap N_{k+1} = \{s_1, \dots, s_u\}$, $s_1 < s_2 < \dots < s_u$. We construct successively $t_1^{\tilde{\gamma}_1}, t_2^{\tilde{\gamma}_1}, \dots, t_r^{\tilde{\gamma}_1}$ in such a way that at each step after the corresponding extension of γ_1 (9) is satisfied for the system S' with the extended γ_1 instead of the original γ_1 .

Further, we choose successively v_1, \dots, v_u and set $v_p^{\tilde{\gamma}_1} = s_p$ in such a way that after each step the condition (9) is fulfilled. For $\tilde{\gamma}_1$ we have:

$$D_{\tilde{\gamma}_1} = D_{\gamma_1} \cup \{t_1, \dots, t_r\} \cup \{v_1, \dots, v_u\},$$

$$R_{\tilde{\gamma}_1} = R_{\gamma_1} \cup \{t_1^{\tilde{\gamma}_1}, \dots, t_r^{\tilde{\gamma}_1}\} \cup \{s_1, \dots, s_u\}, \quad \tilde{\gamma}_1 = \gamma_1 \text{ on } D_{\gamma_1}.$$

This construction can be applied successively to the systems

$$S'_2 = \{\gamma_2, \dots, \gamma_{m+l}, \tilde{\gamma}_1\}, \dots, S'_{m+l} = \{\gamma_{m+l}, \hat{\gamma}_1, \dots, \gamma'_{m+l-1}\}$$

and after $m+l$ steps we obtain the desired \tilde{S} .

To every admissible set S_k of order k we can construct a sequence of admissible sets $\{S_{k+i}\}_{i=0}^{\infty}$ satisfying:

- (i) order of $S_{k+i} \geq k+i$.
- (ii) There exists an injection $I_p^q : S_{k+p} \rightarrow S_{k+q}$, $p < q$ such that $I_p^q(\gamma)$ is an extension of γ for every γ .

The system $\{S_{k+i}, I_p^q\}$ is an inductive system of sets and the inductive limit $\varinjlim S_{k+i} = S_{\infty}$ exists. An element from S_{∞} is a permutation of N and the set S_{∞} satisfies the conditions (9) and (10), i.e., S_{∞} is a sharply doubly transitive set of permutations of N .

Special examples. A. If we take $S_2 = \{\gamma_1, \gamma_2, \gamma_3\}$, $\gamma_1 = \text{Id}$,

$$0^{\gamma_2} = 1, \quad 1^{\gamma_2} = 2, \quad 2^{\gamma_2} = 0 \quad (2r-1)^{\gamma_2} = 2r, \quad (2r)^{\gamma_2} = 2r-1,$$

$$(2r+1)^{\gamma_3} = 2r, \quad (2r)^{\gamma_3} = 2r+1$$

so we have

$$\gamma_1, \gamma_2, \gamma_3 \in S_{\infty}, \quad 0^{\gamma_2} = 1, \quad 0^{\gamma_3} = 1, \quad \gamma_2 \gamma_1^{-1} \in \mathcal{R}, \quad \gamma_3 \gamma_1^{-1} \in \mathcal{R}.$$

B. We take $S = \{\gamma_1, \gamma_2\}$, $\gamma_1 = \text{Id}$,

$$D_{\gamma_2} = \{0, 1, 2\}, \quad 0^{\gamma_2} = 1, \quad 1^{\gamma_2} = 0, \quad 2^{\gamma_2} = 2$$

and complete the construction in the following way: To every permutation γ we add one fixed element, i.e. $x \in N$ with $x^{\gamma} = x$ if there is no one left. There exists no permutation $\gamma \in S_{\infty}$ satisfying $\gamma \gamma_1^{-1} \in \mathcal{R}$.

The examples A and B determine 2-structures which are not affine planes.

Remark. As we extend only finite partial permutations and we have only a finite system of permutations we can always choose the elements of N in the step II so that (9) and (10) are satisfied.

2. CONSTRUCTION OF 2-STRUCTURES

Definition 5. Let (M, G) be a sharply doubly transitive set of permutations. The triple $(a, b, \gamma) \in M \times M \times G$ is called *admissible* ([2] p. 7) if $a^\gamma \neq b$.

Denote by \mathfrak{M} the set of all admissible triples of (M, G) . For $(a, b, \gamma) \in \mathfrak{M}$ define the multiplicity $N(a, b, \gamma)$:

$$\begin{aligned} N(a, b, \gamma) &= 0 \text{ if there is no } \delta \in G \text{ such that } a^\delta = b, \gamma\delta^{-1} \in \mathcal{R}_M; \\ N(a, b, \gamma) &= 1 \text{ if there is only one } \delta \in G \text{ such that } a^\delta = b, \gamma\delta^{-1} \in \mathcal{R}_M; \\ N(a, b, \gamma) &= 2 \text{ if there exist } \delta_1, \delta_2 \in G, \text{ such that } \delta_1 \neq \delta_2, a^{\delta_i} = b, \gamma\delta_i^{-1} \in \mathcal{R}_M, \\ &\quad i = 1, 2. \end{aligned}$$

We say that: (M, G) has the type (j) for $j \in \{0, 1, 2\}$ if every $(a, b, \gamma) \in \mathfrak{M}$ has the multiplicity (j) ;

(M, G) has the type (i, j) if there exist $(a, b, \gamma) \in \mathfrak{M}$ with the multiplicity i , $(a', b', \gamma') \in \mathfrak{M}$ with the multiplicity j and one with the multiplicity k , $\{i, j, k\} = \{0, 1, 2\}$;

(M, G) has the type $(0, 1, 2)$ if there exist triples with all multiplicities.

The notions of Definition 5 have a geometric meaning for the corresponding 2-structure.

If $(\mathcal{P}, \mathcal{L}, \in)$ is the corresponding 2-structure, then the condition $\gamma\delta^{-1} \in \mathcal{R}_M$ means that the corresponding lines $\tilde{\gamma}$ and $\tilde{\delta}$ have $\tilde{\delta}$ have no intersection point.

Theorem 1 and Theorem 3 imply

Theorem 4. a) (M, G) has the type (1) if and only if the corresponding 2-structure is an affine plane.

b) If M is a finite set, then (M, G) has always the type (1).

Similarly, for a 2-structure $(\mathcal{P}, \mathcal{L}, \in) = \mathcal{S}$ we define a pair $(P, l) \in \mathcal{P} \times \mathcal{L}$ to be admissible if $P \notin l$.

We say that an admissible pair (P, l) has the multiplicity j (writing $\tilde{N}(P, l) = j$) for $j \in \{0, 1, 2\}$ in the following cases:

$$\begin{aligned} \tilde{N}(P, l) &= 0 \Leftrightarrow^{\text{Def}} \forall l_1 \in \mathcal{L} (P \in l_1 \Rightarrow l \cap l_1 = \emptyset), \\ \tilde{N}(P, l) &= 1 \Leftrightarrow^{\text{Def}} \exists! l_1 \in \mathcal{L} (P \in l_1 \wedge l \cap l_1 = \emptyset), \\ \tilde{N}(P, l) &= 2 \Leftrightarrow^{\text{Def}} \exists! l_1, l_2 \in \mathcal{L} \ l_1 \neq l_2 \ \forall i \in \{1, 2\} (P \in l_i \wedge l \cap l_i = \emptyset). \end{aligned}$$

A 2-structure \mathcal{S} has the type (j) if every admissible pair has the multiplicity j , type

(j, k) if there exists an admissible pair of the multiplicity j and an admissible pair of the multiplicity k while no admissible pair has the multiplicity m . (Here $\{j, k, m\} = \{0, 1, 2\}$.)

We say it has the type $(0, 1, 2)$ if there exist admissible pairs with all multiplicities.

Theorem 5. a) (M, G) has the type α if and only if the corresponding 2-structure has the same type.

$$(\alpha \in \{(0), (1), (2), (0, 1), (0, 2), (1, 2), (0, 1, 2)\}.)$$

b) If $(\mathcal{P}, \mathcal{L}, \epsilon)$ is a 2-structure of the type 0 then every two distinct lines from \mathcal{L}_0 have an intersection point.

The proof is obvious.

The main result of this paper is

Theorem 6. There exist 2-structures of all types.

We shall construct 2-structures of all types except the type (1) (this type is that of every affine plane). All 2-structures constructed in the sequel are not affine planes. For better description of the construction we take two singular points $\alpha \notin \mathcal{P}$, $\beta \notin \mathcal{P}$ and suppose that α lies exactly on all lines of \mathcal{L}_1 , β lies exactly on the lines of \mathcal{L}_2 and there is no line containing both α and β .

Let K be an arbitrary set, $\#K \geq 2$. Put

$$\mathcal{P}^0 = K \times K \cup \{\alpha, \beta\}, \quad \mathcal{L}^0 = \mathcal{L}_1^0 \cup \mathcal{L}_2^0 \cup \mathcal{L}_0^0, \quad \mathcal{I}^0 = (\mathcal{P}^0, \mathcal{L}^0, \epsilon),$$

\mathcal{L}_1^0 is formed by all subsets of \mathcal{P}^0 of the form

$$\{(k, x) \mid x \in K\} \cup \{\alpha\} \quad \forall k \in K,$$

\mathcal{L}_2^0 is formed by all subsets of \mathcal{P}^0 of the form

$$\{(x, k) \mid x \in K\} \cup \{\beta\}$$

and \mathcal{L}_0^0 will be defined in each case separately.

We proceed by induction constructing step by step the incidence structures $\mathcal{I}^0 = (\mathcal{P}^i, \mathcal{L}^i, \epsilon)$ in the following ways.

Case I. $\mathcal{L}_0^0 = \emptyset$ a) \mathcal{L}^i is obtained from \mathcal{L}^{i-1} by adding all lines joining the points which have not been joined so far (i.e. new lines $\{A, B\}$, $A, B \in \mathcal{P}^{i-1}$ such that there exists no line from \mathcal{L}^{i-1} going through A and B) with the exception of the line joining α and β .

b) \mathcal{P}^i is obtained from \mathcal{P}^{i-1} by adding all intersection points of the pairs of lines which have no intersection point in \mathcal{P}^{i-1} (i.e. new distinct points $\{l, m\}$, $l, m \in \mathcal{L}^i$ with $l \cap m = \emptyset$ in \mathcal{P}^{i-1} , we set $\{l, m\} \in l, m$.)

Case II. $\mathcal{L}_0^0 = \emptyset$, $\#K \geq 3$

a) \mathcal{L}^i is obtained as in Ia.

b) \mathcal{P}^i is obtained by adding to \mathcal{P}^{i-1} all intersection points of lines going through α or β with other lines.

Case III. For $\#K \geq 3$, we fix an arbitrary point $A \in K \times K$.

a) \mathcal{L}^i is obtained as in Ia.

b) \mathcal{P}^i is obtained by adding to \mathcal{P}^{i-1} all intersection points of all lines going through α or β with other lines, and all intersections of lines going through A with other lines.

Case IV. For $\#K \geq 3$. Let $\Delta = \{(k, k) \mid k \in K\}$ and let Δ_1 be a subset of $K \times K$ satisfying

(i) $\forall k \in K \exists! x(k, x) \in \Delta_1$,

(ii) $\forall k \in K \exists! y(y, k) \in \Delta_1$,

(iii) $\Delta \cap \Delta_1 = \emptyset$.

Fix $A \in \Delta_1$ and put $\mathcal{L}_0^0 \in \{\Delta, \Delta_1\}$.

a) \mathcal{L}_i is obtained as in Ia.

b) \mathcal{P}^i is obtained from \mathcal{P}^{i-1} by adding the intersection points of all lines from \mathcal{L}^i with the lines going through α or β and the intersection points of all lines, with the exception of Δ_1 , going through A with the line Δ .

Case V. For $\#K \geq 3$. Choose Δ, Δ_1 and set $\mathcal{L}_0^0 = \{\Delta, \Delta_1\}$ as in IV.

a) \mathcal{L}^i is obtained as in Ia.

b) \mathcal{P}^i is obtained from \mathcal{P}^{i-1} by adding the intersection points of all lines, with the only exception of the intersection point of Δ with Δ_1 .

Case VI. For $\#K \geq 3$. Choose Δ as in IV and take $A \notin \Delta$. Set $\mathcal{L}_0^0 = \{\Delta\}$.

a) \mathcal{L}^i is obtained as in Ia.

b) \mathcal{P}^i is obtained from \mathcal{P}^{i-1} by adding the intersection points of all lines from $\mathcal{L}^i \setminus \{\Delta\}$ and further by adding the intersection points of Δ with the lines going through α or β .

A modification of VI is obtained from VI if we substitute b) by b)':

b)' \mathcal{P}^i is obtained from \mathcal{P}^{i-1} by adding the following intersection points: Intersection points of any line with the lines going through α or β , and intersection points of the lines going through A with all lines from $\mathcal{L}^i \setminus \{\Delta\}$.

In all these cases we add intersection points only if the corresponding lines do not intersect in \mathcal{P}^{i-1} , and we set

$$\mathcal{P} = \bigcup_{i=0}^{\infty} \mathcal{P}^i \setminus \{\alpha, \beta\}, \quad \mathcal{L} = \bigcup_{i=1}^{\infty} \mathcal{L}^i, \quad \mathcal{L}_1 = \{g \in \mathcal{L} \mid \alpha \in g\}, \quad \mathcal{L}_2 = \{g \in \mathcal{L} \mid \beta \in g\}.$$

Theorem 7. *The incidence structures constructed in I–VI are 2-structures which are not affine planes.*

They are of the following types: In the case I of the type (0), in the case II of the type (2), in the case III of the type (0, 2), in the case IV of the type (0, 1), in the case V of the type (1, 2) and in the case VI of the type (0, 1, 2)

Proof. From the part (a) of the constructions it follows that any two points can be joined by a unique line and to each point of \mathcal{P} there exists a unique line from \mathcal{L}_1 and a unique line from \mathcal{L}_2 going through the point. As a consequence of the part b) we get that every line from \mathcal{L}_i intersects every line from $\mathcal{L} \setminus \mathcal{L}_i$ ($i = 1, 2$).

Distinct lines from \mathcal{L}_1 or \mathcal{L}_2 have no intersection point. Moreover, in the individual cases we have:

I. Every pair of lines from \mathcal{L}_0 has an intersection point, every admissible pair has the multiplicity 0 and the 2-structure is of the type (0).

II. Let $l, m \in \mathcal{L}_0$ be arbitrary lines. Then there exists an index k so that $l, m \in \mathcal{L}^k$. If l does not intersect m at a point from \mathcal{P}^k , then l does not intersect m .

If (A, l) is an admissible pair, then there exists k such that $A \in \mathcal{P}^k$, $l \in \mathcal{L}^k$. Let us choose points $A_1, A_2 \in \mathcal{P}^{k+1}$, A_1, A_2, A non-collinear, and $A_1, A_2 \notin l$ (e.g., newly added intersection points). Then AA_1 and AA_2 do not intersect l . The 2-structure is of the type (2).

III. If $B \in \mathcal{P}$, $B \neq A$ and $l \in \mathcal{L}_0$ is a line not going through A , (B, l) is an admissible pair, then (B, l) has the multiplicity 2 by virtue of II. If $m \in \mathcal{L}_0$ is a line going through A and $B \notin m$ an arbitrary point, then every line going through B intersects m . Similarly if (A, t) is an admissible pair then every line going through A_0 intersects t . The pairs $\{(A, l) \mid A \notin l\}$, $\{(B, m) \mid B \notin m, A \in m\}$ are of the multiplicity 0, the 2-structure is thus of the type (0, 2).

IV. If $B \in \mathcal{P}$, $B \neq A$ and (B, l) is an admissible pair then there exist at least two lines going through B and not intersecting l . The pair (B, l) has the multiplicity 2. Similarly, the pair (A, l) for $l \neq \Delta$ has the multiplicity 2. The pair (A, Δ) has the multiplicity 1. The 2-structure is of the type (1, 2).

V. Through every point $A \in \Delta$ there goes a unique line non-intersecting Δ_1 , namely Δ . Similarly, through every point of Δ_1 there goes a unique line non-intersecting Δ , namely Δ_1 . The admissible pairs $\{(A, \Delta_1) \mid A \in \Delta\}$, $\{(B, \Delta), B \in \Delta_1\}$ have the multiplicity 1. The others have the multiplicity 0 (see I). The 2-structure has the type (0, 1).

VI. Let (B, Δ) be an admissible pair and $B \in \mathcal{P}^k$. If $A_1, A_2 \in \mathcal{P}^k$ are such that B, A_1, A_2 are non-collinear and BA_1, BA_2 do not intersect Δ in \mathcal{P}^k then the lines BA_1, BA_2 do not intersect Δ even in \mathcal{P} , and the pair (B, Δ) has the multiplicity 2. Choose now $X \in \Delta$ and a line m non-intersecting Δ . (Such a line surely exists; it is e.g. the line joining points not lying on Δ in \mathcal{L}^i). Then there is a unique line going

through X and non-intersecting m , namely Δ . The pairs (X, Δ) have the multiplicity 1. If l is a line intersecting Δ , then every line going through X intersects l and thus (X, l) has the multiplicity 0. The 2-structure is of the type $(0, 1, 2)$.

VI'. Here we can proceed similarly as in VI. Admissible pairs (A, l) , $l \neq \Delta$ has the multiplicity 0, the admissible pair (A, Δ) has the multiplicity 2. The admissible pairs of the type (X, l) for $X \in \Delta$, $A \in l$, and $l \cap \Delta = \emptyset$ have the multiplicity 1, the admissible pairs (B, l) , $A \in l$, $B \notin \Delta$ have the multiplicity 0, the other admissible pairs have the multiplicity 2. The 2-structure has the type $(0, 1, 2)$.

Definition 6. A regular incidence structure $(\mathcal{P}, \mathcal{L}, \in)$ provided with a decomposition $\mathcal{L} = \mathcal{L}_0 \cup \mathcal{L}_1 \cup \mathcal{L}_2$ satisfying (2), (3) from Definition 1 is called a *partial 2-structure*.

Remark. After a modification, the constructions I–VI can be applied to an arbitrary partial 2-structure.

We arrive at various 2-structures also when studying some well-known algebraic structures. This questions will be discussed in the last part of this paper.

Theorem 8. Let $\mathcal{F} = (Q, +, \cdot)$ be a proper reduced quasifield (i.e. there exist $a, b, c \in Q$, $a \neq b$ such that the equation $-a \cdot x + b \cdot x = c$ has no solution). The corresponding incidence structure is a 2-structure (namely, a weak affine plane) which is not an affine plane.

Further, if \mathcal{F} satisfies the condition

$$(12) \quad \exists b', c' \in Q \setminus \{0\} \quad \forall m \in Q \setminus \{b'\} \quad \exists x \in Q \quad (-m \cdot x + b' \cdot x = c')$$

then the 2-structure has the type $(1, 2)$. If (12) is not satisfied the 2-structures has the type (2) .

Proof. The 2-structure $\mathcal{I} = (\mathcal{P}, \mathcal{L}, \in)$ which corresponds to \mathcal{F} has the form

$$\begin{aligned} \mathcal{P} &= Q \times Q, & \mathcal{L}_1 &= \{(a, x) \mid x \in Q \mid a \in Q\}, & \mathcal{L}_2 &= \{(x, a) \mid x \in Q \mid a \in Q\}, \\ \mathcal{L}_0 &= \{(x, y) \mid y = m \cdot x + b \mid x \in Q\}, & m &\neq 0, & m, b &\in Q. \end{aligned}$$

The pair (P, l) , $P = (0, 0)$ and $l = \{(x, ax + c) \mid x \in Q\}$ has the multiplicity 2, because there exist two lines $l_1 = \{(x, ax) \mid x \in Q\}$ and $l_2 = \{(x, bx) \mid x \in Q\}$ non-intersecting l and going through P .

If the condition (12) is satisfied, then the pair (P, q) , $P = (0, 0)$ and $q = \{(x, mx + c) \mid x \in Q\}$ has the multiplicity 1, because every line $q'_1 \neq q_1$, $q_1 = \{(x, mx) \mid x \in Q\}$ which goes through P intersects q . There is no admissible pair of the multiplicity 0.

If the condition (12) is not satisfied then no lines of the form

$$\{(x, bx + c_1) \mid x \in Q\}, \quad \{(x, mx + c_2) \mid x \in Q\}$$

have an intersection point. Every admissible pair has the multiplicity 2 and the 2-structure has the type (2).

If we define the relation \parallel on \mathcal{L} in such a way that the first class of equivalence is the set \mathcal{L}_1 , the second is \mathcal{L}_2 and for $l_1, l_2 \in \mathcal{L}_0$ of the form

$$l_1 = \{(x, rx + t_1) \mid x \in Q\}, \quad l_2 = \{(x, sx + t_2) \mid x \in Q\}$$

we set $l_1 \parallel l_2$ if $r = s$, we obtain the relation which has all the properties from Definition 3.

Theorem 9. *For a proper nonplanar nearfield $\mathcal{F} = (Q, +, \cdot)$ (i.e., there exist $t, s, t \neq 1$ such that the equation $t \cdot x = x + s$ has no solution) the condition (12) is not satisfied. The corresponding 2-structure has the type 2.*

Proof. Theorem 9 follows immediately from Theorem 8 and Theorem 3,2 of [6].

Let $\mathcal{F} = (Q, +, \cdot)$ be a nonplanar quasifield, $\mathcal{A}(\mathcal{F})$ the corresponding 2-structure. If we construct the sharply doubly transitive set of permutations (M, G) which corresponds to $\mathcal{A}(\mathcal{F})$ with the base line $A = \{(x, x) \mid x \in Q\}$ we obtain the permutations from G in the form

$$\gamma \in G \Rightarrow x^\gamma = m \cdot x + b.$$

Then G is a permutation group and $G \cdot$ is a subgroup of G . $G \cdot$ is the set of all elements \bar{b} of the form

$$x^{\bar{b}} = x + b, \quad b \in Q.$$

Remark. There exists a non-planar nearfield (see e.g. [5, [6]]) and a non-planar quasifield (see [4]).

References

- [1] *H. Karzel:* Zusammenhänge zwischen Fastbereichen, scharf zweifach transitiven Permutationsgruppen und 2-Strukturen mit Rechteckaxiom. *Abh. Math. Sem. Univ. Hamburg* 32 (1968), 191–206.
- [2] *V. Havel:* Geometric interpretation of sharply doubly transitive permutation groups (Czech) Mimeographed text, Palacký University Olomouc 1973.
- [3] *M. Hall:* Projective planes, *Trans. Amer. Math. Soc.* 54 (1943), 224–277.
- [4] *J. Bureš:* Construction of an infinite quasifield, *Geometriae dedicata* 1 (1973), 500–504.
- [5] *W. Kerby:* Projektive und nicht-projektive Fastkörper, *Abh. Math. Sem. Univ. Hamburg* 32 (1968), 20–24.
- [6] *E. H. Davis:* Incidence systems associated with non-planar nearfields, *Canad. Math. J.* 22 (1970), 939–953.

Author's address: 118 00 Praha, Malostranské nám. 25, ČSSR (Matematicko-fyzikální fakulta UK).