

Alois Švec

On the rigidity of certain surfaces in E^5

Czechoslovak Mathematical Journal, Vol. 27 (1977), No. 2, 250–257

Persistent URL: <http://dml.cz/dmlcz/101465>

Terms of use:

© Institute of Mathematics AS CR, 1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE RIGIDITY OF CERTAIN SURFACES IN E^5

ALOIS ŠVEC, Olomouc

(Received May 12, 1975)

E. BOMPIANI [1] has presented the classes of surfaces in E^n which may admit non-trivial higher order deformations (at least locally). In E^5 , the general surfaces of this class are those possessing a conjugate net. In what follows, I just show the global infinitesimal rigidity of a subclass of surfaces with a conjugate net. Of course, Bompiani's results deserve a further study.

1. Let $G \subset \mathcal{R}^2$ be a bounded domain, ∂G its boundary and $M : \bar{G} \rightarrow E^5$, $\bar{G} = G \cup \partial G$, a surface in the 5-dimensional Euclidean space. To each point $m \in M \equiv M(\bar{G})$, associate an orthonormal frame $\{v_1, \dots, v_5\}$ such that $v_1, v_2 \in T(M)$. Then

$$(1) \quad \begin{aligned} dm &= \omega^1 v_1 + \omega^2 v_2, \\ dv_1 &= \omega_1^2 v_2 + \omega_1^3 v_3 + \omega_1^4 v_4 + \omega_1^5 v_5, \\ dv_2 &= -\omega_1^2 v_1 + \omega_2^3 v_3 + \omega_2^4 v_4 + \omega_2^5 v_5, \\ dv_3 &= -\omega_1^3 v_1 - \omega_2^3 v_2 + \omega_3^4 v_4 + \omega_3^5 v_5, \\ dv_4 &= -\omega_1^4 v_1 - \omega_2^4 v_2 - \omega_3^4 v_3 + \omega_4^5 v_5, \\ dv_5 &= -\omega_1^5 v_1 - \omega_2^5 v_2 - \omega_3^5 v_3 - \omega_4^5 v_4 \end{aligned}$$

with the usual integrability conditions $d\omega^i = \omega^j \wedge \omega_j^i$, $d\omega_i^j = \omega_i^k \wedge \omega_k^j$. From

$$(2) \quad \omega^3 = \omega^4 = \omega^5 = 0,$$

we get

$$(3) \quad \begin{aligned} \omega^1 \wedge \omega_1^3 + \omega^3 \wedge \omega_2^3 &= 0, \quad \omega^1 \wedge \omega_1^4 + \omega^2 \wedge \omega_2^4 = 0, \\ \omega^1 \wedge \omega_1^5 + \omega^2 \wedge \omega_2^5 &= 0 \end{aligned}$$

and the existence of functions a_1, \dots, c_3 such that

$$(4) \quad \begin{aligned} \omega_1^3 &= a_1\omega^1 + a_2\omega^2, & \omega_2^3 &= a_2\omega^1 + a_3\omega^2, \\ \omega_1^4 &= b_1\omega^1 + b_2\omega^2, & \omega_2^4 &= b_2\omega^1 + b_3\omega^2, \\ \omega_1^5 &= c_1\omega^1 + c_2\omega^2, & \omega_2^5 &= c_2\omega^1 + c_3\omega^2. \end{aligned}$$

Let $T_m^2(M)$ denote the 2-osculating space of M at m , and suppose $\dim T_m^2(M) = 4$ for each $m \in M$. The frames $\{v_i\}$ be chosen in such a way that $v_3, v_4 \in T^2(M)$, i.e.,

$$(5) \quad \omega_1^5 = \omega_2^5 = 0.$$

Let $\{w_i\}$ be another field of moving frames, and let

$$(6) \quad \begin{aligned} v_1 &= \varepsilon_1(\cos \alpha \cdot w_1 - \sin \alpha \cdot w_2), & v_2 &= \sin \alpha \cdot w_1 + \cos \alpha \cdot w_2, \\ v_3 &= \varepsilon_2(\cos \beta \cdot w_3 - \sin \beta \cdot w_4), & v_4 &= \sin \beta \cdot w_3 + \cos \beta \cdot w_4, \\ v_5 &= \varepsilon_3 w_5; & \varepsilon_1^2 &= \varepsilon_2^2 = \varepsilon_3^2 = 1. \end{aligned}$$

The equations (1) being now $dm = \Omega^i w_i$, $dw_i = \Omega_j^i w_j$, we get

$$(7) \quad \begin{aligned} \omega^1 &= \varepsilon_1(\cos \alpha \cdot \Omega^1 - \sin \alpha \cdot \Omega^2), & \omega^2 &= \sin \alpha \cdot \Omega^1 + \cos \alpha \cdot \Omega^2, \\ \omega_1^3 &= \varepsilon_1 \varepsilon_2 (\cos \alpha \cos \beta \cdot \Omega_1^3 - \sin \alpha \cos \beta \cdot \Omega_2^3 - \cos \alpha \sin \beta \cdot \Omega_1^4 + \\ & \quad + \sin \alpha \sin \beta \cdot \Omega_2^4), \\ \omega_1^4 &= \varepsilon_1 (\cos \alpha \sin \beta \cdot \Omega_1^3 - \sin \alpha \sin \beta \cdot \Omega_2^3 + \cos \alpha \cos \beta \cdot \Omega_1^4 - \\ & \quad - \sin \alpha \cos \beta \cdot \Omega_2^4), \\ \omega_2^3 &= \varepsilon_2 (\sin \alpha \cos \beta \cdot \Omega_1^3 + \cos \alpha \cos \beta \cdot \Omega_2^3 - \sin \alpha \sin \beta \cdot \Omega_1^4 - \\ & \quad - \cos \alpha \sin \beta \cdot \Omega_2^4), \\ \omega_2^4 &= \sin \alpha \sin \beta \cdot \Omega_1^3 + \cos \alpha \sin \beta \cdot \Omega_2^3 + \sin \alpha \cos \beta \cdot \Omega_1^4 + \\ & \quad + \cos \alpha \cos \beta \cdot \Omega_2^4. \end{aligned}$$

Consider the functions

$$(8) \quad K = a_1 a_3 - a_2^2 + b_1 b_3 - b_2^2, \quad k = a_1 b_2 - a_2 b_1 + a_2 b_3 - a_3 b_2$$

defined by

$$(9) \quad \omega_1^3 \wedge \omega_2^3 + \omega_1^4 \wedge \omega_2^4 = K \omega^1 \wedge \omega^2, \quad \omega_1^3 \wedge \omega_1^4 + \omega_2^3 \wedge \omega_2^4 = k \omega^1 \wedge \omega^2$$

resp. Similarly, K^* and k^* be defined by

$$(10) \quad \Omega_1^3 \wedge \Omega_2^3 + \Omega_1^4 \wedge \Omega_2^4 = K^* \Omega^1 \wedge \Omega^2, \quad \Omega_1^3 \wedge \Omega_1^4 + \Omega_2^3 \wedge \Omega_2^4 = k^* \Omega^1 \wedge \Omega^2$$

resp. From (7), we get

$$(11) \quad K^* = K, \quad k^* = \varepsilon_1 \varepsilon_2 k.$$

Thus K is an invariant; it is easy to see that it is the Gauss curvature of M . The expression k depends just on the orientation of the moving frames. Let us look for the existence of a tangent vector field $V = \xi v_1 + \eta v_2$ such that there is a non-trivial tangent vector field $W = \xi' v_1 + \eta' v_2$ satisfying $WV \subset T(M)$. Because of

$$dV = dx \cdot v_1 + dy \cdot v_2 + x(\omega_1^2 v_2 + \omega_1^3 v_3 + \omega_1^4 v_4) + \\ + y(-\omega_1^2 v_2 + \omega_2^3 v_3 + \omega_2^4 v_4),$$

we get

$$WV \equiv (a_1 \xi \xi' + a_2 \xi \eta' + a_2 \eta \xi' + a_3 \eta \eta') v_3 + \\ + (b_1 \xi \xi' + b_2 \xi \eta' + b_2 \eta \xi' + b_3 \eta \eta') v_4 \pmod{T(M)};$$

because of $WV \equiv 0$ and $\xi'^2 + \eta'^2 \neq 0$, we get

$$(12) \quad (a_1 b_2 - a_2 b_1) \xi^2 + (a_1 b_3 - a_3 b_1) \xi \eta + (a_2 b_3 - a_3 b_2) \eta^2 = 0.$$

The equation (12) is the equation of the so-called conjugate directions $\{\xi v_1 + \eta v_2\}$ of M .

2. Let Φ be an infinitesimal second order deformation of our surface M ; Φ is obviously given by (1) where we have to replace $\omega_3^4, \omega_3^5, \omega_4^5$ by $\omega_3^4 + t\varphi_3^4 + \dots$, $\omega_3^5 + t\varphi_3^5 + \dots$, $\omega_4^5 + t\varphi_4^5 + \dots$ resp. Comparing the terms at t in the integrability conditions, we get

$$(13) \quad \omega_1^4 \wedge \varphi_3^4 + \omega_1^5 \wedge \varphi_3^5 = 0, \quad \omega_1^3 \wedge \varphi_3^4 - \omega_1^5 \wedge \varphi_4^5 = 0, \\ \omega_2^4 \wedge \varphi_3^4 + \omega_2^5 \wedge \varphi_3^5 = 0, \\ \omega_2^3 \wedge \varphi_3^4 - \omega_2^5 \wedge \varphi_4^5 = 0, \quad \omega_1^3 \wedge \varphi_3^5 + \omega_1^4 \wedge \varphi_4^5 = 0, \\ \omega_2^3 \wedge \varphi_3^5 + \omega_2^4 \wedge \varphi_4^5 = 0;$$

$$(14) \quad d\varphi_3^4 = -\omega_3^5 \wedge \varphi_4^5 - \varphi_3^5 \wedge \omega_4^5, \\ d\varphi_3^5 = \omega_3^4 \wedge \varphi_4^5 + \varphi_3^4 \wedge \omega_4^5, \quad d\varphi_4^5 = -\omega_3^4 \wedge \varphi_3^5 - \varphi_3^4 \wedge \omega_3^5.$$

Because of (4) and (13), we get the existence of functions A_1, \dots, C_4 such that

$$(15) \quad b_1 \varphi_3^4 + c_1 \varphi_3^5 = A_1 \omega^1 + A_2 \omega^2, \quad a_1 \varphi_3^4 - c_1 \varphi_4^5 = B_1 \omega^1 + B_2 \omega^2, \\ b_2 \varphi_3^4 + c_2 \varphi_3^5 = A_2 \omega^1 + A_3 \omega^2, \quad a_2 \varphi_3^4 - c_2 \varphi_4^5 = B_2 \omega^1 + B_3 \omega^3, \\ b_3 \varphi_3^4 + c_3 \varphi_3^5 = A_3 \omega^1 + A_4 \omega^2, \quad a_3 \varphi_3^4 - c_3 \varphi_4^5 = B_3 \omega^1 + B_4 \omega^2, \\ a_1 \varphi_3^5 + b_1 \varphi_4^5 = C_1 \omega^1 + C_2 \omega^2, \\ a_2 \varphi_3^5 + b_2 \varphi_4^5 = C_2 \omega^1 + C_3 \omega^2, \\ a_3 \varphi_3^5 + b_3 \varphi_4^5 = C_3 \omega^1 + C_4 \omega^2.$$

Write

$$(16) \quad \varphi_3^4 = x_1\omega^1 + x_2\omega^2, \quad \varphi_3^5 = y_1\omega^1 + y_2\omega^2, \quad \varphi_4^5 = z_1\omega^1 + z_2\omega^2;$$

from (15), we obtain

$$(17) \quad \begin{aligned} b_2x_1 - b_1x_2 + c_2y_1 - c_1y_2 &= 0, \\ b_3x_1 - b_2x_2 + c_3y_1 - c_2y_2 &= 0, \\ a_2x_1 - a_1x_2 &\quad - c_2z_1 + c_1z_2 = 0, \\ a_3x_1 - a_2x_2 &\quad - c_3z_1 + c_2z_2 = 0, \\ a_2y_1 - a_1y_2 + b_2z_1 - b_1z_2 &= 0, \\ a_3y_1 - a_2y_2 + b_3z_1 - b_2z_2 &= 0. \end{aligned}$$

We are now in the position to prove the following local

Theorem 1. *Let $M \equiv M(G) \subset E^5$ be a surface such that $\dim T_m^2(M) = 5$ for each $m \in M$. Then each infinitesimal second order deformation Φ of M is trivial.*

Proof. From (1), we obtain

$$(18) \quad v_1m = v_1, \quad v_2m = v_2,$$

$$(19) \quad v_1v_1m = (\cdot)v_2 + a_1v_3 + b_1v_4 + c_1v_5,$$

$$v_2v_1m = (\cdot)v_2 + a_2v_3 + b_2v_4 + c_2v_5,$$

$$v_2v_2m = (\cdot)v_1 + a_3v_3 + b_3v_4 + c_3v_5.$$

The vectors (19) are linearly independent because of the condition $\dim T_m^2(M) = 5$, and we may choose the frames in such a way that

$$(20) \quad b_1 = c_1 = c_2 = 0, \quad a_1b_2c_3 \neq 0.$$

The system (17) is then reduced to

$$(21) \quad b_2x_1 = 0, \quad b_3x_1 - b_2x_2 + c_3y_1 = 0, \quad a_2x_1 - a_1x_2 = 0,$$

$$a_3x_1 - a_2x_2 - c_3z_1 = 0,$$

$$a_2y_1 - a_1y_2 + b_2z_1 = 0, \quad a_3y_1 - a_2y_2 + b_3z_1 - b_2z_2 = 0,$$

and we get $x_1 = x_2 = y_1 = y_2 = z_1 = z_2 = 0$, i.e.,

$$(22) \quad \varphi_3^4 = \varphi_3^5 = \varphi_4^5 = 0.$$

Thus Φ is trivial. QED.

3. Now, let us prove a global result.

Theorem 2. *Let $G \subset \mathbb{R}^2$ be a bounded domain and let $M : G \cup \partial G \rightarrow E^5$ be a surface such that: (i) $\dim T_m^2(M) = 4$ for each $m \in M$; (ii) there is $K \neq k$ or $K \neq -k$ on M ; (iii) there are no real conjugate directions on M . Let Φ be an infinitesimal second order deformation of M which is trivial on the boundary of M . Then Φ is trivial on M .*

Proof. Choose moving frames $\{v_i\}$ of M and suppose $K \neq k$ on M ; of course, we have (5). From (17) and (i), $x_1 = x_2 = 0$, i.e.,

$$(23) \quad \varphi_3^4 = 0.$$

From (13), (14) and (17),

$$(24) \quad \omega_1^3 \wedge \varphi_3^5 + \omega_1^4 \wedge \varphi_4^5 = 0, \quad \omega_2^3 \wedge \varphi_3^5 + \omega_2^4 \wedge \varphi_4^5 = 0,$$

$$(25) \quad d\varphi_3^5 = \omega_3^4 \wedge \varphi_4^5, \quad d\varphi_4^5 = -\omega_3^4 \wedge \varphi_3^5,$$

$$(26) \quad a_2y_1 - a_1y_2 + b_2z_1 - b_1z_2 = 0, \quad a_3y_1 - a_2y_2 + b_3z_1 - b_2z_2 = 0.$$

From (16_{2,3}) and (25), we get the existence of functions S_1, \dots, S_6 such that

$$(27) \quad \begin{aligned} dy_1 - y_2\omega_1^2 - z_1\omega_3^4 &= S_1\omega^1 + S_2\omega^2, \\ dy_2 + y_1\omega_1^2 - z_2\omega_3^4 &= S_2\omega^1 + S_3\omega^2, \\ dz_1 - z_2\omega_1^2 + y_1\omega_3^4 &= S_4\omega^1 + S_5\omega^2, \\ dz_2 + z_1\omega_1^2 + y_2\omega_3^4 &= S_5\omega^1 + S_6\omega^2. \end{aligned}$$

From (4₁₋₄) + (5), we get the existence of functions α_1, \dots, β_4 such that

$$(28) \quad \begin{aligned} da_1 - 2a_2\omega_1^2 - b_1\omega_3^4 &= \alpha_1\omega^1 + \alpha_2\omega^2, \\ da_2 + (a_1 - a_3)\omega_1^2 - b_2\omega_3^4 &= \alpha_2\omega^1 + \alpha_3\omega^2, \\ da_3 + 2a_2\omega_1^2 - b_3\omega_3^4 &= \alpha_3\omega^1 + \alpha_4\omega^2, \\ db_1 - 2b_2\omega_1^2 + a_1\omega_3^4 &= \beta_1\omega^1 + \beta_2\omega^2, \\ db_2 + (b_1 - b_3)\omega_1^2 + a_2\omega_3^4 &= \beta_2\omega^1 + \beta_3\omega^2, \\ db_3 + 2b_2\omega_1^2 + a_3\omega_3^4 &= \beta_3\omega^1 + \beta_4\omega^2. \end{aligned}$$

The differential consequences of (26) are then

$$(29) \quad \begin{aligned} a_2S_1 - a_1S_2 + b_2S_4 - b_1S_5 &= -\alpha_2y_1 + \alpha_1y_2 - \beta_2z_1 + \beta_1z_2, \\ a_2S_2 - a_1S_3 + b_2S_5 - b_1S_6 &= -\alpha_3y_1 + \alpha_2y_2 - \beta_3z_1 + \beta_2z_2, \\ a_3S_1 - a_2S_2 + b_3S_4 - b_2S_5 &= -\alpha_3y_1 + \alpha_2y_2 - \beta_3z_1 + \beta_2z_2, \\ a_3S_2 - a_2S_3 + b_3S_5 - b_2S_6 &= -\alpha_4y_1 + \alpha_3y_2 - \beta_4z_1 + \beta_3z_2. \end{aligned}$$

Consider the system

$$(30) \quad a_1X + b_1Y - a_2Z - b_2T = Q_1, \quad a_2X + b_2Y - a_3Z - b_3T = Q_2$$

for X, Y, Z, T . It is easy to show that the convenient combinations of the equations (30) lead to

$$(31) \quad \begin{aligned} (K - k)X &= (a_3b_2 - a_2b_3 + b_1b_3 - b_2^2)(X + T) + \\ &+ (a_2b_2 - a_3b_1)(Y - Z) + (a_3 - b_2)Q_1 + (b_1 - a_2)Q_2, \\ (K - k)Y &= (a_2b_2 - a_1b_3)(X + T) + (a_1a_3 - a_2^2 + a_3b_2 - a_2b_3)(Y - Z) + \\ &+ (a_2 + b_3)Q_1 - (a_1 + b_2)Q_2, \\ (K - k)Z &= (a_2b_2 - a_1b_3)(X + T) + (a_1b_2 - a_2b_1 - b_1b_3 + b_2^2)(Y - Z) + \\ &+ (a_2 + b_3)Q_1 - (a_1 + b_2)Q_2, \\ (K - k)T &= (a_1a_3 - a_2^2 - a_1b_2 + a_2b_1)(X + T) + (a_3b_1 - a_2b_2)(Y - Z) + \\ &+ (b_2 - a_3)Q_1 + (a_2 - b_1)Q_2. \end{aligned}$$

Applying this auxiliary result to the systems (26), (29_{1,3}) and (29_{2,4}) resp., we get

$$(32) \quad \begin{aligned} (K - k)y_2 &= (a_3b_2 - a_2b_3 + b_1b_3 - b_2^2)(y_2 + z_1) + \\ &+ (a_2b_2 - a_3b_1)(z_2 - y_1), \\ (K - k)z_2 &= (a_2b_2 - a_1b_3)(y_2 + z_1) + \\ &+ (a_1a_3 - a_2^2 + a_3b_2 - a_2b_3)(z_2 - y_1), \\ (K - k)y_1 &= (a_2b_2 - a_1b_3)(y_2 + z_1) + \\ &+ (a_1b_2 - a_2b_1 - b_1b_3 + b_2^2)(z_2 - y_1), \\ (K - k)z_1 &= (a_1a_3 - a_2^2 - a_1b_2 + a_2b_1)(y_2 + z_1) + \\ &+ (a_3b_1 - a_2b_2)(z_2 - y_1); \end{aligned}$$

$$(33) \quad \begin{aligned} (K - k)S_2 &= (a_3b_2 - a_2b_3 + b_1b_3 - b_2^2)(S_2 + S_4) + (a_2b_2 - a_3b_1) \cdot \\ &\cdot (S_5 - S_1) + (a_3 - b_2)q_1 + (b_1 - a_2)q_2, \\ (K - k)S_5 &= (a_2b_2 - a_1b_3)(S_2 + S_4) + (a_1a_3 - a_2^2 + a_3b_2 - a_2b_3) \cdot \\ &\cdot (S_5 - S_1) + (a_2 + b_3)q_1 - (a_1 + b_2)q_2, \\ (K - k)S_1 &= (a_2b_2 - a_1b_3)(S_2 + S_4) + (a_1b_2 - a_2b_1 - b_1b_3 + b_2^2) \cdot \\ &\cdot (S_5 - S_1) + (a_2 + b_3)q_1 - (a_1 + b_2)q_2, \\ (K - k)S_4 &= (a_1a_3 - a_2^2 - a_1b_2 + a_2b_1)(S_2 + S_4) + (a_3b_1 - a_2b_2) \cdot \\ &\cdot (S_5 - S_1) + (b_2 - a_3)q_1 + (a_2 - b_1)q_2 \end{aligned}$$

with

$$\begin{aligned}
 q_1 &= \alpha_2 y_1 - \alpha_1 y_2 + \beta_2 z_1 - \beta_1 z_2, & q_2 &= \alpha_3 y_1 - \alpha_2 y_2 + \beta_3 z_1 - \beta_2 z_2; \\
 (34) \quad (K - k) S_3 &= (a_3 b_2 - a_2 b_3 + b_1 b_3 - b_2^2)(S_3 + S_5) + (a_2 b_2 - a_3 b_1) \cdot \\
 &\quad \cdot (S_6 - S_2) + (a_3 - b_2) q'_1 + (b_1 - a_2) q'_2, \\
 (K - k) S_6 &= (a_2 b_2 - a_1 b_3)(S_3 + S_5) + (a_1 a_3 - a_2^2 + a_3 b_2 - a_2 b_3) \cdot \\
 &\quad \cdot (S_6 - S_2) + (a_2 + b_3) q'_1 - (a_1 + b_2) q'_2, \\
 (K - k) S_2 &= (a_2 b_2 - a_1 b_3)(S_3 + S_5) + (a_1 b_2 - a_2 b_1 - b_1 b_3 + b_2^2) \cdot \\
 &\quad \cdot (S_6 - S_2) + (a_2 + b_3) q'_1 - (a_1 + b_2) q'_2, \\
 (K - k) S_5 &= (a_1 a_3 - a_2^2 - a_1 b_2 + a_2 b_1)(S_3 + S_5) + (a_3 b_1 - a_2 b_2) \cdot \\
 &\quad \cdot (S_6 - S_2) + (b_2 - a_3) q'_1 + (a_2 - b_1) q'_2
 \end{aligned}$$

with

$$q'_1 = \alpha_3 y_1 - \alpha_2 y_2 + \beta_3 z_1 - \beta_2 z_2, \quad q'_2 = \alpha_4 y_1 - \alpha_3 y_2 + \beta_4 z_1 - \beta_3 z_2.$$

It follows from (32) and $K \neq k$ that any linear combination of y_1, y_2, z_1, z_2 may be written as a combination of $y_2 + z_1$ and $y_1 - z_2$.

From (27), we get

$$\begin{aligned}
 (35) \quad d(y_1 - z_2) - (y_2 + z_1)(\omega_1^2 + \omega_3^4) &= (S_1 - S_5) \omega^1 + (S_2 - S_6) \omega^2, \\
 d(y_2 + z_1) + (y_1 - z_2)(\omega_1^2 + \omega_3^4) &= (S_2 + S_4) \omega^1 + (S_3 + S_5) \omega^2.
 \end{aligned}$$

Let G be covered by a system of isothermic coordinates (u, v) ; let

$$(36) \quad I = r^2(du^2 + dv^2), \quad r(u, v) > 0, \quad \text{i.e.,} \quad \omega^1 = r du, \quad \omega^2 = r dv.$$

Then

$$\begin{aligned}
 (37) \quad \frac{\partial(y_1 - z_2)}{\partial u} &= (S_1 - S_5) r + \varrho_1(y_2 + z_1), \\
 \frac{\partial(y_1 - z_2)}{\partial v} &= (S_2 - S_6) r + \varrho_2(y_2 + z_1), \\
 \frac{\partial(y_2 + z_1)}{\partial u} &= (S_2 + S_4) r + \varrho_3(y_1 - z_2), \\
 \frac{\partial(y_2 + z_1)}{\partial v} &= (S_3 + S_5) r + \varrho_4(y_1 - z_2),
 \end{aligned}$$

$\varrho_1, \dots, \varrho_4$ being easy to calculate. From (37) and $(33_1) + (34_3)$ or $(33_2) + (34_4)$ resp., we get a system of the form ($i = 1, 2$)

$$(38) \quad a_{i1} \frac{\partial(y_1 - z_2)}{\partial u} + a_{i2} \frac{\partial(y_1 - z_2)}{\partial v} + b_{i1} \frac{\partial(y_2 + z_1)}{\partial u} + \\ + b_{i2} \frac{\partial(y_2 + z_1)}{\partial v} = c_{i1}(y_1 - z_2) + c_{i2}(y_2 + z_1)$$

with

$$(39) \quad a_{11} = a_2 b_2 - a_3 b_1, \quad a_{12} = b_1 b_3 - b_2^2 + a_2 b_1 - a_1 b_2, \\ a_{21} = a_1 a_3 - a_2^2 + a_3 b_2 - a_2 b_3, \quad a_{22} = a_2 b_2 - a_3 b_1, \\ b_{11} = b_2^2 - b_1 b_3 + a_2 b_3 - a_3 b_2, \quad b_{12} = a_2 b_2 - a_1 b_3, \\ b_{21} = a_1 b_3 - a_2 b_2, \quad b_{22} = a_1 a_3 - a_2^2 + a_2 b_1 - a_1 b_2.$$

Recall that the system (38) is called elliptic if the quadratic form

$$(40) \quad \varphi = (a_{12} b_{22} - a_{22} b_{12}) \mu^2 - (a_{11} b_{22} - a_{21} b_{12} + a_{12} b_{21} - a_{22} b_{11}) \mu v + \\ + (a_{11} b_{21} - a_{21} b_{11}) v^2$$

is definite. In our case,

$$(41) \quad \varphi = (k - K) \{ (a_1 b_2 - a_2 b_1) \mu^2 + (a_1 b_3 - a_3 b_1) \mu v + (a_2 b_3 - a_3 b_2) v^2 \},$$

and φ is definite because of the suppositions of our Theorem. Thus $y_1 - z_2 = y_2 + z_1 = 0$ in G and (32) implies $y_1 = y_2 = z_1 = z_2 = 0$ in G .

In the case $K \neq -k$, we have to use $y_1 + z_2, y_2 - z_1$ instead of $y_1 - z_2, y_2 + z_1$ resp. QED.

Let us remark that locally Φ does not need to be trivial.

Bibliography

- [1] *E. Bompiani*: Geometrie riemanniane di specie superiore. R. Acc. Italia, Mem. Cl. di Sci. Fiz., Mat. e Nat., Vol. VI, 1935.

Author's address: 771 46 Olomouc, Leninova 26, ČSSR (Přírodovědecká fakulta University Palackého).