

Oldřich Kopeček

Existence of monomorphisms of partial unary algebras

Czechoslovak Mathematical Journal, Vol. 28 (1978), No. 3, 462–473

Persistent URL: <http://dml.cz/dmlcz/101551>

Terms of use:

© Institute of Mathematics AS CR, 1978

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

EXISTENCE OF MONOMORPHISMS OF PARTIAL UNARY ALGEBRAS

OLDŘICH KOPEČEK, Brno
 (Received December 10, 1976)

1. MONOMORPHISMS

1.0. Notation. We denote by Ord the class of all ordinals and by N the set of all finite ordinals.

Let $\infty_1, \infty_2 \notin \text{Ord}$. We suppose that $\alpha < \infty_1 < \infty_2$ for each $\alpha \in \text{Ord}$.

If $\alpha \in \text{Ord}$ then we put $W(\alpha) = \{\beta \in \text{Ord}; \beta < \alpha\}$. Further, $W(\infty_1) = N, W(\infty_2) = N \cup \{\infty_1\}$.

If A is a set we denote by $|A|$ the cardinal number of A . Let φ be a partial map from A into a set B . We put $\text{dom } \varphi = \{x \in A; \text{there exists } y \in B \text{ such that } (x, y) \in \varphi\}$. If $\text{dom } \varphi = A$ then we write $\varphi : A \rightarrow B$. Further, if $C \subseteq A, D \subseteq B$ then we put $\varphi C = \{\varphi x; x \in C\}, \varphi^{-1}D = \{x \in A; \varphi x \in D\}, \varphi \upharpoonright C = \varphi \cap (C \times B)$ (the restriction of φ on C).

Let A_i be an ordered set with an order $\leq_i (i = 1, 2)$. We put $B = \bigcup_{a \in A_1} (A_1, a) \cup \bigcup_{a' \in A_2} (A_2, a')$ and define an order \leq on B in the following way: if $(t, a), (u, b) \in B$ then we put

$$(t, a) \leq (u, b) \text{ iff } t = u = A_1, \quad a \leq_1 b \text{ or } t = u = A_2, \\ a \leq_2 b \text{ or } t = A_1, \quad u = A_2.$$

Then we write $B = A_1 \oplus A_2$ and the ordered set B is called the *ordinal sum* of A_1 and A_2 .

1.1. Definition. Let A, B be sets, $S_1 : A \rightarrow \text{Ord} \cup \{\infty_1, \infty_2\}, S_2 : B \rightarrow \text{Ord} \cup \{\infty_1, \infty_2\}$ arbitrary maps. Let $F : A \rightarrow B$ be a map such that $S_1 x \leq S_2 Fx$ for each $x \in A$. Then F is called a *degree map* (abbreviation a *d-map*) with respect to S_1, S_2 . Further, if F is an injection then we speak about a *d-injection*.

1.2. Theorem. Let A, B be finite sets, $S_1 : A \rightarrow N \cup \{\infty_1, \infty_2\}, S_2 : B \rightarrow N \cup \{\infty_1, \infty_2\}$ maps. We put $A^i = S_1^{-1}i, B^i = S_2^{-1}i$ for each $i \in N \cup \{\infty_1, \infty_2\}$. Then there is a *d-injection* $F : A \rightarrow B$ with respect to S_1, S_2 if and only if $|A - \bigcup_{i \in W(n)} A^i| \leq |B - \bigcup_{i \in W(n)} B^i|$ for each $n \in N \cup \{\infty_1, \infty_2\}$.

Proof. Necessity. Let there be a d-injection $F : A \rightarrow B$. Let $n \in N \cup \{\infty_1, \infty_2\}$ be arbitrary. Then, for each $x \in A - \bigcup_{i \in W(n)} A^i$, $S_2 Fx \geq S_1 x \notin W(n)$ holds and thus, $Fx \in B - \bigcup_{i \in W(n)} B^i$. Hence $F(A - \bigcup_{i \in W(n)} A^i) \subseteq B - \bigcup_{i \in W(n)} B^i$ and we obtain $|A - \bigcup_{i \in W(n)} A^i| = |F(A - \bigcup_{i \in W(n)} A^i)| \leq |B - \bigcup_{i \in W(n)} B^i|$.

Sufficiency. Since $S_1 A$ is finite there is $n_0 \in N \cup \{\infty_1, \infty_2\}$ such that $n_0 = \max S_1 A$. Then $|A^{n_0}| = |A - \bigcup_{i \in W(n_0)} A^i| \leq |B - \bigcup_{i \in W(n_0)} B^i|$. Thus, there is a d-injection $F_0 : A^{n_0} \rightarrow B$.

Let now $p \in N$, let $n_0 > n_1 > \dots > n_p$ be the last $p+1$ elements in $S_1 A$ and let there be a d-injection $F_p : \bigcup_{i \in W(p+1)} A^{n_i} \rightarrow B$.

If $A = \bigcup_{i \in W(p+1)} A^{n_i}$ then the proof is complete. Therefore, let $A \neq \bigcup_{i \in W(p+1)} A^{n_i}$, i.e. $S_1 A \cap W(n_p) \neq \emptyset$. Then we put $n_{p+1} = \max(S_1 A \cap W(n_p))$. Hence $|A^{n_{p+1}}| + |\bigcup_{i \in W(p+1)} A^{n_i}| = |A - \bigcup_{i \in W(n_{p+1})} A^i| \leq |B - \bigcup_{i \in W(n_{p+1})} B^i|$. Further, $\bigcup_{i \in W(p+1)} A^{n_i} = A - \bigcup_{i \in W(n_p)} A^i$ and since F_p is a d-map we have $F_p \bigcup_{i \in W(p+1)} A^{n_i} = F_p(A - \bigcup_{i \in W(n_p)} A^i) \subseteq B - \bigcup_{i \in W(n_p)} B^i \subseteq B - \bigcup_{i \in W(n_{p+1})} B^i$. Thus, since F_p is an injection we obtain $|A^{n_{p+1}}| \leq |B - \bigcup_{i \in W(n_{p+1})} B^i| - |\bigcup_{i \in W(p+1)} A^{n_i}| = |B - \bigcup_{i \in W(n_{p+1})} B^i| - |F_p \bigcup_{i \in W(p+1)} A^{n_i}| \leq |(B - \bigcup_{i \in W(n_{p+1})} B^i) - F_p \bigcup_{i \in W(p+1)} A^i| = |B - (\bigcup_{i \in W(p+1)} B^i \cup F_p \bigcup_{i \in W(p+1)} A^{n_i})|$. It follows that there is an injection $G : A^{n_{p+1}} \rightarrow B - (\bigcup_{i \in W(n_{p+1})} B^i \cup F_p \bigcup_{i \in W(p+1)} A^{n_i})$. Further, G is a d-injection and since $F_p \bigcup_{i \in W(p+1)} A^{n_i}$ and $GA^{n_{p+1}}$ are disjoint we obtain that $F_{p+1} = F_p \cup G$ is a d-injection $F_{p+1} : \bigcup_{i \in W(p+2)} A^{n_i} \rightarrow B$.

1.3. Lemma. Let A_1, A_2 be finite sets, $S_i : A_i \rightarrow N \cup \{\infty_1, \infty_2\}$ maps ($i = 1, 2$). We put $A_i^n = S_i^{-1} n$ for each $n \in N \cup \{\infty_1, \infty_2\}$ ($i = 1, 2$). Let $x_i \in A_i$ ($i = 1, 2$) be such that $S_1 x_1 = S_2 x_2$. We put $\bar{A}_i = A_i - \{x_i\}$, $\bar{A}_i^n = (S_i | \bar{A}_i)^{-1} n$ for each $n \in N \cup \{\infty_1, \infty_2\}$ ($i = 1, 2$). Let $n \in N \cup \{\infty_1, \infty_2\}$ be arbitrary. If $|A_1 - \bigcup_{i \in W(n)} A_1^i| \leq |A_2 - \bigcup_{i \in W(n)} A_2^i|$ then $|\bar{A}_1 - \bigcup_{i \in W(n)} \bar{A}_1^i| \leq |\bar{A}_2 - \bigcup_{i \in W(n)} \bar{A}_2^i|$.

Proof. We put $S_1 x_1 = S_2 x_2 = p$. Then

$$\bar{A}_i - \bigcup_{j \in W(n)} \bar{A}_i^j = \begin{cases} (A_i - \bigcup_{j \in W(n)} A_i^j) - \{x_i\} & \text{if } n \leq p \\ A_i - \bigcup_{j \in W(n)} A_i^j & \text{if } n > p \end{cases} \quad \text{where } i = 1, 2.$$

$$\text{If } n \in N \cup \{\infty_1, \infty_2\}, \quad n \leq p \text{ then } |\bar{A}_1 - \bigcup_{i \in W(n)} \bar{A}_1^i| = |A_1 - \bigcup_{i \in W(n)} A_1^i| - 1 \leq \\ \leq |A_2 - \bigcup_{i \in W(n)} A_2^i| - 1 = |\bar{A}_2 - \bigcup_{i \in W(n)} \bar{A}_2^i|.$$

$$\text{If } n \in N \cup \{\infty_1, \infty_2\}, \quad n > p \text{ then } |\bar{A}_1 - \bigcup_{i \in W(n)} \bar{A}_1^i| = |A_1 - \bigcup_{i \in W(n)} A_1^i| \leq |A_2 - \\ - \bigcup_{i \in W(n)} A_2^i| = |\bar{A}_2 - \bigcup_{i \in W(n)} \bar{A}_2^i|.$$

1.4. Definition. (a) Let A be a non-empty set, f a partial map from the set A into A . Then the ordered pair $\mathbf{A} = (A, f)$ is called a *partial unary algebra*.

(b) Let $\mathbf{A} = (A, f)$, $\mathbf{B} = (B, g)$ be partial unary algebras, let $F: A \rightarrow B$ be a map. Then F is called a *homomorphism* of \mathbf{A} into \mathbf{B} if $x \in \text{dom } f$ implies $Fx \in \text{dom } g$ and $Ffx = gFx$ for each $x \in A$; we write $F: \mathbf{A} \rightarrow \mathbf{B}$. If F is a homomorphism and an injective map then F is called a *monomorphism*.

(c) The category of all partial unary algebras where morphisms are homomorphisms is denoted by \mathcal{U} .

1.5. Definition. Let $\mathbf{A} = (A, f) \in \mathcal{U}$.

(a) We put $D\mathbf{A} = A - \text{dom } f$.

(b) If $A' \subseteq A$ then we put $\langle A' \rangle_{\mathbf{A}} = f^{-1}A'$; especially, we put $\langle x \rangle_{\mathbf{A}} = f^{-1}x$ for every $x \in A$.

(c) We put $f^0 = \text{id}_A$. Suppose that we have defined a partial map f^{n-1} from A into A for $n \in N - \{0\}$. We denote by f^n the following partial map from A into A : if $x \in \text{dom } f^{n-1}$ and $f^{n-1}x \in \text{dom } f$ then we put $f^n x = ff^{n-1}x$.

(d) Let $x \in A$ be arbitrary. Then we define $[x]_{\mathbf{A}} = \{y \in A; \text{ there is } n \in N \text{ with } x \in \text{dom } f^n \text{ and } y = f^n x\}$.

(e) Let $x, y \in A$ be arbitrary. Then we put $\varrho\mathbf{A} = \{(x, y) \in A^2; [x]_{\mathbf{A}} \cap [y]_{\mathbf{A}} \neq \emptyset\}$. If $\varrho\mathbf{A} = A^2$ then \mathbf{A} is called a *connected partial unary algebra* (abbreviation a *c-algebra*). The category of all c-algebras where morphisms are homomorphism defined in 1.4(b) is denoted by \mathcal{U}^c .

(f) Let $\mathbf{A} \in \mathcal{U}^c$, $x \in A$ be arbitrary. Then we put $[x]_{\mathbf{A}}^0 = [x]_{\mathbf{A}}$, $[x]_{\mathbf{A}}^{-1} = \langle [x]_{\mathbf{A}} \rangle_{\mathbf{A}} - [x]_{\mathbf{A}}$. Suppose that we have defined the set $[x]_{\mathbf{A}}^{-n}$ for $n \in N - \{0\}$. Then we put $[x]_{\mathbf{A}}^{-(n+1)} = \langle [x]_{\mathbf{A}}^{-n} \rangle_{\mathbf{A}}$.

(g) Let $\mathbf{A} \in \mathcal{U}^c$, $x, y \in A$ be arbitrary. Then we put $\langle y \rangle_{\mathbf{A}, x} = \langle y \rangle_{\mathbf{A}} - [x]_{\mathbf{A}}$.

(i) Let $\mathbf{A} = (A, f) \in \mathcal{U}^c$, $x \in A$ be arbitrary. Then the following assertions hold:

(a) $[x]_{\mathbf{A}}^{-(n+1)} = \bigcup_{y \in [x]_{\mathbf{A}}^{-n}} \langle y \rangle_{\mathbf{A}, x}$ with disjoint terms for each $n \in N$,

(b) $A = \bigcup_{n \in N} [x]_{\mathbf{A}}^{-n}$ with disjoint terms.

(See [7], 2.1 and [4], 3.9 (c).)

1.6. Definition. Let $A = (A, f) \in \mathcal{U}^c$. Then we define $ZA = \{x \in A; \text{there is } n \in N - \{0\} \text{ such that } f^n x = x\}$, $RA = [ZA]$.

(ii) Let $A = (A, f) \in \mathcal{U}^c$. If $ZA \neq \emptyset$, $x \in A$ then there is $n \in N$ such that $f^n x \in ZA$. (See [3], 1.17 (a).)

1.7. Lemma. Let $A = (A, f) \in \mathcal{U}^c$, $x, y \in A$ be arbitrary. We put $p_0 = \min \{n \in N; f^n x \in ZA\}$. Then the following assertions hold:

(a) If $y \in \bigcup_{n \in N - \{0\}} [x]_A^{-n}$ then $\langle y \rangle_A = \langle y \rangle_{A,x}$.

(b) $\langle x \rangle_A = \begin{cases} \langle x \rangle_{A,x} & \text{if } x \notin ZA, \\ \langle x \rangle_{A,x} \cup \{f^{RA-1}x\} & \text{if } x \in ZA. \end{cases}$

(c) If $y = f^p x$, $p \in N - \{0, p_0\}$ then $\langle y \rangle_A = \langle y \rangle_{A,x} \cup \{f^{p-1}x\}$.

(d) If $y = f^{p_0} x$ where $p_0 \in N - \{0\}$ then $\langle y \rangle_A = \langle y \rangle_{A,x} \cup \{f^{p_0-1}x, f^{RA+p_0-1}x\}$.

(e) If $x \in ZA$ then $\langle y \rangle_A = \begin{cases} \langle y \rangle_{A,x} & \text{if } y \in \bigcup_{n \in N - \{0\}} [x]_A^{-n} \\ \langle y \rangle_{A,x} \cup \{f^{p-1}x\} & \text{if } y = f^p x, p \in N - \{0\}. \end{cases}$

(f) If $x \notin ZA$ then

$$\langle y \rangle_A = \begin{cases} \langle y \rangle_{A,x} & \text{if } y \in \{x\} \cup \bigcup_{n \in N - \{0\}} [x]_A^{-n}, \\ \langle y \rangle_{A,x} \cup \{f^{p-1}x\} & \text{if } y = f^p x, p \in N - \{0, p_0\}, \\ \langle y \rangle_{A,x} \cup \{f^{p_0-1}x, f^{RA+p_0-1}x\} & \text{if } y = f^{p_0} x. \end{cases}$$

Proof of (a). Let $y \in [x]_A^{-n}$ where $n \in N - \{0\}$. Then $\langle y \rangle_A \subseteq [x]_A^{-(n+1)}$ by 1.5 (f). Hence $\langle y \rangle_A \cap [x]_A = \emptyset$ by (i) (b) which implies $\langle y \rangle_{A,x} = \langle y \rangle_A - [x]_A = \langle y \rangle_A$.

Proof of (b). If $x \notin ZA$ then $\langle x \rangle_A \cap [x]_A = \emptyset$ by 1.5 (d) and 1.6 and thus $\langle x \rangle_{A,x} = \langle x \rangle_A - [x]_A = \langle x \rangle_A$.

If $x \in ZA$ then $\langle x \rangle_A \cap [x]_A = \{f^{RA-1}x\}$ by 1.5 (d) and 1.6 which implies $\langle x \rangle_{A,x} = \langle x \rangle_A - \{f^{RA-1}x\}$.

Proof of (c). Let $y = f^p x$, $p \in N - \{0, p_0\}$. Then $\langle y \rangle_A \cap [x]_A = \{f^{p-1}x\}$ and therefore, $\langle y \rangle_{A,x} = \langle y \rangle_A - \{f^{p-1}x\}$.

Proof of (d). Let $y = f^{p_0} x$ where $p_0 \in N - \{0\}$. Then $\langle y \rangle_A \cap [x]_A = \{f^{p_0-1}x, f^{RA+p_0-1}x\}$ and we have $\langle y \rangle_{A,x} = \langle y \rangle_A - \{f^{p_0-1}x, f^{RA+p_0-1}x\}$.

(e) and (f) are consequences of (a), (b), (c), (d) and (i) (b).

1.8. Definition. Let $A = (A, f) \in \mathcal{U}^c$. Then we define

(a) the set $KA = \{x \in A - ZA; \text{there is a sequence } (x_i)_{i \in N} \text{ such that } x_i \in \text{dom } f \text{ for each } i \in N - \{0\}, x_0 = x \text{ and } fx_{i+1} = x_i \text{ for each } i \in N\}$;

(b) $A^{\infty_2} = ZA$, $A^{\infty_1} = KA$, $A^0 = \{x \in A; \langle x \rangle_A = \emptyset\}$; if $\alpha \in \text{Ord} - \{0\}$ is arbitrary and if the sets A^α have been defined for each $\alpha \in W(\alpha)$ then we put $A^\alpha = \{x \in A - \bigcup_{\lambda \in W(\alpha)} A^\lambda, \langle x \rangle_A \subseteq \bigcup_{\lambda \in W(\alpha)} A^\lambda\}$;

(c) $\mathfrak{A} = \min \{\alpha \in \text{Ord}; A^\alpha = \emptyset\}$;

(d) a map $SA : A \rightarrow \text{Ord} \cup \{\infty_1, \infty_2\}$ by the condition $SAx = \alpha$ for each $x \in A^\alpha$, $\alpha \in W(\mathfrak{A}) \cup \{\infty_1, \infty_2\}$; SAx is called the *degree* of x .

(iii) Let $A = (A, f) \in \mathcal{U}^c$. Then

(a) $(KA \cup ZA, f \upharpoonright KA \cup ZA)$, $(ZA, f \upharpoonright ZA)$ are subalgebras of A ,

(b) $|DA| \leq 1$,

(c) $DA \neq \emptyset$ iff the following conditions hold: $RA = 0$ and there is $x \in A$ such that $|\langle x \rangle_A| < \aleph_0$.

(See [4], 2.10, 2.15 (a), 2.1 and 2.9.)

1.9. Definition. Let $A \in \mathcal{U}^c$.

(a) The set $DA \cup KA \cup ZA$ is called the *kernel* of A . If $DA \cup KA \cup ZA \neq \emptyset$ then we say that A has the kernel.

(b) If $DA \neq \emptyset$ then we denote by dA the only point with the property $\{dA\} = DA$.

(iv) Let $A = (A, f) \in \mathcal{U}^c$, $x \in A$ be arbitrary. Then

(a) $SAx = \infty_1$ implies $\langle x \rangle_A \cap A^{\infty_1} \neq \emptyset$,

(b) $SAx = \infty_2$ implies $\langle x \rangle_A \cap A^{\infty_2} \neq \emptyset$,

(c) $SAx = \alpha \in \text{Ord}$ implies that $W(\alpha)$ is cofinal with $SA\langle x \rangle_A$,

(d) $SAx \in \text{Ord}$, $x \in \text{dom } f^n$, where $n \in \mathbb{N}$, implies $SAf^n x \geq SAx + n$.

(See [4], 2.17 (a), (b), 2.25 (b) and 2.26 (a).)

1.10. Lemma. Let $A = (A, f) \in \mathcal{U}^c$, $x \in A$ be such that $SAx \in \text{Ord}$. Let $\alpha \in \text{Ord}$. Then the following assertions hold:

(a) $SAx > \alpha$ iff there is $y \in \langle x \rangle_A$ such that $SAy \geq \alpha$.

(b) $\langle x \rangle_A - \bigcup_{\lambda \in W(\alpha)} A^\lambda \neq \emptyset$ iff $x \in A - \bigcup_{\lambda \in W(\alpha+1)} A^\lambda$.

Proof of (a). Let $y \in \langle x \rangle_A$ be such that $SAy \geq \alpha$. Then $SAx = SAy > SAy \geq \alpha$ by (iv) (d).

Let, on the other hand, $SAx > \alpha$. Then there is $y \in \langle x \rangle_A$ such that $SAy \geq \alpha$ by (iv) (c).

(b) is a consequence of (a).

1.11. Convention. Let $A = (A, f) \in \mathcal{U}^c$, $B = (B, g) \in \mathcal{U}^c$. If we speak about a d-map (d-injection) $F : A \rightarrow B$ we already mean a d-map (d-injection) with respect to SA, SB (see 1.1).

1.12. Definition. Let $A = (A, f) \in \mathcal{U}^c$, $B = (B, g) \in \mathcal{U}^c$. Then

(a) we put $H(A, B) = \{(x, x') \in A \times B; \text{ for each } n \in N, x \in \text{dom } f^n \text{ implies } x' \in \text{dom } g^n \text{ and } SAf^n x \leq SBg^n x'\}$,

(b) we define

$$(A, B) \in h\text{-Ad} \text{ iff } \begin{aligned} (1) & RB \neq 0 \text{ implies } RB \mid RA^*, \\ (2) & RB = 0 \text{ implies } H(A, B) \neq \emptyset; \end{aligned}$$

(c) we put $M_0(A, B) = \{(x, x') \in H(A, B); RB \neq 0 \text{ implies } x \in ZA \cup \bigcup_{i \in W(RB)} [dA]_A^{-i}\}$,

(d) supposing that $DA = \emptyset$ implies $RA = RB$ we define $(x, x') \in M(A, B)$ iff the following conditions hold: $(x, x') \in M_0(A, B)$ and if we define a map $F_0: [x]_A \rightarrow [x']_B$ such that $F_0 f^n x = g^n x'$ for each $f^n x \in [x]_A$ then there is a sequence of maps $\{F_n: [x]_A^{-n} \rightarrow [x']_B^{-n}; n \in N\}$ such that, for each $n \in N$ and each $y \in [x]_A^{-n}$, $F_{n+1} \mid \langle y \rangle_{A,x}$ is a d-injection and $F_{n+1} \langle y \rangle_{A,x} \subseteq \langle F_n y \rangle_{B,x'}$;

(e) we define

$$(A, B) \in m\text{-Ad} \text{ iff } \begin{aligned} (1) & DA = \emptyset \text{ implies } RA = RB, \\ (2) & M(A, B) \neq \emptyset. \end{aligned}$$

(v) $m\text{-Ad} \subseteq h\text{-Ad}$. (See [7], 3.17.)

(vi) Let $A, B \in \mathcal{U}^c$. Then there is a monomorphism of A into B if and only if $(A, B) \in m\text{-Ad}$. (See [7], 3.21.)

2. EXISTENCE OF MONOMORPHISMS

The characteristic condition of the existence of monomorphisms contained in (vi) is rather complicated because of the definition of $M(A, B)$ (1.12(d)). In the following special cases, it can be simplified.

2.1. Definition. We put $\mathcal{U}_0^c = \{A = (A, f) \in \mathcal{U}^c; |\langle x \rangle_A| < \aleph_0 \text{ for each } x \in A\}$.

2.2. Lemma. Let $A = (A, f) \in \mathcal{U}_0^c$. Then

(a) $\mathfrak{A}A \leq \omega_0$,

(b) $SAA \subseteq N \cup \{\infty_1, \infty_2\}$.

Proof of (a). Let, on the contrary, $\mathfrak{A}A > \omega_0$. Then there is $x \in A$ such that $SAx = \omega_0$ by 1.8(c). Further, $W(\omega_0)$ is cofinal with $SA\langle x \rangle_A$ by (iv) (c) and thus $|\langle x \rangle_A| \geq |SA\langle x \rangle_A| = \aleph_0$ which is a contradiction to $A \in \mathcal{U}_0^c$.

(b) follows immediately from (a).

2.3. Definition. Let $d, \bar{d} \notin (N - \{0\}) \cup \{\omega_0\}$. We put $N_1 = (N - \{0\}) \cup \{\omega_0, d, \bar{d}\}$, $N_2 = N - \{0\}$.

* $p \mid q$ for $p, q \in N$ means that p is a divisor of q .

(a) We define a relation

(α) \leq_1 on N_1 in this way: if $a, b \in N_1$ then we put $a \leq_1 b$ iff (1) $a, b \in N - \{0\}$, $a \leq b$ or (2) $a \in (N - \{0\}) \cup \{\omega_0\}$, $b = \omega_0$ or (3) $a \in (N - \{0\}) \cup \{d\}$, $b = d$ or (4) $b = \bar{d}$;

(β) \leq_2 on N_2 in this way: if $a, b \in N_2$ then we put $a \leq_2 b$ iff $b \mid a$.
(N_1, N_2 are ordered sets. Compare [5], 1.16, 1.17.)

(b) We put $C_0 = N_1 \oplus N_2$ (see 1.0) and we denote the order on C_0 by \leq_0 . Further, if $(N_1, a), (N_2, b) \in C_0$ (where $a \in N_1, b \in N_2$) are arbitrary then we put (for brevity) $a = (N_1, a), b = (N_2, b)$.

(c) If $a, b \in C_0$ are arbitrary then we put $a \leq b$ iff (1) $a \leq_0 b$ and (2) $b \in N_2$ implies $a \in N_1 - \{\omega_0, \bar{d}\}$ or $a = b$.

(d) Let $A \in \mathcal{U}_0^c$. Then we define

$$\chi A = \begin{cases} RA \in N_2 & \text{if } RA \neq 0, \\ \bar{d} \in N_1 & \text{if } RA = 0, KA \neq \emptyset, DA = \emptyset, \\ d \in N_1 & \text{if } RA = 0, KA \neq \emptyset, DA \neq \emptyset, \\ \emptyset A \in N_1 & \text{if } RA = 0, KA = \emptyset. \end{cases}$$

(vii) Let $A, B \in \mathcal{U}_0^c$. Then

(a) $DA = \emptyset$ iff $\chi A \in \{\omega_0, \bar{d}\} \cup N_2$,

(b) $(A, B) \in h\text{-Ad}$ implies $\chi A \leq_0 \chi B$.

(See [5], 1.12 and 1.23.)

2.4. Lemma. Let $A, B \in \mathcal{U}_0^c$, Then

(a) $\chi A \leq \chi B$ iff (1) $\chi A \leq_0 \chi B$ and (2) $DA = \emptyset$ implies $RA = RB$,

(b) $(A, B) \in m\text{-Ad}$ implies $\chi A \leq \chi B$.

Proof of (a). Let $\chi A \leq \chi B$. Then $\chi A \leq_0 \chi B$. Further, let $DA = \emptyset$. Then $\chi A \in \{\omega_0, \bar{d}\} \cup N_2$ by (vii) (a).

Now, if $RB = 0$ then $\chi B \in N_1$ which implies $\chi A \in N_1$, i.e. $RA = 0$ and we have $RA = RB$.

If $RB \neq 0$ then $\chi B = RB \in N_2$ which implies $\chi A \in N_1 - \{\omega_0, \bar{d}\}$ or $\chi B = \chi A$ by 2.3 (c). Since $\chi A \notin N_1 - \{\omega_0, \bar{d}\}$ we have $RB = \chi B = \chi A = RA$.

Let, on the other hand, $\chi A \leq_0 \chi B$ and let $DA = \emptyset$ implies $RA = RB$.

Let $\chi B \in N_2$. Then $RB \neq 0$. If $DA \neq \emptyset$ then $\chi A \in N_1 - \{\omega_0, \bar{d}\}$ by (vii) (a) and if $DA = \emptyset$ then $RA = RB \neq 0$ which implies $\chi A = RA = RB = \chi B$. It follows that $\chi A \leq \chi B$.

Proof of (b). Let $(A, B) \in m\text{-Ad}$. Then $\chi A \leq_0 \chi B$ by (v) and (vii) (b). Hence $\chi A \leq \chi B$ by 1.12 (e) and by (a) of this lemma.

2.5. Definition. Let $n \in N \cup \{\infty_1, \infty_2\}$ be arbitrary. Then we put

$$W^*(n) = \begin{cases} W(n+1) & \text{if } n \in N \\ W(n) & \text{if } n \in \{\infty_1, \infty_2\} \end{cases} \quad (\text{see 1.0}).$$

2.6. Lemma. Let $A = (A, f) \in \mathcal{U}_0^c$, $x \in A$, $n \in N \cup \{\infty_1, \infty_2\}$ be arbitrary. Then $\langle x \rangle_A - \bigcup_{i \in W(n)} A^i \neq \emptyset$ iff $x \in A - \bigcup_{i \in W^*(n)} A^i$.

Proof. Let $n \in N$. If $S_A x \in W(\mathcal{A})$ then the assertion holds by 1.10 (b). If $S_A x \in \{\infty_1, \infty_2\}$ then $x \in A^{\infty_1} \cup A^{\infty_2} \subseteq A - \bigcup_{i \in W^*(n)} A^i$ and $\langle x \rangle_A - \bigcup_{i \in W(n)} A^i \supseteq \langle x \rangle_A \cap (A^{\infty_1} \cup A^{\infty_2}) \neq \emptyset$ by (iv) (a), (b).

Let $n = \infty_1$. If $y \in \langle x \rangle_A - \bigcup_{i \in W(n)} A^i$ then $y \in A^{\infty_1} \cup A^{\infty_2}$ which implies $x = fy \in A^{\infty_1} \cup A^{\infty_2}$ by (iii) (a) and 1.8 (b). If, on the other hand, $x \in A - \bigcup_{i \in W^*(n)} A^i = A^{\infty_1} \cup A^{\infty_2}$ then $\langle x \rangle_A - \bigcup_{i \in W(n)} A^i \supseteq \langle x \rangle_A \cap (A^{\infty_1} \cup A^{\infty_2}) \neq \emptyset$ by (iv) (a), (b).

Similarly, the assertion holds for $n = \infty_2$ by (iii) (a) and (iv) (b).

2.7. Definition. Let $A = (A, f) \in \mathcal{U}_0^c$. If there is a sequence $\{c_n; n \in N \cup \{\infty_1, \infty_2\}\}$ such that for each $n \in N \cup \{\infty_1, \infty_2\}$ and each $x \in A - \bigcup_{i \in W^*(n)} A^i$, $|\langle x \rangle_A - \bigcup_{i \in W(n)} A^i| = c_n$ holds then A is called *homogeneous* (with the sequence $\{c_n\}$).

2.8. Definition. Let $A = (A, f) \in \mathcal{U}_0^c$, $B = (B, g) \in \mathcal{U}_0^c$, let B be homogeneous with the sequence $\{c_n\}$. Then B is said to be a *majorant* of A if $\chi_A \leq \chi_B$ and if, for arbitrary $n \in N \cup \{\infty_1, \infty_2\}$, $|\langle x \rangle_A - \bigcup_{i \in W(n)} A^i| \leq c_n$ holds for each $x \in A$.

2.9. Theorem. Let $A, B \in \mathcal{U}_0^c$, B be homogeneous. If $(A, B) \in m\text{-Ad}$ then B is a majorant of A .

Proof. $\chi_A \leq \chi_B$ by 2.4 (b).

Further, let B be homogeneous with the sequence $\{c_n\}$. Let $(x, x') \in M(A, B)$. Then $(x, x') \in M_0(A, B)$ and there is a sequence of maps $\{F_n : [x]_A^{-n} \rightarrow [x']_B^{-n}; n \in N\}$ such that for each $f^n x \in [x]_A$, $F_0 f^n x = g^n x'$ and for each $n \in N$ and each $y \in [x]_A^{-n}$, $F_{n+1} | \langle y \rangle_{A,x}$ is a d-injection and $F_{n+1} \langle y \rangle_{A,x} \subseteq \langle F_n y \rangle_{B,x'}$.

1. Now let $y \in A$ be arbitrary. Then is $n \in N$ such that $y \in [x]_A^{-n}$ by (i) (b). It will be shown that there is a d-injection of $\langle y \rangle_A$ into $\langle F_n y \rangle_B$.

1a. Let $x \in ZA$. Then $x' \in ZB$ because $(x, x') \in H(A, B)$.

If $\langle y \rangle_A = \langle y \rangle_{A,x}$ then $n \in N - \{0\}$ by 1.7 (e). Hence $F_n y \in [x']_B^{-n}$ and $\langle F_n y \rangle_B = \langle F_n y \rangle_{B,x'}$ by 1.7 (e). It follows that $F_{n+1} | \langle y \rangle_{A,x}$ is a d-injection of $\langle y \rangle_A$ into $\langle F_n y \rangle_B$.

If, by 1.7 (e), $\langle y \rangle_A = \langle y \rangle_{A,x} \cup \{f^{p-1}x\}$ then $n = 0$, $y = f^p x$; hence $F_0 y = F_0 f^p x = g^p x'$ and thus $\langle F_0 y \rangle_B = \langle F_0 y \rangle_{B,x'} \cup \{g^{p-1}x'\}$ by 1.7 (e). Further, $F_0 f^{p-1}x = g^{p-1}x'$ and we obtain that $F_1 | \langle y \rangle_{A,x} \cup F_0 | \{f^{p-1}x\}$ is a d-injection of $\langle y \rangle_A$ into $\langle F_0 y \rangle_B$.

1b. Let, on the other hand, $x \notin ZA$. Then $ZA = \emptyset$ because if we had $ZA \neq \emptyset$ then we should have $DA = \emptyset$ by (iii) (c) and $RB = RA = 0$ by 1.12 (e) which is a contradiction to $(x, x') \in M_0(A, B)$ by 1.12 (c). Thus,

$$\langle y \rangle_A = \begin{cases} \langle y \rangle_{A,x} & \text{if } y \in \{x\} \cup \bigcup_{n \in N - \{0\}} [x]_A^{-n} \\ \langle y \rangle_{A,x} \cup \{f^{p-1}x\} & \text{if } y = f^p x, \quad p \neq 0 \end{cases} \quad \text{by 1.7 (f)}.$$

1ba. Let now $\langle y \rangle_A = \langle y \rangle_{A,x}$. Then $F_{n+1} \langle y \rangle_A = F_{n+1} \langle y \rangle_{A,x} \subseteq \langle F_{n+1} y \rangle_{B,x'} \subseteq \langle F_n y \rangle_B$ and thus $F_{n+1} | \langle y \rangle_{A,x}$ is a d-injection of $\langle y \rangle_A$ into $\langle F_n y \rangle_B$.

1bb. Let $\langle y \rangle_A = \langle y \rangle_{A,x} \cup \{f^{p-1}x\}$ where $y = f^p x$, $p \neq 0$. Then $F_0 y = F_0 f^p x = g^p x'$.

If $x' \in ZB$ then $\langle F_0 y \rangle_B = \langle F_0 y \rangle_{B,x'} \cup \{g^{p-1}x'\}$ by 1.7 (e).

If $x' \notin ZB$ and if we put $p_0 = \min \{n \in N; g^n x' \in ZB\}$ then

$$\langle F_0 y \rangle_B = \begin{cases} \langle F_0 y \rangle_{B,x'} \cup \{g^{p-1}x'\} & \text{if } p \neq p_0 \\ \langle F_0 y \rangle_{B,x'} \cup \{g^{p-1}x', g^{RB+p-1}x'\} & \text{if } p = p_0 \end{cases} \quad \text{by 1.7 (f)}.$$

Since $(F_0 \cup F_1) \langle y \rangle_A = F_1 \langle y \rangle_{A,x} \cup \{F_0 f^{p-1}x\} \subseteq \langle F_0 y \rangle_{B,x'} \cup \{g^{p-1}x'\} \subseteq \langle F_0 y \rangle_B$ we obtain that $F_1 | \langle y \rangle_{A,x} \cup F_0 | \{f^{p-1}x\}$ is a d-injection of $\langle y \rangle_A$ into $\langle F_0 y \rangle_B$.

2. Let $y \in A$, $m \in N \cup \{\infty_1, \infty_2\}$ be arbitrary.

Let $y \in [x]_A^{-n}$. Then by 1 there exists a d-injection of $\langle y \rangle_A$ into $\langle F_n y \rangle_B$ which implies $|\langle y \rangle_A - \bigcup_{i \in W(m)} A^i| \leq |\langle F_n y \rangle_B - \bigcup_{i \in W(m)} B^i|$ by 1.2.

If $y \in A - \bigcup_{i \in W^*(m)} A^i$ then $SAy \notin W^*(m)$ and we have $SBF_n y \notin W^*(m)$ because $SBF_n y \geq SAy$. Thus $F_n y \in B - \bigcup_{i \in W^*(m)} B^i$ and we obtain $|\langle F_n y \rangle_B - \bigcup_{i \in W(m)} B^i| = c_m$ by 2.7. It follows that $|\langle y \rangle_A - \bigcup_{i \in W(m)} A^i| \leq c_m$.

Now if $y \in \bigcup_{i \in W^*(m)} A^i$ then $|\langle y \rangle_A - \bigcup_{i \in W(m)} A^i| = 0$ by 2.6; therefore $|\langle y \rangle_A - \bigcup_{i \in W(m)} A^i| \leq c_m$.

Consequently, B is a majorant of A .

2.10. Lemma. Let $A = (A, f) \in \mathcal{Q}_0^c$, $B = (B, g) \in \mathcal{Q}_0^c$ be such that $RA = RB$. Let $x \in DA \cup KA \cup ZA$, $(x, x') \in M_0(A, B)$ be such that $DA \neq \emptyset$ implies $x = dA$. Let $n \in N$, $y \in [x]_A^{-n}$, $y' \in [x']_B^{-n}$ be such that $n = 0$ implies $y = f^p x$, $y' = g^p x'$ for some $p \in N$. If $|\langle y \rangle_A - \bigcup_{i \in W(m)} A^i| \leq |\langle y' \rangle_B - \bigcup_{i \in W(m)} B^i|$ for each $m \in N \cup \{\infty_1, \infty_2\}$ then there is a d-injection of $\langle y \rangle_{A,x}$ into $\langle y' \rangle_{B,x'}$.

Proof. 1. Let $RA \neq 0$. Then $RB \neq 0$ and $(x, x') \in ZA \times ZB$ by 1.12 (c).

If $n = 0$ then $y = f^p x \in ZA$, $y' = g^p x' \in ZB$ where $p \in N$. We can suppose that $p \neq 0$. Thus $\langle y \rangle_A = \langle y \rangle_{A,x} \cup \{f^{p-1}x\}$, $\langle y' \rangle_B = \langle y' \rangle_{B,x'} \cup \{g^{p-1}x'\}$ by 1.7 (e) and since $SAf^{p-1}x = \infty_2 = SBg^{p-1}x'$ it follows that there is a d-injection of $\langle y \rangle_{A,x}$ into $\langle y' \rangle_{B,x'}$ by 1.3 and 1.2.

If $n > 0$ then $\langle y \rangle_A = \langle y \rangle_{A,x}$, $\langle y' \rangle_B = \langle y' \rangle_{B,x'}$ by 1.7 (e) and therefore there is a d-injection of $\langle y \rangle_{A,x}$ into $\langle y' \rangle_{B,x'}$ by 1.2.

2. Let $RA = 0$. Then $RB = 0$.

2a. Let $DA = \emptyset$. Then $x = dA$.

In this case, $\langle y \rangle_A = \langle y \rangle_{A,x}$ by 1.7 (f) because $y \in A = \{dA\} \cup \bigcup_{i \in N - \{0\}} [dA]_A^{-i}$

by (i) (b). Further, $y' \in \{x'\} \cup \bigcup_{i \in N - \{0\}} [x']_B^{-i}$ because $y \in [dA]_A$, $y' \in [x']_B$ imply

$y = f^0 dA$, $y' = g^0 x' = x'$ and since $x' \notin ZB$ we have $\langle y' \rangle_B = \langle y' \rangle_{B,x'}$ by 1.7 (f).

Therefore, there is a d-injection of $\langle y \rangle_{A,x}$ into $\langle y' \rangle_{B,x'}$ by 1.2.

2b. Let $DA = \emptyset$. Then $x \in KA$ which implies $[x]_A \subseteq KA$, $[x']_B \subseteq KB$ because $(x, x') \in H(A, B)$, $ZA = ZB = \emptyset$.

If $y \in \{x\} \cup \bigcup_{i \in N - \{0\}} [x]_A^{-i}$ then $\langle y \rangle_A = \langle y \rangle_{A,x}$ by 1.7 (f). Further, $y' \in \{x'\} \cup \bigcup_{i \in N - \{0\}} [x']_B^{-i}$ because $y \in [x]_A$, $y' \in [x']_B$ and we complete the proof again by 1.2.

If $y = f^p x$ where $p > 0$ then $y' = g^p x'$ and thus $\langle y \rangle_A = \langle y \rangle_{A,x} \cup \{f^{p-1}x\}$, $\langle y' \rangle_B = \langle y' \rangle_{B,x'} \cup \{g^{p-1}x'\}$ by 1.7 (f) because $ZA = ZB = \emptyset$. Since $SAf^{p-1}x = \infty_1 = SBg^{p-1}x'$ there is a d-injection of $\langle y \rangle_{A,x}$ into $\langle y' \rangle_{B,x'}$ by 1.3 and 1.2.

2.11. Theorem. Let $A, B \in \mathcal{U}_0^c$ be such that A has the kernel, B is homogeneous and $DA \neq \emptyset$ implies $RB = 0$. If B is a majorant of A then $(A, B) \in m\text{-Ad}$.

Proof. Let $A = (A, f)$, $B = (B, g)$ and let B be homogeneous with the sequence $\{c_n; n \in N \cup \{\infty_1, \infty_2\}\}$. Since $\chi A \preceq \chi B$ we obtain that $DA = \emptyset$ implies $RA = RB$ by 2.4 (a). Further, we take $x \in DA \cup KA \cup ZA$ such that $RA \neq 0$ implies $x \in ZA$ and $DA \neq \emptyset$ implies $x = dA$.

If $RA \neq 0$ then $RB = RA \neq 0$ by (iii) (c) and 1.12 (e) and we take $x' \in ZB$. Then $(x, x') \in H(A, B)$ and further $(x, x') \in M_0(A, B)$ by 1.12 (c).

Let now $RA = 0$. Then $RB = 0$ because $RB \neq 0$ implies $DA = \emptyset$ and $RA = RB$ which is a contradiction to $RA = 0$. Since $\chi A \preceq \chi B$ there is $x' \in B$ such that $SAx \preceq SBx'$ by 2.3 (d), (c) and (iv) (e). If $x = dA$ then directly $(x, x') \in H(A, B) = M_0(A, B)$ by 1.12 (a), (c) because $ZA = ZB = \emptyset$ and it follows that $(x, x') \in H(A, B) = M_0(A, B)$.

We define a d-map $F_0 : [x]_A \rightarrow [x']_B$ such that $F_0 f^n x = g^n x'$ for each $f^n x \in [x]_A$ (see 1.12 (a)). Let $n \in N$ be arbitrary. Suppose that we have defined a d-map $F_n : [x]_A^{-n} \rightarrow [x']_B^{-n}$. It will be shown that there is a map $F_{n+1} : [x]_A^{-(n+1)} \rightarrow [x']_B^{-(n+1)}$ such that for each $y \in [x]_A^{-n}$, $F_{n+1} \mid \langle y \rangle_{A,x}$ is a d-injection and $F_{n+1} \langle y \rangle_{A,x} \subseteq \langle F_n y \rangle_{B,x'}$.

Let $y \in [x]_{\mathbf{A}}^{-n}$ be arbitrary. Since \mathbf{B} is a majorant of \mathbf{A} we have $|\langle y \rangle_{\mathbf{A}} - \bigcup_{i \in W(m)} A^i| \leq c_m$ for each $m \in N \cup \{\infty_1, \infty_2\}$. Further, $SAy \leq SBF_n y$ because F_n is a d-map.

Let $m \in N \cup \{\infty_1, \infty_2\}$ be arbitrary.

If $F_n y \in B - \bigcup_{i \in W^*(m)} B^i$ then $|\langle F_n y \rangle_{\mathbf{B}} - \bigcup_{i \in W(m)} B^i| = c_m$ because \mathbf{B} is homogeneous.

Thus $|\langle y \rangle_{\mathbf{A}} - \bigcup_{i \in W(m)} A^i| \leq |\langle F_n y \rangle_{\mathbf{B}} - \bigcup_{i \in W(m)} B^i|$.

If $F_n y \in \bigcup_{i \in W^*(m)} B^i$ then $SBF_n y \in W^*(m)$ which implies $SAy \in W^*(m)$. It follows that $|\langle y \rangle_{\mathbf{A}} - \bigcup_{i \in W(m)} A^i| = 0$ by 2.6. Hence $|\langle y \rangle_{\mathbf{A}} - \bigcup_{i \in W(m)} A^i| = 0 = |\langle F_n y \rangle_{\mathbf{B}} - \bigcup_{i \in W(m)} B^i|$.

Thus for each $m \in N \cup \{\infty_1, \infty_2\}$, $|\langle y \rangle_{\mathbf{A}} - \bigcup_{i \in W(m)} A^i| \leq |\langle F_n y \rangle_{\mathbf{B}} - \bigcup_{i \in W(m)} B^i|$ and we obtain that there is a d-injection $F_{n+1}^y : \langle y \rangle_{\mathbf{A}, x} \rightarrow \langle F_n y \rangle_{\mathbf{B}, x'}$ by 2.10.

If we put $F_{n+1} = \bigcup_{y \in [x]_{\mathbf{A}}^{-n}} F_{n+1}^y$ then F_{n+1} is the map sought for.

Consequently, $(x, x') \in M(\mathbf{A}, \mathbf{B})$ and finally, $(\mathbf{A}, \mathbf{B}) \in m\text{-Ad}$.

2.12. Theorem. Let $\mathbf{A}, \mathbf{B} \in \mathcal{U}_0^c$ be such that \mathbf{A} has the kernel, \mathbf{B} is homogeneous and $D\mathbf{A} \neq \emptyset$ implies $R\mathbf{B} = 0$. Then the following assertions are equivalent:

(α) There is a monomorphism of \mathbf{A} into \mathbf{B} .

(β) $(\mathbf{A}, \mathbf{B}) \in m\text{-Ad}$.

(γ) \mathbf{B} is a majorant of \mathbf{A} .

The assertion is a consequence of (vi), 2.9 and 2.11.

2.13. Corollary. Let $\mathbf{A} \in \mathcal{U}^c$ be finite and complete, $\mathbf{B} \in \mathcal{U}_0^c$ homogeneous. Then the following assertions are equivalent:

(α) There is a monomorphism of \mathbf{A} into \mathbf{B} .

(β) $(\mathbf{A}, \mathbf{B}) \in m\text{-Ad}$.

(γ) \mathbf{B} is a majorant of \mathbf{A} .

Indeed, if \mathbf{A} is finite then $\mathbf{A} \in \mathcal{U}_0^c$ and it has the kernel.

Bibliography

- [1] *M. Novotný*: Sur un problème de la théorie des applications, Publ. Fac. Sci. Univ. Masaryk, No. 344 (1953), 53–64.
- [2] *M. Novotný*: Über Abbildungen von Mengen, Pac. J. Math., 13 (1963), 1359–1369.
- [3] *O. Kopeček, M. Novotný*: On some invariants of unary algebras, Czech. Math. J., 24 (99) (1974), 219–246.

- [4] *O. Kopeček*: Homomorphisms of partial unary algebras, *Czech. Math. J.* 26 (101) (1976), 108—127.
- [5] *O. Kopeček*: The category of connected partial unary algebras, *Czech. Math. Journal* 27 (102) (1977), 415—423.
- [6] *O. Kopeček*: Construction of all machine homomorphisms, *Bull. Acad. Polon. Sci., Sér. Sci. Math. Astronom. Phys.* 24 (1976), 655—658.
- [7] *O. Kopeček*: Monomorphisms of partial unary algebras, to appear in *Czech. Math. J.*
- [8] *W. Stucky, H. Walter*: Minimal linear realizations of autonomous automata, *Information and Control* 16 (1970), 66—84.

Author's address: 662 95 Brno, Janáčkovo nám. 2a, ČSSR (Universita J. E. Purkyně).