

Bedřich Pondělíček

Note on homomorphisms of direct products of algebras

Czechoslovak Mathematical Journal, Vol. 29 (1979), No. 3, 500–501

Persistent URL: <http://dml.cz/dmlcz/101629>

Terms of use:

© Institute of Mathematics AS CR, 1979

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

NOTE ON HOMOMORPHISMS OF DIRECT PRODUCTS OF ALGEBRAS

BEDŘICH PONDĚLÍČEK, Praha

(Received June 2, 1978)

Let A be a non-void set and F a set of (algebraic) operations on A . An algebra (A, F) is said to be *without zero-divisors* if

(i) there exist $0 \in A$ and $\oplus \in F$ (where $\text{ar } \oplus = 2$) such that $a \oplus 0 = a = 0 \oplus a$ for each $a \in A$ and

(ii) at least one $\omega \in F$ (where $\omega \neq \oplus$) is regular on (A, F) , i.e. $\text{ar } \omega = n \geq 2$ and for each $a_1, \dots, a_n \in A$ we have $a_1 \dots a_n \omega = 0$ iff $a_i = 0$ for at least one $i \in \{1, \dots, n\}$.

The element 0 is called a *zero* of (A, F) .

I. CHAJDA in [1] has investigated homomorphisms of algebras, which are direct products of algebras without zero-divisors. In this note we shall show that in Theorem 9 of [1] and in its Corollary the author omits the following assumption:

(iii) $0 \dots 0\omega = 0$ for arbitrary $\omega \in F$.

Let A, B be algebras of the same type. The algebras A, B are called *r-similar* if they are without zero-divisors and have the same set of regular operations. If $f(0) = 0$ for each $f \in \text{Hom}(A, B)$, then the r-similar algebras A, B are said to be *super similar*. See [1].

Remark 1. The following example shows that there exist r-similar algebras A, B of the same type such that the zero mapping $o : A \rightarrow \{0\} \subset B$ is not a homomorphism of A into B . See Notation, p. 167, [1].

Example 1. By I we denote the set of all integers. Put $a \oplus b = a + b$, $a \circ b = ab$ and $a * b = 1$ for every $a, b \in I$. It is clear that 0 is a zero of the algebra $\mathcal{Z} = (I, F)$, where $F = \{\oplus, \circ, *\}$; \oplus fulfils (i), \circ fulfils (ii). This implies that the algebra \mathcal{Z} is without zero-divisors and so \mathcal{Z}, \mathcal{Z} are r-similar.

Now we shall show that $\text{Hom}(\mathcal{Z}, \mathcal{Z}) = \{\text{id}_I\}$.

Indeed, if $\varphi \in \text{Hom}(\mathcal{Z}, \mathcal{Z})$, then $\varphi(1) = \varphi(1 * 1) = \varphi(1) * \varphi(1) = 1$ and so we can prove by induction that $\varphi(n) = n$ for every positive integer n . It is clear that $\varphi(0) = 0$ and so $\varphi(-n) = -\varphi(n) = -n$.

Remark 2. The following example shows that Theorem 9 [1] is not true.

Example 2. It follows from Example 1 that the algebras \mathcal{L}, \mathcal{L} are super similar. By h we denote the projection of $\mathcal{L} \times \mathcal{L}$ onto the first factor \mathcal{L} . It is clear that $h \in \text{Hom}(\mathcal{L} \times \mathcal{L}, \mathcal{L})$.

Now we shall show that there exists no matrix representing h .

On the contrary, let us assume that h is represented by a matrix $H = \|h_{i1}\|$, where $h_{i1} \in \text{Hom}(\mathcal{L}, \mathcal{L})$ and $i = 1, 2$. It follows from Example 1 that $h_{i1} = \text{id}_{\mathcal{L}}$ and so $0 = h(0, 1) = h_{11}(0) \oplus h_{21}(1) = 0 + 1 = 1$, which is a contradiction.

Remark 3. The following example shows that Corollary to Theorem 9 [1] is false.

Example 3. Let $s = \text{card Hom}(\mathcal{L} \times \mathcal{L}, \mathcal{L})$, where \mathcal{L} is the same as in Example 1 and 2. Since both projections of $\mathcal{L} \times \mathcal{L}$ onto \mathcal{L} are homomorphisms, we have $s \geq 2$. On the other hand, it follows from Example 1 that $\text{card Hom}(\mathcal{L}, \mathcal{L}) = 1$ and so

$$s \neq 1 = \prod_{j=1}^m \left(1 + \sum_{i=1}^n (p_{ij} - 1)\right), \text{ where } m = 1, n = 2 \text{ and } p_{11} = p_{21} = 1.$$

Remark 4. Let A_i, B_j be super similar algebras for $i = 1, \dots, n; j = 1, \dots, m$ and $A = \prod_{i=1}^n A_i, B = \prod_{j=1}^m B_j$. If we define a matrix $H = \|h_{ij}\|$ representing a mapping h of A into B such that either $h_{ij} \in \text{Hom}(A_i, B_j)$ or h_{ij} is a zero mapping of A_i into B_j , then h need not be a homomorphism nor a zero mapping. Compare with Theorem 8 of [1].

Example 4. Let h be a mapping of \mathcal{L} into $\mathcal{L} \times \mathcal{L}$ (see Examples 1 and 2) represented by a matrix $H = \|h_{ij}\|$, where $j = 1, 2$ and $h_{11} = \text{id}_{\mathcal{L}}, h_{12} = 0$. Evidently $h(1) = (1, 0) \neq (0, 0)$ and so h is no zero mapping. We shall show that h is no homomorphism. On the contrary, let us suppose that $h \in \text{Hom}(\mathcal{L}, \mathcal{L} \times \mathcal{L})$. Then $(1, 0) = h(1) = h(1 * 1) = h(1) * h(1) = (1, 0) * (1, 0) = (1 * 1, 0 * 0) = (1, 1)$. a contradiction.

References

- [1] I. Chajda: Homomorphisms of direct products of algebras. Czech. Math. J. 28 (1978), 155–170.

Author's address: 166 27 Praha 6, Suchbátarova 2, ČSSR (České vysoké učení technické).