

Samuel Jezný; Marián Trenkler  
Characterization of magic graphs

*Czechoslovak Mathematical Journal*, Vol. 33 (1983), No. 3, 435–438

Persistent URL: <http://dml.cz/dmlcz/101893>

## Terms of use:

© Institute of Mathematics AS CR, 1983

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

CHARACTERIZATION OF MAGIC GRAPHS

SAMUEL JEZŇY and MARIÁN TRENKLER, Košice

(Received January 22, 1982)

I. INTRODUCTION

We shall consider a non-orientable finite graph  $G = [V(G), E(G)]$  without loops, multiple edges or isolated vertices. If there exists a mapping  $f$  from the set of edges  $E(G)$  into positive real numbers such that

$$(i) f(e_i) \neq f(e_j) \text{ for all } e_i \neq e_j; e_i, e_j \in E(G),$$

$$(ii) \sum_{e \in E(G)} \eta(v, e) f(e) = r \text{ for all } v \in V(G),$$

$$\text{where } \eta(v, e) = \begin{cases} 1 & \text{when vertex } v \text{ and edge } e \text{ are incident} \\ 0 & \text{in the opposite case,} \end{cases}$$

then the graph  $G$  is called *magic*. The mapping  $f$  is called a *labelling* of  $G$  and the value  $r$  is the index of the label  $f$ . We say that a graph  $G$  is *semimagic* if there exists a mapping  $f$  into positive real number which satisfies only the condition (ii). If the semimagic graph  $G$  has a label with the index  $r$  we shall say that  $G$  has index  $r$ .

To study magic graphs was suggested by J. Sedláček [3]. Some sufficient conditions for the existence of magic graphs are established in [2], [4] and [5]. A characterization of regular magic graphs in terms of circuits is given by M. Doob [1]. J. Mühlbacher [2] used matrix theory to prove two necessary conditions for the existence of magic graph. These conditions are weaker than that of theorem 2 of this paper.

First we shall formulate several necessary definitions.

A subgraph  $F = [V(F), E(F)]$  of the graph  $G = [V(G), E(G)]$  is called a *factor* of  $G$  if the sets  $V(G)$  and  $V(F)$  are the same. A factor  $F$  is a  $(1-2)$ -factor of  $G$  if each of its components is a regular graph of degree one or two. By the symbol  $F^1$ , resp.  $F^2$  we denote the subgraph of  $F$  which consists of all isolated edges, or of all circuits of  $F$  and the necessary vertices, respectively. We say that a  $(1-2)$ -factor separates the edge  $e_1$  and  $e_2$ , if at least one of them belongs to  $F$  and neither  $F^1$  nor  $F^2$  contains both of them.

The aim of this paper is to characterize all magic graph using the notion of separating edges by a  $(1-2)$ -factor.

## II. SEMIMAGIC GRAPHS

In this part we state some results about semimagic graphs which we shall use to prove the main result.

**Lemma 1.** *If  $G$  is a semimagic graph with the index  $r$ , then*

- a) *each isolated edge of  $G$  has the label  $r$ ,*
- b) *a connected part of  $G$  having more than one edge contains no vertex of degree one.*

The proofs of these statements follow from the definition of a semimagic graph.

**Lemma 2.** *Let a semimagic graph  $G$  contain an even circuit  $C$ , then there exists a semimagic factor  $H$  of  $G$  which does not contain all edges of  $C$ .*

*Proof.* Let  $f$  be a semimagic labelling of  $G$  and let  $m = \min \{f(e); e \in E(C)\}$ . We denote the edges of  $C$  by  $e_1, e_2, \dots, e_{2n}$  and suppose that  $f(e_1) = m$ . We define a new labelling  $h$  of  $G$ :


$$\begin{aligned} h(e_{2i-1}) &= f(e_{2i-1}) - m, \\ h(e_{2i}) &= f(e_{2i}) + m \text{ for } i = 1, 2, \dots, n, \\ h(e_j) &= f(e_j) \text{ for all } e_j \notin E(C). \end{aligned}$$

Obviously  $h(e_i) = 0$ . By omitting all edges with  $h(e) = 0$  from  $G$  we obtain a factor which does not contain all edges of the circuit  $C$  and has the same index as the graph  $G$ .

The graph  $D$  is called a *dumbbell* if it consists of two odd circuits  $C_1$  and  $C_2$  without common vertices joined by a path  $P$  or if it consists only of two odd circuits  $C_1$  and  $C_2$  with only one common vertex.

**Lemma 3.** *Let a semimagic graph  $G$  contain as a subgraph a dumbbell  $D$ , then there exists a semimagic factor  $H$  of  $G$  which does not contain all edges of the subgraph  $D$ .*

*Proof.* Let  $f$  be a semimagic labelling of  $G$  with a dumbbell  $D$  which consists of two circuits  $C_1, C_2$  and a path  $P$  or only of two circuits  $C_1, C_2$ . We denote  $m = \min \{m_1, m_2\}$  where  $m_1 = \min \{f(e); e \in E(C_1) \cup E(C_2)\}$  and  $m_2 = 1/2 \min \{f(e);$



$e \in E(P)\}$ . Let  $e'$  be an edge of  $D$  such that  $f(e') = m$ . We define an auxiliary labelling  $p$ . The edges of  $C_i$  have alternating values 1 and  $-1$  and the edges of  $P$  the values 2 and  $-2$  such that the sum at each vertex is zero, and the value of the edge  $e'$  is negative.

All the other edges of  $G$  have value 0. We consider the labelling

$$h(e) = f(e) + m p(e) \quad \text{for all } e \in E(G).$$

All edges having  $h(e) > 0$  form a semimagic factor  $H$  of  $G$  which has the same index as  $G$ .

From the lemmas 2 and 3 it follows:

**Lemma 4.** *If  $G$  is a semimagic graph, then there exists a semimagic  $(1-2)$ -factor  $F$  of  $G$  with the same index.*

**Lemma 5.** *If  $G$  is a semimagic graph, then every edge  $e'$  of  $G$  is contained in a  $(1-2)$ -factor.*

*Proof.* Let  $e'$  be an arbitrary edge of  $G$  and  $F$  some  $(1-2)$ -factor of  $G$ . There are two possible cases: either  $e' \in E(F)$  or  $e' \notin E(F)$ . We must consider only the second case.

Let  $q$  be an auxiliary labelling such that

$$\begin{aligned} q(e) &= 2 & \text{for all } e \in E(F^1), \\ q(e) &= 1 & \text{for all } e \in E(F^2), \\ q(e) &= 0 & \text{for all } e \notin E(F), \end{aligned}$$

and

$$m = \min \{f(e)/q(e) : e \in E(F)\}.$$

We consider a new labelling

$$h(e) = f(e) - m q(e) \quad \text{for all } e \in E(G).$$

Omitting from the graph  $G$  all edges for which  $h(e) = 0$  we obtain a semimagic factor  $H$  which contains the edge  $e'$ . Let  $F'$  be a  $(1-2)$ -factor of  $H$ . (Note that  $F'$  is also a  $(1-2)$ -factor of  $G$ .) If  $e' \notin E(F')$  we repeat the construction described after. By a finite number of repetitions we obtain a  $(1-2)$ -factor of  $G$  which contains the edge  $e'$ .

**Lemma 6.** *If every edge of  $G$  belongs to a  $(1-2)$ -factor, then  $G$  is semimagic.*

*Proof.* A semimagic labelling of  $G$  is obtained by a finite number of repetitions of the following construction.

Let  $f$  be a labelling with nonnegative numbers such that the sum of the labels of edges incident with each vertex is the same. (Note that every graph has such a labelling.) Let  $e$  be an edge with  $f(e) = 0$  and  $F$  one  $(1-2)$ -factor such that  $e \in E(F)$ . We define a new labelling

$$\begin{aligned} h(e) &= f(e) + 2m & \text{for all } e \in E(F^1), \\ h(e) &= f(e) + m & \text{for all } e \in E(F^2), \\ h(e) &= f(e) & \text{for all } e \notin E(F), \end{aligned}$$

where  $m = \max \{f(e) : e \in E(G)\} + 1$ .

From the previous lemmas it follows:

**Theorem 1.** *The graph  $G$  is semimagic if and only if every edge is contained in a  $(1-2)$ -factor.*

### III. CHARACTERIZATION OF MAGIC GRAPH

**Lemma 7.** *If every couple of edges  $e_1, e_2$  of a semimagic graph  $G$  is separated by a  $(1-2)$ -factor, then  $G$  is magic.*

*Proof.* Let  $f$  be a semimagic labelling of  $G$ . If  $f(e_1) \neq f(e_2)$  for all couples of edges  $e_1, e_2$ , then  $G$  is magic. In the opposite case we choose a  $(1-2)$ -factor  $F$  which separates  $e_1$  and  $e_2$  and define a new labelling  $h$  as in the proof of lemma 6. After a finite number of repetitions of the previous step we obtain a magic graph.

The previous lemmas yield the proof of our main result.

**Theorem 2.** *A graph  $G$  is magic if and only if (i) every edge of  $G$  belongs to a  $(1-2)$ -factor, and (ii) every couple of edges  $e_1, e_2$  is separated by a  $(1-2)$ -factor.*

**Consequence.** *If  $G$  is magic graph then there exists a magic labelling of  $G$  with positive integers.*

#### References

- [1] *M. Doob*: Characterizations of Regular Magic Graphs, *J. Combinatorial Theory, Ser. B*, 25, 94–104 (1978).
- [2] *J. Mühlbacher*: Magische Quadrate und ihre Verallgemeinerung: ein graphentheoretisches Problem, *Graph, Data Structures, Algorithms*, Hansen Verlag 1979, München.
- [3] *J. Sedláček*: Problem 27, in "Theory of Graphs and Its Applications", *Proc. Symp. Smolenice* 1963, 163–167.
- [4] *J. Sedláček*: On magic graphs, *Math. Slov.* 26 (1976), 329–335.
- [5] *B. M. Stewart*: Magic Graphs, *Canad. J. Math.*, 18 (1966), 1031–1059.

*Authors' address*: 041 54 Košice, Jesenná 5, ČSSR (Univerzita P. J. Šafárika).