

William F. Trench

Linear perturbations of a constant coefficient differential equation subject to mild integral smallness conditions

Czechoslovak Mathematical Journal, Vol. 36 (1986), No. 4, 623–633

Persistent URL: <http://dml.cz/dmlcz/102121>

Terms of use:

© Institute of Mathematics AS CR, 1986

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

LINEAR PERTURBATIONS OF A CONSTANT COEFFICIENT
DIFFERENTIAL EQUATION SUBJECT TO MILD INTEGRAL
SMALLNESS CONDITIONS

WILLIAM F. TRENCH, Philadelphia

(Received August 19, 1985)

1. INTRODUCTION

We consider the equation

$$(1) \quad x^{(n)} + [a_1 + p_1(t)] x^{(n-1)} + \dots + [a_n + p_n(t)] x = f(t), \quad t > 0,$$

assuming throughout that a_1, \dots, a_n are complex constants and p_1, \dots, p_n, f are complex-valued and continuous on $(0, \infty)$. We give conditions implying that if λ_m is a simple zero of

$$Q(\lambda) = \lambda^n + a_1 \lambda^{n-1} + \dots + a_n,$$

then (1) has a solution x_m which behaves asymptotically like $ce^{\lambda_m t}$.

We use “ o ” and “ O ” in the standard way to denote behavior as $t \rightarrow \infty$. The following theorem is due to Dunkel [1]; see also Hartman [2; Thm 17.2, p. 316].

Theorem 1. *Suppose that*

$$(2) \quad \int_0^\infty t^q |p_k(t)| dt < \infty, \quad 1 \leq k \leq n,$$

for some $q \geq 0$, that λ_m is a simple zero of $Q(\lambda)$, and that if λ_j is any other zero of $Q(\lambda)$, then $\operatorname{Re}(\lambda_j - \lambda_m) \neq 0$. Then the equation

$$(3) \quad x^{(n)} + [a_1 + p_1(t)] x^{(n-1)} + \dots + [a_n + p_n(t)] x = 0, \quad t > 0,$$

has a solution x_m such that

$$(4) \quad x_m^{(r)}(t) = (\lambda_m^r + o(t^{-q})) e^{\lambda_m t}, \quad 0 \leq r \leq n-1.$$

Šimša [3] has recently given conditions which imply that (3) has a fundamental system x_1, x_2, \dots, x_n which satisfies (4) for $1 \leq m \leq n$. His proof easily implies the following result for a given m in $\{1, \dots, n\}$.

Theorem 2. *Suppose that $Q(\lambda)$ has simple roots $\lambda_j = \mu_j + i\nu_j$ ($1 \leq j \leq n$), and*

that

$$(5) \quad \int_0^\infty |p_1(t)| dt < \infty.$$

Suppose also that for some integer m ($1 \leq m \leq n$) and nonnegative constants q and ϱ , the integrals

$$(6) \quad \int_0^\infty t^q p_k(t) e^{[\varrho + i(\nu_m - \nu_j)]t} dt, \quad 1 \leq k \leq n$$

converge (perhaps conditionally), for j in

$$(7) \quad S = \{j \mid j = m \text{ or } \operatorname{Re}(\lambda_j - \lambda_m) + \varrho = 0\}.$$

Finally, suppose that at least one of the following is true:

(i) $\varrho > 0$; (ii) $q \geq 1$; or

$$(8) \quad \int_0^\infty t^{-q} \left| \int_t^\infty s^q p_k(s) ds \right| dt < \infty, \quad 2 \leq k \leq n.$$

Then (3) has a solution x_m such that

$$x_m^{(r)}(t) = (\lambda_m^r + o(e^{-\alpha t} t^{-q})) e^{2mt}, \quad 0 \leq r \leq n-1.$$

Except for the assumption that all zeros of $Q(\lambda)$ be distinct (which is required for Šimša's stronger result), Theorem 2 is a considerable extension of Theorem 1, since under assumptions (i) and (ii) there are no integral smallness conditions on p_2, \dots, p_n which require absolute convergence, and (8) is weaker than (2) for $2 \leq k \leq n$, while (5) is weaker than (2) with $k = 1$ and $q > 0$.

Šimša [4] has given an example showing that some additional assumption such as (8) must be imposed to obtain the conclusion of Theorem 2 with $\varrho = 0$ and $0 \leq q < 1$; however, we will show below that (8) can be weakened. (Very recently, Šimša [5] has obtained results for this case without assuming (8); however, they do not seem to be directly related to the results that we present here.)

2. THE MAIN THEOREM

It is convenient to state the following assumption separately from our main theorem.

Assumption A. Let

$$(9) \quad Q(\lambda) = (\lambda - \lambda_1)^{d_1} \dots (\lambda - \lambda_L)^{d_L},$$

where

$$\lambda_j = \mu_j + i\nu_j, \quad 1 \leq j \leq L,$$

are distinct, and

$$\mu_1 \leq \mu_2 \leq \dots \leq \mu_L.$$

Let m be a fixed integer in $\{1, \dots, L\}$ such that $d_m = 1$. Let ϱ be a nonnegative constant, and suppose that $d_j = 1$ if

$$(10) \quad \mu_j - \mu_m + \varrho = 0.$$

Let N be the unique integer in $\{1, \dots, L\}$ such that

$$(11) \quad \mu_j - \mu_m + \varrho < 0, \quad 1 \leq j \leq N - 1$$

(which is vacuous if $N = 1$) and

$$(12) \quad \mu_j - \mu_m + \varrho \geq 0, \quad N \leq j \leq L.$$

Let ϕ be positive and nonincreasing on $(0, \infty)$ and, if $N \geq 2$, let $e^{\alpha t} \phi(t)$ be non-decreasing for t sufficiently large (say $t \geq T_1$) for some α such that

$$0 < \alpha < \mu_m - \mu_{N-1} - \varrho.$$

Finally, let c be a given constant, and define

$$(13) \quad g(t) = f(t) - ce^{\lambda_m t} \sum_{k=1}^n \lambda_m^{n-k} p_k(t).$$

Notice that λ_j need not be a simple root of $Q(\lambda)$ except for those j 's (if any) that satisfy (10); thus, if $\varrho > 0$, then λ_m itself need not be simple.

Improper integrals occurring in hypotheses below are assumed to converge, and the convergence may be conditional, except, of course, where the integrands are necessarily nonnegative.

The following is our main result.

Theorem 3. *Suppose that Assumption A holds and*

$$(14) \quad \int_t^\infty g(s) e^{(e^{-\mu_m - i\nu_j})s} ds = O(\phi(t))$$

(see (13)) for j in S (see (7)). Suppose also that

$$(15) \quad \int_t^\infty |p_1(s)| \phi(s) ds = o(\phi(t))$$

and

$$(16) \quad \int_t^\infty \left| \int_s^\infty p_k(\lambda) d\lambda \right| \phi(s) ds = o(\phi(t)), \quad 2 \leq k \leq n.$$

Then (1) has a solution x_m such that

$$(17) \quad x_m^{(r)}(t) = (c\lambda_m^r + O(e^{-\varrho t} \phi(t))) e^{\lambda_m t}, \quad 0 \leq r \leq n - 1.$$

Moreover, if "O" can be replaced by "o" in (14), then

$$(18) \quad x_m^{(r)}(t) = (c\lambda_m^r + o(e^{-\varrho t} \phi(t))) e^{\lambda_m t}, \quad 0 \leq r \leq n - 1.$$

3. PROOF OF THE MAIN THEOREM

Following Šimša [3], we use the Banach contraction principle to prove Theorem 3. It is convenient to introduce the new dependent variable

$$h(t) = x(t) - ce^{\lambda mt},$$

in terms of which (1) becomes

$$(19) \quad Q(D)h = g - Mh$$

(see (9) and (13)), with

$$(20) \quad Mh = \sum_{k=1}^n p_k h^{(n-k)}.$$

Now suppose that $t_0 \geq 0$ and let $B(t_0)$ be the Banach space of functions h in $C^{(n-1)}[t_0, \infty)$ such that

$$h^{(r)}(t) = O(e^{(\mu_m - \sigma)t} \phi(t)), \quad 0 \leq r \leq n-1,$$

with norm

$$(21) \quad \|h\| = \sup_{t \geq t_0} \{e^{(\sigma - \mu_m)t} (\phi(t))^{-1} \sum_{r=0}^{n-1} |h^{(r)}(t)|\}.$$

Clearly, if (19) has a solution h_m in $B(t_0)$, then the function

$$(22) \quad x_m(t) = ce^{\lambda mt} + h_m(t)$$

satisfies (1) on $[t_0, \infty)$ (and can be continued as a solution of (1) over $(0, \infty)$), and has the asymptotic behavior (17). We will now define a transformation which we will show to be a contraction of $B(t_0)$ if t_0 is sufficiently large, whose fixed point (function) h_m satisfies (19) on $[t_0, \infty)$.

To this end, let $A_1(t), \dots, A_L(t)$ be the unique polynomials such that $\deg A_j < d_j$ ($1 \leq j \leq L$) and

$$\sum_{j=1}^L [A_j(t) e^{\lambda_j t}]^{(r)} \Big|_{t=0} = \delta_{r, n-1}, \quad 0 \leq r \leq n-1,$$

and define the associated polynomials

$$A_{jr}(t) = e^{-\lambda_j t} [A_j(t) e^{\lambda_j t}]^{(r)}, \quad 0 \leq r \leq n-1, \quad 1 \leq j \leq L.$$

Then

$$\deg A_{jr} = \deg A_j < d_j, \quad 0 \leq r \leq n-1, \quad 1 \leq j \leq L,$$

and the standard variation of parameters argument shows that if $w \in C[t_0, \infty)$ and

$$(23) \quad v'(t; w) = \sum_{j=1}^{N-1} \int_{t_0}^t A_j(t-\tau) e^{\lambda_j(t-\tau)} w(\tau) d\tau - \\ - \sum_{j=N}^L \int_t^\infty A_j(t-\tau) e^{\lambda_j(t-\tau)} w(\tau) d\tau$$

(where the first sum is vacuous if $N = 1$), then

$$(24) \quad v^{(r)}(t; w) = \sum_{j=1}^{N-1} \int_{t_0}^t A_{jr}(t-\tau) e^{\lambda_j(t-\tau)} w(\tau) d\tau - \\ - \sum_{j=N}^L \int_t^\infty A_{jr}(t-\tau) e^{\lambda_j(t-\tau)} w(\tau) d\tau, \quad 0 \leq r \leq n-1,$$

and

$$(25) \quad Q(D) v(t; w) = w,$$

provided that the improper integrals in (23) and (24) converge. This prompts us to consider the transformation \mathcal{T} defined by

$$\mathcal{T}h = G - \mathcal{L}h,$$

where

$$(26) \quad G(t) = v(t; g)$$

and

$$(27) \quad (\mathcal{L}h)(t) = v(t; Mh);$$

thus,

$$(\mathcal{T}h)(t) = v(t; g - Mh),$$

and (25) with $w = g - Mh$ implies that

$$Q(D) \mathcal{T}h = g - Mh.$$

Therefore, h_m satisfies (19) if $\mathcal{T}h_m = h_m$.

We assume henceforth that $t_0 > 0$ or, if $N \geq 2$, that $t_0 \geq T_1$, so that $e^{\lambda t} \phi(t)$ is nondecreasing on $[t_0, \infty)$. (See Assumption A.) The proof of Theorem 3 reduces to showing that \mathcal{T} is a contraction mapping of $B(t_0)$ into itself provided that t_0 is sufficiently large, since this implies that there is and h_m in $B(t_0)$ such that $\mathcal{T}h_m = h_m$. We will do this by showing that

$$(28) \quad G \in B(t_0),$$

$$(29) \quad \mathcal{L}(B(t_0)) \subset B(t_0),$$

and that there is a positive function σ on $(0, \infty)$ such that

$$(30) \quad \lim_{t \rightarrow \infty} \sigma(t) = 0$$

and

$$(31) \quad \|\mathcal{L}h\| \leq \sigma(t_0) \|h\|.$$

The following lemma is needed for these proofs.

Lemma 1. *Suppose that u is complex-valued and continuous on $[t_0, \infty)$ and the integral*

$$U(t) = \int_t^\infty u(s) ds$$

converges for $t \geq t_0$. Denote

$$(32) \quad \psi(t) = \sup_{\tau \geq t} \left| \int_{\tau}^{\infty} u(s) ds \right|.$$

Let A be a polynomial, and suppose that γ is a complex constant, with $\operatorname{Re}(\gamma) = \xi$.

(i) If $\xi > 0$, then

$$(33) \quad \left| \int_t^{\infty} A(t-s) e^{\gamma(t-s)} u(s) ds \right| \leq K_1 \psi(t), \quad t \geq t_0,$$

where K_1 is a constant which depends only on γ and A .

(ii) If $\xi < 0$ and there is an α such that $0 < \alpha < -\xi$ and $e^{\alpha t} \psi(t)$ is nondecreasing on $[t_0, \infty)$, then

$$(34) \quad \left| \int_{t_0}^t A(t-s) e^{\gamma(t-s)} u(s) ds \right| \leq K_2 \psi(t), \quad t \geq t_0,$$

where K_2 is a constant which depends only on α, γ , and A .

Proof. (i) Integrating by parts yields

$$(35) \quad \int_t^{\infty} A(t-s) e^{\gamma(t-s)} u(s) ds = A(0) U(t) - \int_t^{\infty} [A(t-s) e^{\gamma(t-s)}]' U(s) ds.$$

From (32),

$$(36) \quad |[A(t-s) e^{\gamma(t-s)}]' U(s)| \leq \psi(t) B(s-t) e^{\xi(t-s)}, \quad s \geq t,$$

where B is a polynomial with nonnegative coefficients determined by γ and the coefficients of A ; therefore, (32) and (35) imply (33), with

$$K_1 = |A(0)| + \int_0^{\infty} e^{-\xi t} B(\tau) d\tau.$$

(ii) Integration by parts yields

$$(37) \quad \int_{t_0}^t A(t-s) e^{\gamma(t-s)} u(s) ds = A(t-t_0) e^{\gamma(t-t_0)} U(t_0) - A(0) U(t) - \int_{t_0}^t [A(t-s) e^{\gamma(t-s)}]' U(s) ds.$$

Our assumptions regarding α imply that

$$\psi(t_0) \leq e^{\alpha(t-t_0)} \psi(t), \quad t \geq t_0;$$

hence,

$$(38) \quad |A(t-t_0) e^{\gamma(t-t_0)} U(t_0)| = |A(t-t_0)| e^{\xi(t-t_0)} \psi(t_0) \leq |A(t-t_0)| e^{(\xi+\alpha)(t-t_0)} \psi(t), \quad t \geq t_0.$$

With B as in (36), the assumption regarding α also implies that

$$(39) \quad \left| \int_{t_0}^t [A(t-t_0) e^{\gamma(t-s)}]^\tau U(s) ds \right| \leq \int_{t_0}^t B(t-s) e^{\xi(t-s)} \psi(s) ds \leq \psi(t) \int_{t_0}^t B(t-s) e^{(\xi+\alpha)(t-s)} ds.$$

Now (32), (37), (38), and (39) imply (34), with

$$K_2 = |A(0)| + \int_0^\infty B(\tau) e^{(\xi+\alpha)\tau} d\tau + \sup_{\tau \geq 0} |B(\tau)| e^{(\xi+\alpha)\tau}.$$

This completes the proof of Lemma 1.

We now turn to the proof of (28). We must show that $G \in C^{(n-1)}[t_0, \infty)$ and that

$$G^{(r)}(t) = O(e^{(\mu_m - \varrho)t} \phi(t)), \quad 0 \leq r \leq n-1.$$

Because of (24) with $w = g$ (see also (26)), this will follow if we show that for $0 \leq r \leq n-1$,

$$(40) \quad \int_{t_0}^t A_{jr}(t-s) e^{\lambda_j(t-s)} g(s) ds = O(e^{(\mu_m - \varrho)t} \phi(t)), \quad 1 \leq j \leq N-1,$$

and

$$(41) \quad \int_t^\infty A_{jr}(t-s) e^{\lambda_j(t-s)} g(s) ds = O(e^{(\mu_m - \varrho)t} \phi(t))$$

if $N \leq j \leq L$. To this end, notice that

$$(42) \quad e^{\lambda_j(t-s)} g(s) = e^{(\lambda_m - \varrho)t} e^{(\lambda_j - \lambda_m + \varrho)(t-s)} [e^{(\varrho - \lambda_m)s} g(s)].$$

Since the integral (14) converges with $j = m$, we can infer (40) for $1 \leq j \leq N-1$ (recall (11) and our condition on α in Assumption A) and (41) for those j 's such that $N \leq j \leq L$ and strict inequality holds in (12), from Lemma 1 with $A = A_{jr}$, $\gamma = \lambda_j - \lambda_m + \varrho$, $u(t) = g(t) e^{(\varrho - \lambda_m)t}$, and $\psi(t) = O(\phi(t))$. If the equality holds in (12), then $A_{jr} = \text{constant}$ (by Assumption A) and (42) reduces to

$$e^{\lambda_j(t-s)} g(s) = e^{(\mu_m - \varrho + i\nu_j)t} [e^{(\varrho - \mu_m - i\nu_j)s} g(s)],$$

so (14) implies (41). This proves (28).

The next lemma will be used to prove (29) and to establish the existence of the function σ satisfying (30) and (31).

Lemma 2. *Suppose that the integrals in (15) and (16) converge, that $h \in B(t_0)$, and that β is a real constant. Then the functions*

$$W_k(t; h) = \int_t^\infty p_k(s) e^{(\varrho - \mu_m + i\beta)s} h^{(n-k)}(s) ds, \quad 1 \leq k \leq n,$$

are defined on $[t_0, \infty)$, and they satisfy the inequalities

$$(43) \quad |W_1(t; h)| \leq \|h\| \int_t^\infty |p_1(s)| \phi(s) ds$$

and

$$(44) \quad |W_k(t; h)| \leq \|h\| \left[\phi(t) \left| \int_t^\infty p_k(\lambda) d\lambda \right| + (1 + |\varrho - \mu_m + i\beta|) \int_t^\infty \left| \int_s^\infty p_k(\lambda) d\lambda \right| \phi(s) ds \right], \quad 2 \leq k \leq n.$$

Proof. The existence of $W_1(t; h)$ and (43) follow from (21) and the assumed existence of the integral on the right side of (43). If $2 \leq k \leq n$, integration by parts yields

$$\begin{aligned} & \int_t^T p_k(s) e^{(\varrho - \mu_m + i\beta)s} H^{(n-k)}(s) ds = \\ & = - \left[e^{(\varrho - \mu_m + i\beta)s} H^{(n-k)}(s) \int_s^\infty p_k(\lambda) d\lambda \right]_t^T + \\ & + \int_t^T [e^{(\varrho - \mu_m + i\beta)s} H^{(n-k)}(s)]' \left(\int_s^\infty p_k(\lambda) d\lambda \right) ds, \end{aligned}$$

and routine estimates based on (21) imply (44), given the assumed convergence of the integrals on the right side. This completes the proof of Lemma 2.

We now turn to the proof of (29). Lemma 2 implies that if $h \in B(t_0)$ and β is a real constant, then

$$\left| \int_t^\infty e^{(\varrho - \mu_m - i\beta)s} M h(s) ds \right| \leq \|h\| \sigma(t; \beta)$$

(see (20)), where

$$(45) \quad \begin{aligned} \sigma(t; \beta) &= \int_t^\infty |p_1(s)| \phi(s) ds + \phi(t) \sum_{k=2}^n \left| \int_t^\infty p_k(s) ds \right| + \\ &+ (1 + |\varrho - \mu_m + i\beta|) \sum_{k=2}^n \int_t^\infty \left| \int_s^\infty p_k(\lambda) d\lambda \right| \phi(s) ds. \end{aligned}$$

Therefore, since

$$e^{\lambda_j(t-s)} M h(s) = e^{(\lambda_m - \varrho)t} e^{(\lambda_j - \lambda_m + \varrho)(t-s)} [e^{(\varrho - \lambda_m)s} M h(s)].$$

Assumption A and Lemma 1 with $A = A_{jr}$, $\gamma = \lambda_j - \lambda_m + \varrho$, $u(t) = e^{(\varrho - \lambda_m)t} M h(t)$, and $\psi(t) = \|h\| \sup_{\tau \geq t} \sigma(\tau; 0)$ imply that there is a constant K (independent of h and t_0) such that

$$(46) \quad \begin{aligned} \left| \int_{t_0}^t A_{jr}(t-s) e^{\lambda_j(t-s)} M h(s) ds \right| &\leq K \|h\| e^{(\mu_m - \varrho)t} \sup_{\tau \geq t} \sigma(\tau; 0), \\ 0 \leq r \leq n-1, \quad 1 \leq j \leq N-1, \end{aligned}$$

and

$$(47) \quad \left| \int_t^\infty A_{jr}(t-s) e^{\lambda_j(t-s)} M h(s) ds \right| \leq K \|h\| e^{(\mu_m - \varrho)t} \sup_{\tau \geq t} \sigma(\tau; 0), \quad 0 \leq r \leq n-1,$$

If $N \leq j \leq L$ and (10) does not hold. On the other hand, if (10) holds, then

$$e^{\lambda_j(t-s)} M h(s) = e^{(\mu_m - \varrho + i\nu_j)t} [e^{(\varrho - \mu_m - i\nu_j)s} M h(s)]$$

and $A_{jr} = \text{constant}$ (Assumption A), so we can choose K so that (47) also holds if $\sigma(\tau; 0)$ is replaced by $\sigma(\tau; \nu_j)$. Since (15), (16), and (45) imply that $\sigma(t; \beta) = o(\phi(t))$ for all real β , (24) with $w = Mh$, (27), (46), and (47) imply that

$$(48) \quad |(\mathcal{L}h)^{(r)}(t)| \leq \|h\| e^{(\mu_m - \varrho)t} \phi(t) / n, \quad 0 \leq r \leq n-1, \quad t \geq t_0,$$

where σ satisfies (30). Since (21) and (48) imply (31), we now conclude that \mathcal{T} has a fixed point h_m in $B(t_0)$ if t_0 is sufficiently large; hence x_m as defined in (22) satisfies (1) and (17). To deduce the improved estimate (18) in the case where (14) is replaced by

$$(49) \quad \int_t^\infty g(s) e^{(\varrho - \mu_m - i\nu_j)s} ds = o(\Phi(t)),$$

it suffices to show that

$$(50) \quad h_m^{(r)}(t) = o(e^{(\mu_m - \varrho)t} \phi(t)), \quad 0 \leq r \leq n-1,$$

in this case. Since $h_m = G - \mathcal{L}h_m$, we see from (30) and (48) (with $h = h_m$) that (50) will follow if

$$(51) \quad G^{(r)}(t) = o(e^{(\mu_m - \varrho)t} \phi(t)), \quad 0 \leq r \leq n-1.$$

To see that this is so, define

$$\phi_1(t) = \sup_{\tau \geq t} \left\{ \max_{j \in S} \left| \int_\tau^\infty g(s) e^{(\varrho - \mu_m - i\nu_j)s} ds \right| \right\}$$

(see (7)). Applying the argument used earlier to prove (28), now with ϕ replaced by ϕ_1 , shows that

$$G^{(r)}(t) = O(e^{(\mu_m - \varrho)t} \phi_1(t)), \quad 0 \leq r \leq n-1.$$

Since (49) implies that $\phi_1(t) = o(\phi(t))$, this implies (51) and completes the proof of Theorem 1.

3. RELATIONSHIP OF THE MAIN THEOREM WITH ŠIMŠA'S RESULT

We first deduce the following corollary from Theorem 3.

Corollary 1. *Suppose that Assumption A and (15) hold, and that*

$$(52) \quad \int_t^\infty f(s) e^{(\varrho - \mu_m - i\nu_j)s} ds = O(\phi(t))$$

and

$$(53) \quad \int_t^\infty p_k(s) e^{[\varrho + i(v_m - v_j)]s} ds = O(\phi(t)), \quad 1 \leq k \leq n$$

for j in S (see (7)). Then (1) has a solution x_m which satisfies (17), provided that (16), holds. Moreover, (16) holds automatically if any one of the following is true:

(i) $\varrho > 0$; (ii) "O" can be replaced by "o" in (53) and

$$(54) \quad \int_t^\infty \phi^2(s) ds = O(\phi(t));$$

or (iii)

$$(55) \quad \int_t^\infty \phi^2(s) ds = o(\phi(t)).$$

Finally, if (52) and (53) hold with "O" replaced by "o", then x_m satisfies (18).

Proof. From (13), (52) and (53) imply (14). Therefore, Theorem 3 implies the conclusion if (16) holds. To complete the proof, we need only show that (16) follows from each of (i), (ii), and (iii). Integrating (53) (with $j = m$) by parts shows that

$$(56) \quad \int_t^\infty p_k(s) ds = O(e^{-\varrho t} \phi(t)), \quad 2 \leq k \leq n,$$

and therefore

$$(57) \quad \int_t^\infty \left| \int_s^\infty p_k(\lambda) d\lambda \right| \phi(s) ds = O\left(\int_t^\infty e^{-\varrho s} \phi^2(s) ds \right), \quad 2 \leq k \leq n.$$

The right side of (57) is $o(\phi(t))$ if either $\varrho > 0$ or (55) holds; hence, (i) and (iii) imply (16). To see that (ii) also implies (16), we have only to observe that if (53) holds with "o" on the right, then so does (56) and therefore (57). Given this, (54) implies (16) even if $\varrho = 0$. This proves Corollary 1.

We conclude by showing that Corollary 1 implies Theorem 2. It suffices to show that the integrability conditions of the latter imply those of the former, with $\phi(t) = t^{-q}$. Obviously, (5) implies (15) for any nonincreasing ϕ . Integrating by parts shows that if (6) converges, then

$$\int_t^\infty p_k(s) e^{[\varrho + i(v_m - v_j)]s} ds = o(t^{-q}), \quad j \in S,$$

which verifies (53) with "O" replaced by "o". Since

$$\int_t^\infty s^{-2q} ds = \frac{t^{-2q+1}}{(2q-1)} \quad (q > 1/2),$$

(54) holds if $q \geq 1$. Finally, an argument using integration by parts shows that if (8)

holds for some $q > 0$, then

$$(58) \quad \int^{\infty} \left| \int_s^{\infty} p_k(\lambda) d\lambda \right| ds < \infty .$$

(The converse is false.) Obviously, (58) implies (15) for any nonincreasing ϕ .

Since the integrability conditions of Theorem 1 imply those of Theorem 2 with $\varrho = 0$, Corollary 1 also implies Theorem 1.

References

- [1] *O. Dunkel*: Regular singular points of a system of homogeneous linear differential equations of the first order, Proc. Amer. Acad. Arts Sci. 38 (1902–3), 341–370.
- [2] *P. Hartman*: Ordinary Differential Equations, John Wiley, New York, 1964.
- [3] *J. Šimša*: Asymptotic integration of perturbed linear differential equations under conditions involving ordinary integral convergence, SIAM J. Math. Anal. 15 (1984), 116–123.
- [4] *J. Šimša*: The second order differential equation with oscillatory coefficient, Arch. Math. (Brno) 18 (1982), 95–100.
- [5] *J. Šimša*: The condition of ordinary integral convergence in the asymptotic theory of linear differential equations with almost constant coefficients, SIAM J. Math. Anal., 16 (1985), 757–769.

Author's address: Department of Mathematics and Computer Science, Drexel University, Philadelphia, Pennsylvania 19104, U.S.A.