

Jaroslav Kurzweil; Garyfalos Pappaschinopoulos
Structural stability of linear discrete systems via the exponential dichotomy

Czechoslovak Mathematical Journal, Vol. 38 (1988), No. 2, 280–284

Persistent URL: <http://dml.cz/dmlcz/102223>

Terms of use:

© Institute of Mathematics AS CR, 1988

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

STRUCTURAL STABILITY OF LINEAR DISCRETE SYSTEMS VIA
THE EXPONENTIAL DICHOTOMY

JAROSLAV KURZWEIL, Praha, and GARYFALOS PAPASCHINOPOULOS, Xanthi

(Received April 1, 1986)

INTRODUCTION

Consider the difference equation

$$(1) \quad x(n+1) = A(n)x(n), \quad n \in N = \{0, 1, \dots\}$$

where $A(n)$ is an invertible $k \times k$ -matrix for $n \in N$ such that

$$|A(n)| \leq M, \quad |A^{-1}(n)| \leq M, \quad M > 0, \quad n \in N.$$

We denote by W the space of the systems of the form (1) and by $|\cdot|$ the Euclidean norm.

Equation (1) is said to possess an exponential dichotomy if there exist a projection P ($P^2 = P$) and constants $K > 0$, $0 < p < 1$, such that

$$|X(n)PX^{-1}(m)| \leq Kp^{n-m}, \quad n \geq m$$

$$|X(n)(I-P)X^{-1}(m)| \leq Kp^{m-n}, \quad m \geq n$$

where $X(n)$ is the fundamental matrix solution of (1) such that $X(0) = I$.

Consider a system in W

$$(2) \quad y(n+1) = B(n)y(n).$$

According to [5, p. 17], (1) and (2) are said to be topologically equivalent if there exists a function $h: N \times R^k \rightarrow R^k$ with the following properties:

- (i) if $|x| \rightarrow \infty$, then $|h(n, x)| \rightarrow \infty$ uniformly with respect to n ,
- (ii) the map $h_n(\cdot) = h(n, \cdot)$ from R^k to R^k is a homeomorphism for each n ,
- (iii) the map $g_n(\cdot) = h_n^{-1}(\cdot)$ from R^k to R^k also has property (i),
- (iv) if $x(n)$ is a solution of (1) then $h(n, x(n))$ is a solution of (2).

Equation (1) is called structurally stable if there exists $\delta > 0$ such that if (2) belongs to W and $|B(n) - A(n)| < \delta$ then (2) is topologically equivalent to (1).

The results of this paper are:

- (i) If equation (1) has an exponential dichotomy then it is topologically equivalent

From [1, p. 39] we have

$$\log B_1(n) = \frac{1}{2\pi i} \int_{\gamma} (zI - B_1(n))^{-1} \log z \, dz$$

where γ is any simple closed curve which contains in its interior every characteristic root of $B_1(n)$ but not the origin.

We claim that $\log B_1(n)$ is a real valued bounded matrix for $n \in N$. It is obvious that it is a real matrix since $B_1(n)$ is an upper triangular matrix with real positive diagonal elements. Since the matrices $B_1(n), B_1^{-1}(n)$ are bounded for $n \in N$ there exist constants $\lambda, \mu > 0$ such that $\lambda \leq \lambda_i(n) \leq \mu, n \in N, i = 1, \dots, l$ and $\lambda_i(n)$ are the eigenvalues of $B_1(n)$. We choose $r > \max\{\frac{1}{2}(\varepsilon + \mu - \lambda), \frac{1}{2}(\varepsilon + \mu)\}, \lambda > \varepsilon > 0$. So $-r + \varepsilon + \mu < r < r - \varepsilon + \lambda$. Consider $z_0 \in R^+ : r < z_0 < r - \varepsilon + \lambda$. It is obvious that $|z_0 - \lambda| < r - \varepsilon$ and $|z_0 - \mu| < r - \varepsilon$. Let γ be the sphere $|z - z_0| = r$. Then we have $|z - \lambda_i(n)| = |z - z_0 + z_0 - \lambda_i(n)| \geq |z - z_0| - |z_0 - \lambda_i(n)| \geq r - (r - \varepsilon) = \varepsilon, i = 1, \dots, l$. So if $|B_1(n)| \leq L, L > 0, n \in N$ then [2, p. 47] implies

$$|(zI - B_1(n))^{-1}| \leq \frac{c|zI - B_1(n)|^{l-1}}{|z - \lambda_1(n)| \dots |z - \lambda_l(n)|} \leq \frac{c(z_0 + r + L)^{l-1}}{\varepsilon^l}$$

c a constant.

Therefore $\log B_1(n)$ is a bounded matrix for $n \in N$. Hence our claim is proved.

Let $Z_1(t)$ be the fundamental matrix of (8) such that $Z_1(0) = I$. Then

$$\begin{aligned} Z_1(t) &= e^{\int_n^{t \log B_1(n) ds}} Z_1(n) = e^{(t-n) \log B_1(n)} Z_1(n), \quad n \leq t < n+1, \\ Z_1(t) &= e^{(t-n-1) \log B_1(n+1)} Z_1(n+1), \quad n+1 \leq t < n+2. \end{aligned}$$

The above relations yield

$$Z_1(t) = e^{(t-n-1) \log B_1(n+1)} e^{\log B_1(n)} Z_1(n), \quad n+1 \leq t < n+2.$$

Take $t = n+1$. Then we obtain

$$Z_1(n+1) = B_1(n) Z_1(n), \quad n \in N.$$

Therefore $Z_1(n)$ is a fundamental matrix solution of (7).

We prove that equation (8) has an exponential dichotomy. If $z(t)$ is a solution of (8) then for $t \in R^+$ we have

$$z(t) = Z_1(t) Z_1^{-1}([t]) z([t]), \quad [t] \text{ being the integral part of } t.$$

If $|B(t)| \leq \bar{L}$ then for $t \in R^+$ we have $|Z_1(t) Z_1^{-1}([t])| \leq e^{L(t-[t])} \leq e^{\bar{L}}$. So for $t \geq s \geq 0$ and provided $e^{-a}, a > 0, \bar{K} > 0$ are the constants of the exponential dichotomy of (7) we have

$$\begin{aligned} |z(t)| &\leq e^{\bar{L}} |z([t])| \leq \bar{K} e^{\bar{L}} e^{-a([t]-[s])} |z([s])| \leq \\ &\leq \bar{K} e^{\bar{L}} e^{-a(t-1-s-1)} |z([s])| \leq \bar{K} e^{2\bar{L}} e^{2a} e^{-a(t-s)} |z(s)|, \quad t \geq s \geq 0. \end{aligned}$$

Then by Palmer's Theorem in [4, p. 9] and arguing as in [5, p. 20] we prove that (8)

and $x'_1 = -x_1$ are topologically equivalent (c.f. [5, p. 17]). Let $h_1: R^+ \times R^1 \rightarrow R^1$ be the corresponding homeomorphism. If $z(t, 0, x)$ is a solution of (8) such that $z(0) = x$ we get

$$h_1(t, z(t, 0, x)) = e^{-t}h_1(0, x), \quad t \in R^+, \quad x \in R^1.$$

Take $t = n, n \in N$. So (7) and $x_1(n+1) = e^{-1}x_1(n)$ are topologically equivalent.

Now consider system (6). Proceeding as in (5) we obtain a unitary matrix $U_2(n)$ such that the change of variables $y = U_2(n)z$ transforms (6) into a system

$$(9) \quad z(n+1) = B_2(n)z(n)$$

where $B_2(n)$ is a $(k-l) \times (k-l)$ upper triangular matrix in which the diagonal elements are real and positive functions on N . Arguing as in (7) we prove that (9) and $x_2(n+1) = ex_2(n)$ are topologically equivalent. Let $h_2: N \times R^{k-1} \rightarrow R^{k-1}$ be the corresponding homeomorphism. Therefore the system

$$z(n+1) = \text{diag}(B_1(n), B_2(n))z(n)$$

is topologically equivalent to (3). Hence (4) is topologically equivalent to (3). The corresponding homeomorphism is $\bar{h}(n, x) = (h_1(n, \{U^{-1}(n)x\}_1), h_2(n, \{U^{-1}(n)x\}_2))$ where $\{U^{-1}(n)x\}_1$ and $\{U^{-1}(n)x\}_2$ are the components of $U^{-1}(n)x$ in R^1 and R^{k-1} , respectively, $U(n) = \text{diag}(U_1(n), U_2(n))$. So (1) and (3) are topologically equivalent. The corresponding homeomorphism is $h(n, x) = \bar{h}(n, S^{-1}(n)x)$ and the proof of the proposition is complete.

Proposition 2. Equation (1) is structurally stable if and only if it has an exponential dichotomy.

Proof. The proof of necessity is given in [10, Proposition 2]. Using Proposition 1, the roughness of the exponential dichotomy [3, p. 232] and the same argument as in [5, p. 20] we can easily prove sufficiency.

References

- [1] *W. A. Coppel: Stability and Asymptotic Behaviour of Differential Equations*, Heath, Boston, 1965.
- [2] *W. A. Coppel: Dichotomies in Stability Theory*, Lecture Notes in Mathematics, No. 629, Springer Verlag, Berlin, 1978.
- [3] *D. Henry: Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Mathematics, No. 840, Springer-Verlag, Berlin, 1981.
- [4] *K. J. Palmer: A characterization of exponential dichotomy in terms of topological equivalence*. J. Math. Anal. Appl. 69 (1979), 8–16.
- [5] *K. J. Palmer: The structurally stable linear systems on the half-line are those with exponential dichotomies*. J. Differential Equations, 33 (1979), 16–25.
- [6] *G. Papaschinopoulos and J. Schinas: Criteria for an exponential dichotomy of difference equations*, Czechoslovak Math. J. 35 (110) 1985, 295–299.
- [7] *G. Papaschinopoulos and J. Schinas: A criterion for the exponential dichotomy of difference equations*, Rend. Sem. Fac. Sci. Univ. Cagliari, Vol. 54, fasc. 1 (1984), 61–71.
- [8] *G. Papaschinopoulos and J. Schinas: Multiplicative separation, diagonalizability and struc-*

tural stability of linear difference equations, *Differential Equations: Qualitative theory* (Szeged 1984), *Colloq. Math. Soc. Janos Bolyai*, 47, North-Holland, Amsterdam—New York.

- [9] *G. Papaschinopoulos*: Exponential separation, exponential dichotomy and almost periodicity of linear difference equations, *J. Math. Anal. Appl.* 120 (1986), 276—287.
- [10] *G. Papaschinopoulos* and *J. Schinas*: Structural stability via the density of a class of linear discrete systems, *J. Math. Anal. Appl.* (to appear).
- [11] *J. Schinas* and *G. Papaschinopoulos*: Topological equivalence for linear discrete systems via dichotomies and Lyapunov functions, *Boll. Un. Math. Ital.* 6, 4 (1985), 61—70.

Authors' addresses: J. Kurzweil, 115 67 Praha 1, Žitná 25, Czechoslovakia (Matematický ústav ČSAV), G. Papaschinopoulos, Democritus University of Thrace, 671 00 Xanthi, Greece.