

Jiří Rachůnek

Lattice-ordered groups with minimal prime subgroups satisfying a certain condition

Archivum Mathematicum, Vol. 9 (1973), No. 3, 147--149

Persistent URL: <http://dml.cz/dmlcz/104805>

Terms of use:

© Masaryk University, 1973

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

LATTICE-ORDERED GROUPS WITH MINIMAL PRIME SUBGROUPS SATISFYING A CERTAIN CONDITION

JIŘÍ RACHŮNEK, OLOMOUČ

(Received May 10, 1973)

In this paper one problem of P. Conrad's book [2] is partially solved in connection with one problem of F. Šik. There is proved (Theorem 1) that the set of all cardinal summands of an l -group G is equal to the set of all polars of this group if and only if G is projectable and satisfies a certain property. Further a connection between minimal prime subgroups and cardinal summands and also a connection between minimal prime subgroups and polars is shown here.

Let $G = [G, +, \vee]$ be an l -group. For $x \in G$ we shall denote $|x| = x \vee -x$. If $|a| \wedge |b| = 0$, then elements $a, b \in G$ will be called *disjoint*. If $\emptyset \neq A \subseteq G$, then we denote $A' = \{x \in G: |x| \wedge |a| = 0 \text{ for each } a \in A\}$. Now $A \subseteq G$ is called *polar* if $A'' = A$. (A'' denotes $(A')'$.) Instead of $\{a'\}, \{a''\}$ we write a', a'' , respectively. It is known that any polar is a convex l -subgroup of G . The set of all polars of G will be denoted by $\Gamma = \Gamma(G)$. If $B \in \Gamma$, then B, B' are called *complementary polars*.

The following theorem has been proved by F. Šik in [3] (Teorema 1):

Theorem A. (1) *Polars of an l -group form a complete Boolean algebra Γ (ordered by inclusion, an infimum is formed by an intersection).*

(2) *Polars that are l -ideals form a closed subalgebra Γ_1 of Γ .*

(3) *Cardinal summands of G form a subalgebra Γ_2 of Γ_1 (not always complete), where a supremum is formed by a sum of summands.*

It holds that for $B \in \Gamma_2(G)$ it is $G = B \oplus B'$. An l -group G is called an r -group if it is isomorphic to a subdirect product of totally ordered groups. By [4], an l -group is an r -group if and only if each its polar is an l -ideal. A convex l -subgroup P is called *prime* if the following is satisfied:

- (i) *If $x \notin P$, then $x' \subseteq P$.*
- (i) and the following conditions are equivalent:
 - (ii) *P contains at least one of polars a'', a' ($a \in G$).*
 - (iii) *P contains at least one of complementary polars.*

Any prime subgroup contains at least one minimal prime subgroup. In $G \neq \{0\}$, minimal prime subgroups are characterized among convex l -subgroups as: $a \notin P$ iff $a' \subseteq P$. A convex l -subgroup Z is a z -subgroup if from $x \in Z$ and $y' = x'$ it follows $y \in Z$. It is known that every polar and every minimal prime subgroup is a z -subgroup. An l -group is called *projectable* if $G = g' \oplus g''$ for each $g \in G$. Clearly any projectable l -group is an r -group.

The following theorem is proved in [1] (Théorème 3.1):

Theorem B. *An l -group G is projectable if and only if any proper prime subgroup contains exactly one prime z -subgroup.*

The problem how to characterize those l -groups for which $\Gamma_2(G) = \Gamma(G)$ has been given by P. Conrad in the book [2, p. 2.8]. Clearly any such l -group will be projectable.

Note. This problem has been solved by F. Šik in [3, p. 8] yet. He has proved that for an l -group the following are equivalent:

- (1) An arbitrary polar is a direct summand.
- (2) A sum of two arbitrary polars is also a polar.
- (3) A sum of an arbitrary pair of complementary polars is also a polar.
- (4) Any pair of complementary polars forms a direct decomposition of this l -group.

Another characterization is given in [4, Satz 13].

Further denote the following condition:

(*) For each minimal prime subgroup A of an l -group G and for each polar K of G it is satisfied: $K \subseteq A$ iff $K' \not\subseteq A$.

F. Šik has proposed (in a letter) the problem how to characterize l -groups with the property (*).

The following theorem shows a certain connection between both problems.

Theorem 1. For an l -group $G \neq \{0\}$ it holds $\Gamma(G) = \Gamma_2(G)$ if and only if G is projectable and possesses the property (*).

Proof. a) Let $\Gamma(G) = \Gamma_2(G)$ and let A be a minimal prime subgroup of G . Let $K \in \Gamma(G)$, $K, K' \subseteq A$. Since $K \oplus K' = G$, $A = A + A \supseteq K + K' = G$. If $G \neq \{0\}$, then by [5, Folgerung 7.3] $A \neq G$, a contradiction. But since A is a prime subgroup, it contains K or K' . Thus G satisfies (*).

b) Let G be projectable and have the property (*). Let $K \in \Gamma(G)$ such that $K \oplus K' \neq G$. Let P be a proper prime subgroup of G such that $K \oplus K' \subseteq P$. Let us remind yet that the filet of an element $x \in G$ is $x = \{y \in G: y' = x'\}$ and the set of all filets $\mathcal{F}(G)$ form a distributive lattice. Denote thus $\Phi = \{x: x \notin P\}$. Evidently Φ is a filter of $\mathcal{F}(G)$. For each $y \in K \cup K'$ it holds $\bar{y} \notin \Phi$. (If, namely, $y \in K$, $\bar{y} \in \Phi$, then $y'' = a''$ for some $a \notin P$ thus $y'' \not\subseteq P$; but $y'' \subseteq K$, and we have a contradiction. Similarly for $z \in K'$.)

Now if $x \in K \cup K'$, then denote a maximal filter of $\mathcal{F}(G)$ that contains Φ and does not x by Φ^x . It holds Φ^x is a prime filter. Therefore $Z^x = \{u \in G: u \notin \Phi^x\}$ is a prime z -subgroup of G and clearly $Z^x \subseteq P$. Since G is projectable, all prime z -subgroups contained in $P \neq G$ are (by Theorem B) identical, thus for each $x_1, x_2 \in K \cup K'$ $Z^{x_1} = Z^{x_2}$. Further $\Phi^{x_1} = \Phi^{x_2}$ iff $\{u: u \notin \Phi^{x_1}\} = \{v: v \notin \Phi^{x_2}\}$ and this holds iff $Z^{x_1} = Z^{x_2}$. Thus for each $x_1, x_2 \in K \cup K'$ $\Phi^{x_1} = \Phi^{x_2}$ and therefore $\Psi = \bigcap_{x \in K \cup K'} \Phi^x = \Phi^x$ for each $x \in K \cup K'$. Hence Ψ is a prime filter of $\mathcal{F}(G)$ and $Z = \{w: w \notin \Psi\}$ is a prime z -subgroup of G such that $Z \subseteq P$. Consequently, by [1, Proposition 3.1 and its proof], $Z = \bigcup_{a \notin P} a'$.

For each $x \in K \cup K'$ $x \in Z$, therefore $K \subseteq Z$, $K' \not\subseteq Z$ and this contradicts the assumption that G satisfies (*).

Now, it is easy to prove the further

Theorem 2. For a projectable l -group G the following conditions are equivalent:

- (1) Any polar of G is a cardinal summand of G . (Thus $\Gamma(G) = \Gamma_2(G)$.)
- (2) G satisfies the property (*).
- (3) The algebra $\Gamma_2(G)$ is a \vee -closed subalgebra of $\Gamma(G)$.
- (4) The algebra $\Gamma_2(G)$ is a \wedge -closed subalgebra of $\Gamma(G)$.

Proof. (3) \Rightarrow (1): Let $K \in \Gamma(G)$. It holds $K = \bigvee_{a \in K} \Gamma a''$ and $a'' \in \Gamma_2(G)$ implies by (3), $K \in \Gamma_2(G)$.

(4) \Rightarrow (1): If $K \in \Gamma(G)$, then $K = \bigwedge_{b \in K'} \Gamma b'$. We have $b' \in \Gamma_2(G)$, thus by (4), $K \in \Gamma_2(G)$.

If H is a prime subgroup of an l -group G , then we say H has the property (**) if it holds:

(**) If $K \in \Gamma(G)$ then $K \subseteq H$ iff $K' \not\subseteq H$.

Further we say $\emptyset \neq A \subseteq G$ is dense in G if $A' = \{0\}$.

We get

Theorem 3. A prime subgroup H of an l -group G is either a polar in G or it is dense in G .

Proof. Let H not be dense. Then $\{0\} \neq H' \not\subseteq H$. Therefore $H'' \subseteq H$ i.e. H is a polar.

The following theorem is a consequence of Theorems 3 and 1.

Theorem 4. If a projectable l -group G satisfies (*) then each minimal prime subgroup of G is a cardinal summand or it is dense in G .

Denote now the set of all z -subgroups of an l -group G by $\mathcal{Z}(G)$. It is known (see [1, Proposition 2.3]) $\mathcal{Z}(G)$ forms a complete distributive lattice. It holds $\Gamma(G) \subseteq \mathcal{Z}(G)$ but generally $\Gamma(G)$ need not be a sublattice of $\mathcal{Z}(G)$.

We get

Theorem 5. Let G be an l -group and $\Gamma(G)$ a closed sublattice of $\mathcal{Z}(G)$. Then a proper prime subgroup H of G has the property (**) if and only if H is a polar.

Proof. If $H \in \mathcal{Z}(G)$, then (by [1, Proposition 2.1]) $H = \bigcup_{a \in H} a'' = \bigvee_{a \in H} a''$. By the assumption $\bigvee_{a \in H} \Gamma a'' = \bigvee_{a \in H} a''$ thus H is a polar. The converse is evident.

Therefore it holds also

Theorem 6. a) Let an l -group G satisfy (*) and let $\Gamma(G)$ be a closed sublattice of $\mathcal{Z}(G)$. Then each minimal prime subgroup of G is a polar in G .

b) Let, in addition, G be projectable. Then each minimal prime subgroup is a cardinal summand of G .

REFERENCES

- [1] BIGARD, A.: *Contribution à la théorie des groupes réticulés*, Thèse Fac. Sc. Paris, 1969
- [2] CONRAD, P.: *Lattice ordered groups*, Tulane Univ., 1970
- [3] ŠIK, F.: *K teorii strukturno uporjadočennykh grupp*, Czech. Math. J., 6 (1956), 1–25
- [4] ŠIK, F.: *Über subdirekte Summen geordneter Gruppen*, Czech. Math. J., 10 (1960), 400–424
- [5] ŠIK, F.: *Struktur und Realisierungen von Verbandsgruppen III*, Mem. Fac. Cie. Univ. Habana, 1966, 1–20

Jiří Rachánek
 Department of Mathematics
 Palacký University
 Leninova 26, 771 46 Olomouc
 Czechoslovakia