

Ivo Babuška

A remark to the finite element method

Commentationes Mathematicae Universitatis Carolinae, Vol. 12 (1971), No. 2, 367--376

Persistent URL: <http://dml.cz/dmlcz/105352>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1971

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A REMARK TO THE FINITE ELEMENT METHOD

Ivo BABUŠKA, College Park

I. Introduction. The finite element method, its different versions has become very important today in the theory and practice. See e.g. [1] - [17] and others.

One important form of the method is such that the approximate solution $u_{h_n}(X)$ on the domain Ω has the following form

$$(1.1) \quad u_{h_n}(X) = \sum c(h, \underline{k}) \varphi\left(\frac{X}{h} - \underline{k}\right)$$

where the function $\varphi(X)$ has a compact support and the sum is over all multiintegers $\underline{k} = (k_1, \dots, k_n)$, (k_i integers) such that

$$\text{supp } \varphi\left(\frac{X}{h} - \underline{k}\right) \cap \Omega \neq \emptyset.$$

Many times the function $\varphi(x)$, $x \in \mathbb{R}_m$ is taken in a special form

$$(1.2) \quad \varphi(X) = \prod_{i=1}^m \varphi(x_i), \quad i = 1, 2, \dots, m, \quad X \equiv (x_1, \dots, x_m).$$

For some domains e.g. a square, for $\varphi(X)$ smooth enough so called "inverse theorem" is valid for some sequences

$$h_i, \quad i = 1, 2, \dots, h_i \rightarrow 0.$$

$$(1.3) \quad \|u_{h_i}(x)\|_{W_2^l(\Omega)} \leq C(l) h_i^{-l} \|u_{h_i}\|_{W_2^0(\Omega)}.$$

where $W_2^l(\Omega)$ is the usual Sobolev space and $C(l)$ does not depend on i .

Many very important theorems of the theory of the finite elements method are based on the validity of the mentioned "inverse theorem". Under this assumption e.g. the computation of derivatives (see [14]) could be made much easier as in [14], simply by derivating the approximate solution. Without this "inverse theorem" proofs of many theorems will break down.

The purpose of this note is to show that for smooth domains the inverse theorem is not true for any choice of h_i .

To simplify the arguments we shall deal only with the two dimensional case i.e. $\Omega \subset R_2$.

The inverse theorem is in fact equivalent with a geometric property of the domain Ω . Let us explain this property.

Let $\Omega \subset R_2$ be a bounded domain.

For every $h > 0$ let us define the mesh $P_h = \{(kh, kl)\}$ with k, l integers.

Further let Q_h be the set of the all squares $S_{k,l}^h$

$$S_{k,l}^h \equiv \{x = (x_1, x_2); kh < x_1 < h(k+1), \\ hl < x_2 < h(l+1)\}$$

and

$$Q_h^\Omega \equiv \{ \cup_{k,l} S_{k,l}^h; S_{k,l}^h \cap \Omega \neq \emptyset \},$$

$$P_h^\Omega = \{(k, l); S_{k,l}^h \in Q_h^\Omega\}.$$

Let us define now the function $\psi_h(k, l)$ on P_h by the following way:

$$(1.4) \quad \psi_{h_n}(k, l) = \frac{\text{mes}(S_{k,l}^{h_n} \cap \Omega)}{h_n^2}$$

where $(S_{k,l}^{h_n} \cap \Omega)$ means the measure of the set $S_{k,l}^{h_n} \cap \Omega$. Further let

$$(1.5) \quad \chi(h_n) = \min_{k,l} \psi_{h_n}(k, l), \quad (k, l) \in P_{h_n}^{\Omega}.$$

The crucial question is whether we may choose a sequence $h_{n_j} \rightarrow 0, j = 1, 2, \dots$ such that

$$(1.6) \quad \liminf_{j \rightarrow \infty} \chi(h_{n_j}) \geq \alpha.$$

Certainly if Ω is a square then such a sequence h_{n_j} clearly exists.

In the next chapter we shall show that such a sequence does not exist in the case that the boundary has bounded curvature.

For the simplicity we shall prove the statement for a circular domain only. By the same idea it is obviously possible to prove the general statement for domains with bounded curvature in \mathbb{R}_m .

II. Theorem. As we said we shall study the case of the circular domain Ω in \mathbb{R}_2 only. Let

$$(2.1) \quad K(\kappa) \equiv \{(x, y); x^2 + y^2 < \kappa^2\}$$

be a circle with the radius κ .

Further let us denote $\varphi(x, y)$ the distance of the point (x, y) to the boundary of the circle $K(\kappa)$.

Let us prove the following theorem.

Theorem. Let us denote

$$(2.2) \quad \varrho(\kappa) = \min_{(k,l) \in K(\kappa)} \varphi(k, l) ,$$

k, l integers.

Then for every sequence $\kappa_i \rightarrow \infty$ we have

$$(2.3) \quad \lim_{i \rightarrow \infty} \inf \varrho(\kappa_i) = 0 .$$

This theorem is obviously equivalent with the statement which we introduced in the first chapter in the case that Ω is a circle.

Proof. 1) Let us denote by $U(\kappa)$ the number of all (k, l) in $K(\kappa)$,

$$(2.4) \quad U(\kappa) = \sum_{(k,l) \in K(\kappa)} 1 .$$

Obviously $U(\kappa)$ is a not decreasing step function. Let $\xi_i, \xi_{i+1} > \xi_i$ be the sequence of all points of the discontinuity of the function $U(\kappa)$.

Denote further

$$(2.5) \quad \lambda(\kappa) = \sup_{\xi_i \geq \kappa} |\xi_i - \xi_{i-1}| .$$

Our theorem will be proved if we shall show that

$$(2.6) \quad \lim_{\kappa \rightarrow \infty} \lambda(\kappa) = 0 .$$

In fact

$$(2.7) \quad \varrho(\kappa_i) \leq \lambda(\kappa_i) .$$

2) Let us assume on the contrary that there exists a sequence $\kappa_i, i = 1, 2, \dots, \kappa_{i+1} > \kappa_i, \kappa_i \rightarrow \infty$ such that

$$(2.8) \quad \liminf_{i \rightarrow \infty} \lambda(\kappa_i) \geq 3\Delta > 0.$$

Then for all κ_i , $i \geq N$ we have

$$(2.9) \quad \lambda(\kappa_i) \geq 2\Delta.$$

So we may construct the subsequence ξ_{i_j} , such that

$$(2.10) \quad \xi_{i_j} - \xi_{i_{j-1}} \geq \Delta$$

and

$$\xi_{i_j} \geq \kappa_j.$$

Let us denote

$$(2.11) \quad \omega_j = [\xi_{i_{j-1}}]$$

where $[x]$ means integral part of x .

We have

$$(2.12) \quad \omega_j \leq \xi_{i_{j-1}} \leq \omega_j + 1.$$

Because ω_j and $\omega_j + 1$ is certainly the point of the discontinuity of the function $U(\kappa)$ we have

$$(2.12) \quad \omega_j \leq \xi_{i_{j-1}} \leq \xi_{i_{j-1}} + \Delta \leq \xi_{i_j} \leq \omega_j + 1$$

and

$$(2.13) \quad \xi_{i_{j-1}} = \omega_j + \sigma_j \quad \text{where } 0 \leq \sigma_j \leq 1 - \Delta.$$

Let us define now the numbers α_j resp. β_j such that

$$(2.14) \quad \omega_j^2 + \alpha_j^2 = \xi_{i_{j-1}}^2 = (\omega_j + \sigma_j)^2,$$

$$(2.15) \quad \omega_j^2 + \beta_j^2 = \xi_{i_j}^2 \geq (\omega_j + \sigma_j + \Delta)^2.$$

Therefore

$$(2.16) \quad \alpha_j = \sqrt{(\omega_j + \sigma_j)^2 - \omega_j^2} = \omega_j^{\frac{1}{2}} \sigma_j^{\frac{1}{2}} (2 + \sigma_j \omega_j^{-1})^{\frac{1}{2}},$$

$$(2.17) \quad \beta_j \geq \omega_j^{\frac{1}{2}} (\sigma_j + \Delta)^{\frac{1}{2}} (2 + (\sigma_j + \Delta) \omega_j^{-1})^{\frac{1}{2}}.$$

But

$$\omega_j \geq \kappa_j - 1; \quad 0 \leq \sigma_j < 1.$$

Therefore

$$(2.18) \quad \alpha_j \leq \omega_j^{\frac{1}{2}} \sigma_j^{\frac{1}{2}} 2^{\frac{1}{2}} \left(1 + \frac{1}{(\kappa_j - 1)2}\right)^{\frac{1}{2}}$$

and hence

$$(2.19) \quad \beta_j - \alpha_j \geq 2^{\frac{1}{2}} \omega_j^{\frac{1}{2}} \left[(\sigma_j + \Delta)^{\frac{1}{2}} - \sigma_j^{\frac{1}{2}} \left(1 + \frac{1}{(\kappa_j - 1)2}\right)^{\frac{1}{2}} \right].$$

But

$$(2.20) \quad \min_{0 \leq x \leq 1 - \Delta} \left[(x + \Delta)^{\frac{1}{2}} - x^{\frac{1}{2}} \left(1 + \frac{1}{(\kappa_j - 1)2}\right)^{\frac{1}{2}} \right] = \\ = 1 - (1 - \Delta)^{\frac{1}{2}} \left(1 + \frac{1}{(\kappa_j - 1)2}\right)^{\frac{1}{2}}.$$

Therefore for κ_j big enough i.e. $j > N_1$ we have

$$(2.21) \quad \beta_j - \alpha_j \geq 2^{\frac{1}{2}} (\kappa_j - 1)^{\frac{1}{2}} C \quad \text{with } C > 0$$

and hence for $j \cdot N_2 > N_1$ we have $\beta_j - \alpha_j \geq 2$

and therefore there exists an integer k_j such that

$$\beta_j > k_j > \alpha_j \quad \text{for all } j > N_2$$

and hence $\xi_j = (\omega_j^2 + k_j^2)^{\frac{1}{2}}$ is a point of discontinuity of the function $U(\kappa)$.

But obviously

$$\xi_{i,j-1} < f_j < \xi_{i,j-1} + \Delta$$

and this is a contradiction and the theorem is proved.

A closer analysis of the proof shows that obviously by the same idea the general statement introduced in § 1 may be proved i.e. for n -dimensional domain with bounded curvature of the boundary.

R e f e r e n c e s

- [1] G. FIX and G. STRANG: Fourier analysis of the finite element method in Ritz-Galerkin theory, *Studies in Applied Mathematics* 48(1969),265-273.
- [2] G. STRANG and G. FIX: A Fourier analysis of the finite element variational method. To appear.
- [3] J.P. AUBIN: Behavior of the error of the approximate solution of boundary value problems for linear elliptic operators by Galerkin's and finite differences methods, *Annali della Scuola Normale di Pisa* 21(1967),599-637.
- [4] J.P. AUBIN: Some variational methods for non-homogeneous boundary value problems, *Calcolo*, 1969.
- [5] J.H. BRAMBLE and A.H. SCHATZ: Raleigh, Ritz-Galerkin methods for Dirichlet's problem using subspaces without boundary conditions, *Comm.Pure Appl.Math.* (to appear).
- [6] J. NITSCHKE: Über ein Variationsprinzip für Lösung von Dirichlet Problemen bei Verwendung von Teilräumen, die keinen Randbedingungen unterworfen sind (to appear).

- [7] J.H. BRAMBLE and A.H. SCHATZ: Least squares methods for 2nd order elliptic boundary value problems. To appear.
- [8] L.A. OGANESJAN, L.A. RUCHOVEC: A study of rates of convergence of some variational difference schemes for elliptic equations of second order in a two dimensional domain with smooth boundary (Russian). *Ž.Vyčislit.Mat.i Mat. Fiz.*9(1969),1102-1119.
- [9] I. BABUŠKA: Error bounds for finite element method, Tech.Note BN-630(1969),University of Maryland,Institute for Fluid Dynamics and Applied Mathematics. *Numerische Mathematik* 16, 322-333, 1971 .
- [10] I. BABUŠKA: The rate of convergence for the finite element method, Tech.Note BN-646(1970), University of Maryland,Institute for Fluid Dynamics and Applied Mathematics. To appear in *SIAM Journal,Num.Math.*
- [11] I. BABUŠKA: The finite element method for elliptic equations with discontinuous coefficients, Tech.Note BN-631(1969),University of Maryland,Institute for Fluid Dynamics and Applied Mathematics, *Computing* 5, 207-213(1970) .
- [12] I. BABUŠKA: Finite element method for domains with corners, Techn.Note BN-636(1970),University of Maryland,Institute for Fluid Dynamics and Applied Mathematics, to appear in *Computing*.

- [13] I. BABUŠKA: The finite element method for elliptic differential equations, Tech.Note BN-653, (1970), University of Maryland, Institute for Fluid Dynamics and Applied Mathematics. Numerical Solution of Partial Differential Equations II, SYNSPADE 1970. Edited by B. Hubbard, Academic Press 1971, New York-London, 69-107.
- [14] I. BABUŠKA: Computation of derivatives in the finite element method, Tech.Note BN-650(1970), University of Maryland, Institute for Fluid Dynamics and Applied Mathematics, Comment. Math.Univ.Carolinae 11,3(1970), 543-557.
- [15] I. BABUŠKA: Approximation by Hill functions, Tech. Note BN-648(1970), University of Maryland, Institute for Fluid Dynamics and Applied Mathematics, Comment.Math.Univ.Carolinae 11,4(1970), 787-811.
- [16] I. BABUŠKA: The finite element method for infinite domains I, Tech.Note BN-670(1970), University of Maryland, Inst.for Fluid Dynamics and Applied Mathematics.
- [17] I. BABUŠKA and J. SEGETHOVÁ and K. SEGETH: Numerical experiments with finite element method I, Tech.Note BN-669(1970), University of Maryland, Inst.for Fluid Dynamics and Applied Mathematics.

Institute for Fluid Dynamics and
Applied Mathematics
University of Maryland
College Park, Maryland
U.S.A.

(Oblatum 18.12.1970)