

Jaroslav Ježek

Upper semicomplements and a definable element in the lattice of groupoid varieties

Commentationes Mathematicae Universitatis Carolinae, Vol. 12 (1971), No. 3, 565--586

Persistent URL: <http://dml.cz/dmlcz/105366>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1971

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

UPPER SEMICOMPLEMENTS AND A DEFINABLE ELEMENT IN THE
LATTICE OF GROUPOID VARIETIES

Jaroslav JEŽEK, Praha

The variety of semigroups is not generated by any finite number of its proper subvarieties (see Dean and Evans [2]). An analogous statement holds for the lattices of varieties of groups, lattices, loops and commutative semigroups (see Evans [3] for the summary and bibliography). It is proved in [6] that this property is not shared by the variety of all universal algebras of a given type Δ containing at least one at least binary function symbol: there are found in the lattice \mathcal{L}_Δ of varieties of algebras of type Δ some upper semicomplements different from the greatest element ι_Δ of \mathcal{L}_Δ . In the present paper we shall restrict ourselves to the case of the lattice \mathcal{L}_Γ of groupoid varieties and investigate upper semicomplements in \mathcal{L}_Γ .

In § 2 the infimum of the set of all upper semicomplements in \mathcal{L}_Γ is found: it is just the variety of commutative semigroups satisfying $x^2 \cdot y = x \cdot y$. This variety is thus a definable element in \mathcal{L}_Γ .

To prove the result, we must find some further upper

semicomplements in \mathcal{L}_Γ . These are found in § 1.

For the terminology and notation see [6] and § 1 of [4].

§ 1. Some upper semicomplements in \mathcal{L}_Γ

We denote by Γ the type of groupoids, i.e. the type consisting of a single binary function symbol. The terminology given in [4] and [6] can be specialized to the case $\Delta = \Gamma$: e.g. W_Γ denotes the free groupoid freely generated by X . Γ -equations are called equations throughout the paper, etc. If μ and ν are two elements of W_Γ , then the value of the fundamental binary operation of W_Γ , applied to μ and ν , is denoted by $\mu \cdot \nu$ or only $\mu\nu$. We write $\mu\nu \cdot w$ instead of $(\mu \cdot \nu) \cdot w$, etc.

For every $t \in W_\Gamma$ we define two elements \overleftarrow{t} and \overrightarrow{t} of W_Γ in this way: if $t \in X$, then $\overleftarrow{t} = \overrightarrow{t} = t$; if $t = t_1 \cdot t_2$, then $\overleftarrow{t} = t_1$ and $\overrightarrow{t} = t_2$.

For every $t \in W_\Gamma$ we define elements $\sigma_1(t)$, $\sigma_2(t)$, $\sigma_3(t)$, ... of W_Γ in this way: $\sigma_1(t) = tt \cdot t$; $\sigma_{n+1}(t) = (\sigma_n(t) \cdot \sigma_n(t)) \cdot \sigma_n(t)$.

Let us fix two different variables (i.e. elements of X) and denote them by x_0 and y_0 . Put

$$e_1 = \langle (x_0 x_0 \cdot x_0) y_0, x_0 y_0 \rangle ; \quad e_2 = \langle (x_0 \cdot x_0 x_0) x_0, x_0 x_0 \rangle ;$$

$$e^1 = \langle x_0 x_0 \cdot x_0, x_0 x_0 \rangle ; \quad e^2 = \langle x_0 \cdot x_0 x_0, x_0 x_0 \rangle .$$

Let e be any of the four equations e_1, e_2, e^1 and e^2 . It will be useful to notice that the following (tri-

vial) assertion holds: whenever μ_1, μ_2, ν_1 and ν_2 are elements of W_Γ such that $\nu_1 \nu_2$ is a leap-consequence of $\mu_1 \mu_2$ by means of e , then no one of the three cases

(i) $\mu_1 \mu_2 = \nu_1 \nu_2$;

(ii) $\mu_1 = \nu_1$ and either $\mu_2 \in |C_e(\nu_2)$ or $\nu_2 \in |C_e(\mu_2)$;

(iii) $\mu_2 = \nu_2$ and either $\mu_1 \in |C_e(\nu_1)$ or $\nu_1 \in |C_e(\mu_1)$ can take place.

Let e be an arbitrary Γ -equation. We call an e -proof $\lceil t_1, \dots, t_m \rceil$ regular if either $t_i \in LC_e(t_{i+1})$ for all leaps i in $\lceil t_1, \dots, t_m \rceil$ or $t_{i+1} \in LC_e(t_i)$ for all leaps i in $\lceil t_1, \dots, t_m \rceil$. Evidently, if an e -proof has at most one leap, then it is regular.

Lemma 1. Let $a, b \in W_\Gamma$ and $e_1 \vdash \langle a, b \rangle$. Then there exists a regular e_1 -proof of b from a .

Proof. Let $\lceil \mu_1, \dots, \mu_m \rceil$ be an e_1 -proof of b from a with a minimal number of leaps. Suppose that it is not regular. Evidently, it has two leaps i and j ($i < j$) such that there is no leap greater than i and smaller than j (we say that i and j are two neighbouring leaps) and such that either

$$\mu_i = (\alpha \alpha \alpha) \beta \& \mu_{i+1} = \alpha \beta \& \mu_j = \gamma \sigma \& \mu_{j+1} = (\gamma \gamma \gamma) \sigma$$

or

$$\mu_i = \alpha \beta \& \mu_{i+1} = (\alpha \alpha \alpha) \beta \& \mu_j = (\gamma \gamma \gamma) \sigma \& \mu_{j+1} = \gamma \sigma$$

for some $\alpha, \beta, \gamma, \sigma \in W_\Gamma$. If $i + 1 = j$, then $\alpha = \gamma$ and $\beta = \sigma$, so that $\lceil \mu_1, \dots, \mu_i, \mu_{i+3}, \dots, \mu_m \rceil$

is an e_1 -proof of \mathcal{L} from \mathcal{A} which has a smaller number of leaps than $\lceil \mu_1, \dots, \mu_m \rceil$, a contradiction. Let $i + 1 < j$. In the first case

$$\begin{aligned} & \lceil \mu_1, \dots, \mu_i, ((\vec{u}_{i+2} \cdot \alpha) \alpha) \beta, \dots, ((\vec{u}_j \cdot \alpha) \alpha) \beta, \\ & ((\vec{u}_j \cdot \vec{u}_{i+2}) \alpha) \beta, \dots, ((\vec{u}_j \cdot \vec{u}_j) \alpha) \beta, \\ & ((\vec{u}_j \cdot \vec{u}_j) \vec{u}_{i+2}) \beta, \dots, ((\vec{u}_j \cdot \vec{u}_j) \vec{u}_j) \beta, ((\vec{u}_j \cdot \vec{u}_j) \vec{u}_j) \vec{u}_{i+2}, \dots, \\ & ((\vec{u}_j \cdot \vec{u}_j) \vec{u}_j) \vec{u}_j, \mu_{i+2}, \dots, \mu_m \rceil \end{aligned}$$

and in the second case

$$\lceil \mu_1, \dots, \mu_i, \vec{u}_{i+2} \cdot \vec{u}_{i+2}, \dots, \vec{u}_j \cdot \vec{u}_j, \mu_{i+2}, \dots, \mu_m \rceil$$

is an e_1 -proof of \mathcal{L} from \mathcal{A} and it has a smaller number of leaps than $\lceil \mu_1, \dots, \mu_m \rceil$, a contradiction.

Lemma 2. Let $a_1, a_2, \mathcal{L}_1, \mathcal{L}_2 \in W_T$. Then $e_1 \vdash \langle a_1 a_2, \mathcal{L}_1 \mathcal{L}_2 \rangle$ if and only if $e_1 \vdash \langle a_2, \mathcal{L}_2 \rangle$ and one of the following three cases takes place:

- (i) $e_1 \vdash \langle a_1, \mathcal{L}_1 \rangle$;
- (ii) $e_1 \vdash \langle a_1, \mathcal{G}_n(\mathcal{L}_1) \rangle$ for some $n \geq 1$;
- (iii) $e_1 \vdash \langle \mathcal{L}_1, \mathcal{G}_n(a_1) \rangle$ for some $n \geq 1$.

Proof follows easily from Lemma 1.

Lemma 3. For every $t \in W_T$ denote by φ_t the endomorphism of W_T assigning t to every variable. Let $x \in X$, $a \in W_T$ and $w \in T_T(x)$; let $w \neq x$. Then $\{e_1, e_2\} \vdash \langle a, \varphi_a(w) \rangle$ does not hold.

Proof by the induction on a . Everything is evident if $a \in X$. Let $a \notin X$ and suppose $\{e_1, e_2\} \vdash \langle a, \varphi_a(w) \rangle$. Evidently, there exists a finite sequence w_1, \dots, w_m such that $w_1 = w$, $w_m = x$ and $w_{i+1} = \vec{w}_i$ for every

$i = 1, \dots, m-1$. We have evidently $\{e_1, e_2\} \vdash \langle \overline{a^i}, g_a(w_2) \rangle$;
 from this $\{e_1, e_2\} \vdash \langle \overline{a^i}, g_a(w_3) \rangle$; etc; finally,
 $\{e_1, e_2\} \vdash \langle l, g_a(w_m) \rangle = \langle l, a \rangle$ for some $l \in S(\overline{a})$,
 so that $\{e_1, e_2\} \vdash \langle l, g_a(w) \rangle$, a contradiction with
 the induction assumption.

Lemma 4. Let $a, l \in W_\Gamma$ and $e_2 \vdash \langle a, l \rangle$. Then there exists an e_2 -proof of l from a which has at most one leap.

Proof. Let $\Gamma \mu_1, \dots, \mu_m$ be an e_2 -proof of l from a with a minimal number of leaps. Suppose that it has at least two leaps. Then it has two neighbouring leaps i and j ($i < j$). Four cases are possible:

(1) There exist $\alpha, \beta \in W_\Gamma$ such that
 $\mu_i = (\alpha \cdot \alpha \alpha) \alpha$ & $\mu_{i+1} = \alpha \alpha$ & $\mu_j = (\beta \cdot \beta \beta) \beta$ & $\mu_{j+1} = \beta \beta$;
 then $e_2 \vdash \langle \alpha, \beta \cdot \beta \beta \rangle$ and $e_2 \vdash \langle \alpha, \beta \rangle$, so that
 $e_2 \vdash \langle \beta, \beta \cdot \beta \beta \rangle$, a contradiction with Lemma 3.

(2) There exist $\alpha, \beta \in W_\Gamma$ such that
 $\mu_i = \alpha \alpha$ & $\mu_{i+1} = (\alpha \cdot \alpha \alpha) \alpha$ & $\mu_j = \beta \beta$ & $\mu_{j+1} = (\beta \cdot \beta \beta) \beta$;
 then $e_2 \vdash \langle \alpha, \alpha \cdot \alpha \alpha \rangle$, a contradiction.

(3) and (4) The remaining two cases give a contradiction similarly as in the proof of Lemma 1.

Lemma 5. Let $a_1, a_2, l_1, l_2 \in W_\Gamma$. Then
 $e_2 \vdash \langle a_1 a_2, l_1 l_2 \rangle$ if and only if $e_2 \vdash \langle a_2, l_2 \rangle$
 and one of the following three cases takes place:

- (i) $e_2 \vdash \langle a_1, l_1 \rangle$;
- (ii) $e_2 \vdash \langle a_1, a_2 \rangle$ and $e_2 \vdash \langle l_1, a_1 \cdot a_1 a_1 \rangle$;
- (iii) $e_2 \vdash \langle l_1, l_2 \rangle$ and $e_2 \vdash \langle a_1, l_1 \cdot l_1 l_1 \rangle$.

Proof follows easily from Lemma 4.

Lemma 6. Let $\alpha, \beta \in W_T$. Then neither $\{e_1, e_2\} \vdash \langle \alpha\alpha.\alpha, \beta.\beta/\beta \rangle$ nor $\{e_1, e_2\} \vdash \langle \alpha\alpha.\alpha, \beta/\beta \rangle$ takes place.

Proof. Suppose on the contrary that there exist elements $\alpha, \beta \in W_T$ and an $\{e_1, e_2\}$ -proof $\lceil \mu_1, \dots, \mu_n \rceil$ such that the following holds: $\mu_1 = \alpha\alpha.\alpha$; either $\mu_n = \beta.\beta/\beta$ or $\mu_n = \beta/\beta$; whenever $\gamma, \sigma \in W_T$ and $\lceil \nu_1, \dots, \nu_m \rceil$ is an $\{e_1, e_2\}$ -proof of either $\sigma.\sigma/\sigma$ or σ/σ from $\gamma\gamma.\gamma$, then $m \leq n$. This $\lceil \mu_1, \dots, \mu_n \rceil$ has leaps, for if it had not, then in case $\mu_n = \beta.\beta/\beta$ we would have $\{e_1, e_2\} \vdash \langle \alpha\alpha, \beta \rangle$ and $\{e_1, e_2\} \vdash \langle \alpha, \beta/\beta \rangle$, so that $\{e_1, e_2\} \vdash \langle \alpha, \alpha\alpha.\alpha\alpha \rangle$; and in case $\mu_n = \beta/\beta$ we would have $\{e_1, e_2\} \vdash \langle \alpha\alpha, \beta \rangle$ and $\{e_1, e_2\} \vdash \langle \alpha, \beta \rangle$, so that $\{e_1, e_2\} \vdash \langle \alpha, \alpha\alpha \rangle$, a contradiction with Lemma 3. Let i be the first leap in $\lceil \mu_1, \dots, \mu_n \rceil$.

If $\mu_i = (\kappa\kappa.\kappa)\kappa$ & $\mu_{i+1} = \kappa\kappa$ for some $\kappa, \kappa \in W_T$, then $\lceil \mu_i, \mu_{i-1}, \dots, \mu_1 \rceil$ is an $\{e_1, e_2\}$ -proof of $\alpha\alpha$ from $\kappa\kappa.\kappa$, and $i < n$ gives a contradiction.

If $\mu_i = (\kappa.\kappa\kappa)\kappa$ & $\mu_{i+1} = \kappa\kappa$, then $\{e_1, e_2\} \vdash \langle \alpha\alpha, \kappa.\kappa\kappa \rangle$ and $\{e_1, e_2\} \vdash \langle \alpha, \kappa \rangle$, so that $\{e_1, e_2\} \vdash \langle \alpha, \alpha\alpha \rangle$, a contradiction with Lemma 3.

If $\mu_i = \kappa\kappa$ & $\mu_{i+1} = (\kappa.\kappa\kappa)\kappa$, then $\{e_1, e_2\} \vdash \langle \alpha, \alpha\alpha \rangle$, a contradiction.

Let us call a leap l in $\lceil \mu_1, \dots, \mu_n \rceil$ a \ast -leap if there exist $\kappa, \kappa \in W_T$ such that $\mu_l = \kappa\kappa$ & $\mu_{l+1} =$

$= (\mathcal{N}\mathcal{N}\mathcal{N}) \circ$. We have proved that i is a $*$ -leap.
 Suppose that every leap in $\lceil \mu_1, \dots, \mu_m \rceil$ is a $*$ -leap.
 Then $\{e_1, e_2\} \vdash \langle \beta, \bar{\sigma}_m(\alpha\alpha) \rangle$ for some $m \geq 1$;
 in case $\mu_m = \beta \cdot \beta \beta$ we have further
 $\{e_1, e_2\} \vdash \langle \alpha, \beta \beta \rangle$, so that $\{e_1, e_2\} \vdash \langle \alpha, \bar{\sigma}_m(\alpha\alpha), \bar{\sigma}_m(\alpha\alpha) \rangle$,
 a contradiction; in case $\mu_m = \beta \beta$ we have
 $\{e_1, e_2\} \vdash \langle \alpha, \beta \rangle$, so that $\{e_1, e_2\} \vdash \langle \alpha, \bar{\sigma}_m(\alpha\alpha) \rangle$,
 a contradiction again. This proves that $\lceil \mu_1, \dots, \mu_m \rceil$ has
 two neighbouring leaps j and k ($j < k$) such that k
 is not a $*$ -leap and j is a $*$ -leap. There exist a ,
 $b \in W_\Gamma$ such that $\mu_j = ab$ & $\mu_{j+1} = (a \cdot a) b$.

Suppose $\mu_k = (cc \cdot c)d$ & $\mu_{k+1} = cd$ for some c ,
 $d \in W_\Gamma$. Then
 $\lceil \mu_1, \dots, \mu_j, \vec{\mu}_{j+2}, \vec{\mu}_{j+2}, \dots, \vec{\mu}_k, \vec{\mu}_k, \mu_{k+2}, \dots, \mu_m \rceil$
 is an $\{e_1, e_2\}$ -proof, a contradiction with the minimal
 property of $\lceil \mu_1, \dots, \mu_m \rceil$.

Suppose $\mu_k = (c \cdot cc)c$ & $\mu_{k+1} = cc$. Then
 $\lceil \vec{\mu}_{j+1}, \dots, \vec{\mu}_k \rceil$ is an $\{e_1, e_2\}$ -proof of $c \cdot cc$ from
 $a \cdot a \cdot a$, a contradiction with the minimal property of
 $\lceil \mu_1, \dots, \mu_m \rceil$.

The case $\mu_k = cc$ & $\mu_{k+1} = (c \cdot cc)c$ remains.
 $\lceil \mu_1, \dots, \mu_k \rceil$ is an $\{e_1, e_2\}$ -proof of cc from $\alpha\alpha \cdot \alpha$,
 again a contradiction with the minimal property of
 $\lceil \mu_1, \dots, \mu_m \rceil$.

Lemma 7. $Cn(e_1) \vee_\Gamma Cn(e_2) = L_\Gamma$.

Proof. Let us prove the following assertion by in-
 duction on a : whenever $a, b \in W_\Gamma$, $e_1 \vdash \langle a, b \rangle$ and

$e_2 \vdash \langle a, b \rangle$, then $a = b$. This is evident if $a \in X$. Let $a = a_1 a_2$.

Evidently, $b \notin X$; put $b = b_1 b_2$. We get $a_2 = b_2$ easily from the induction assumption, so that it is enough to prove $a_1 = b_1$.

Let $e_1 \vdash \langle a_1, b_1 \rangle$. By Lemma 5, the following three cases are the only possible ones:

(1) $e_2 \vdash \langle a_1, b_1 \rangle$. Then we get $a_1 = b_1$ from the induction assumption.

(2) $e_2 \vdash \langle a_1, a_2 \rangle \& e_2 \vdash \langle b_1, a_1 \cdot a_1 a_1 \rangle$. As $\{e_1, e_2\} \vdash \langle a_1, a_1 \cdot a_1 a_1 \rangle$, we get a contradiction with Lemma 3.

(3) $e_2 \vdash \langle b_1, b_2 \rangle \& e_2 \vdash \langle a_1, b_1 \cdot b_1 b_1 \rangle$. Again, $\{e_1, e_2\} \vdash \langle b_1, b_1 \cdot b_1 b_1 \rangle$, a contradiction.

Let $e_1 \vdash \langle a_1, \sigma_n(b_1) \rangle$ for some $n \geq 1$. (1), (2) and (3) are again the only possible cases. In cases (1) and (2) we get a contradiction with Lemma 3. In case (3) we get a contradiction with Lemma 6 and the definition of σ_n .

By Lemma 5, the case $e_1 \vdash \langle b_1, \sigma_n(a_1) \rangle$ remains. This case is similar to $e_1 \vdash \langle a_1, \sigma_n(b_1) \rangle$.

Lemma 8. If $a \in W_p$, then $e^1 \vdash \langle a, a a \rangle$ does not hold.

Proof by induction on a . It is evident if $a \in X$. Let $a = a_1 a_2$ and suppose $e^1 \vdash \langle a, a a \rangle$. Evidently, $e^1 \vdash \langle a_2, a \rangle$, so that $e^1 \vdash \langle a_2, a_2 a_2 \rangle$ which contradicts to the induction assumption.

Lemma 9. Let $a, b \in W_p$ and $e^1 \vdash \langle a, b \rangle$. Then

there exists an e^1 -proof of \mathcal{L} from a which has at most one leap.

Proof. Let $\ulcorner \mu_1, \dots, \mu_n \urcorner$ be an e^1 -proof of \mathcal{L} from a with a minimal number of leaps. Suppose that it has at least two leaps, so that it has two neighbouring leaps i and j ($i < j$). There are four cases:

(1) $\mu_i = \alpha\alpha$ & $\mu_{i+1} = \alpha\alpha$, α & $\mu_j = \beta\beta$ & $\mu_{j+1} = \beta\beta$, β
 for some $\alpha, \beta \in W_T$. Then $e^1 \vdash \langle \alpha\alpha, \beta \rangle$ and $e^1 \vdash \langle \alpha, \beta \rangle$, so that $e^1 \vdash \langle \alpha, \alpha\alpha \rangle$, a contradiction with Lemma 8.

(2) $\mu_i = \alpha\alpha$, α & $\mu_{i+1} = \alpha\alpha$ & $\mu_j = \beta\beta$, β & $\mu_{j+1} = \beta\beta$.
 We can get a contradiction similarly as in the preceding case.

(3) $\mu_i = \alpha\alpha$ & $\mu_{i+1} = \alpha\alpha$, α & $\mu_j = \beta\beta$, β & $\mu_{j+1} = \beta\beta$.
 Then

$\ulcorner \mu_1, \dots, \mu_i, \overrightarrow{\mu_{i+2}}, \alpha, \dots, \overrightarrow{\mu_j}, \alpha, \overrightarrow{\mu_j}, \overrightarrow{\mu_{i+2}}, \dots, \overrightarrow{\mu_j}, \overrightarrow{\mu_j}, \mu_{j+2}, \dots, \mu_n \urcorner$
 is an e^1 -proof of \mathcal{L} from a which has a smaller number of leaps than $\ulcorner \mu_1, \dots, \mu_n \urcorner$, a contradiction.

(4) $\mu_i = \alpha\alpha$, α & $\mu_{i+1} = \alpha\alpha$ & $\mu_j = \beta\beta$ & $\mu_{j+1} = \beta\beta$, β .
 Then

$\ulcorner \mu_1, \dots, \mu_i, \mu_{i+2}, \alpha, \dots, \mu_j, \alpha, \mu_j, \overrightarrow{\mu_{i+2}}, \dots, \mu_j, \overrightarrow{\mu_j}, \mu_{j+2}, \dots, \mu_n \urcorner$
 is an e^1 -proof of \mathcal{L} from a which has a smaller number of leaps, a contradiction again.

Lemma 10. Let $a_1, a_2, \mathcal{L}_1, \mathcal{L}_2 \in W_T$. Then $e^1 \vdash \langle a_1 a_2, \mathcal{L}_1 \mathcal{L}_2 \rangle$ if and only if $e^1 \vdash \langle a_2, \mathcal{L}_2 \rangle$ and one of the following three cases takes place:

(i) $e^1 \vdash \langle a_1, \mathcal{L}_1 \rangle$;

(ii) $e^1 \vdash \langle l_1, l_2 \rangle$ and $e^1 \vdash \langle a_1, l_1 l_1 \rangle$;

(iii) $e^1 \vdash \langle a_1, a_2 \rangle$ and $e^1 \vdash \langle l_1, a_1 a_1 \rangle$.

Proof follows easily from Lemma 9.

Lemma 11. Let $a_1, a_2, l_1, l_2 \in W_T$. Then $e^2 \vdash \langle a_1 a_2, l_1 l_2 \rangle$ if and only if $e^2 \vdash \langle a_1, l_1 \rangle$ and one of the following three cases takes place:

(i) $e^2 \vdash \langle a_2, l_2 \rangle$;

(ii) $e^2 \vdash \langle l_1, l_2 \rangle$ and $e^2 \vdash \langle a_2, l_2 l_2 \rangle$;

(iii) $e^2 \vdash \langle a_1, a_2 \rangle$ and $e^2 \vdash \langle l_2, a_2 a_2 \rangle$.

Proof is similar to that of Lemma 10.

Lemma 12. Let $a, b \in W_T$. If $\{e^1, e^2\} \vdash \langle a a, b b \rangle$, then $\{e^1, e^2\} \vdash \langle a, b \rangle$, too.

Proof. Suppose that it is not true. There exists an $\{e^1, e^2\}$ -proof $\lceil \mu_1, \dots, \mu_n \rceil$ such that the following holds: there exist $\alpha, \beta \in W_T$ satisfying $\mu_1 = \alpha \alpha$ and $\mu_n = \beta \beta$ and not satisfying $\{e^1, e^2\} \vdash \langle \alpha, \beta \rangle$; whenever $\lceil \nu_1, \dots, \nu_m \rceil$ is an $\{e^1, e^2\}$ -proof with a similar property, then $m \leq n$. Choose such a minimal $\lceil \mu_1, \dots, \mu_n \rceil$ and put $\mu_1 = a a$ and $\mu_2 = b b$. Suppose $\mu_i = c c$ for some i such that $2 \leq i \leq n-1$. As $\lceil \mu_1, \dots, \mu_i \rceil$ is an $\{e^1, e^2\}$ -proof of $c c$ from $a a$ and $i < n$, we have $\{e^1, e^2\} \vdash \langle a, c \rangle$; as $\lceil \mu_i, \dots, \mu_n \rceil$ is an $\{e^1, e^2\}$ -proof of $b b$ from $c c$ and $n-i+1 < n$, we have $\{e^1, e^2\} \vdash \langle b, c \rangle$. Consequently, $\{e^1, e^2\} \vdash \langle a, b \rangle$, a contradiction. From this we infer that no numbers other than 1 and $n-1$ can be leaps in $\lceil \mu_1, \dots, \mu_n \rceil$. If $\lceil \mu_1, \dots, \mu_n \rceil$ had at most one

leap, then either $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_n$ or $\overrightarrow{\mu}_1, \dots, \overrightarrow{\mu}_n$ would be an $\{e^1, e^2\}$ -proof of l from a ; hence, the numbers 1 and $n - 1$ are leaps. We have either $\mu_2 = aa.a$ or $\mu_2 = a.aa$. It is sufficient to consider the case $\mu_2 = aa.a$. If it were $\mu_{n-1} = l.l.l$, then $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_n$ would be an $\{e^1, e^2\}$ -proof of l from a . We get $\mu_{n-1} = l.l.l$. Evidently, $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_{n-1}$ is an $\{e^1, e^2\}$ -proof of $l.l$ from a and $\overleftarrow{\mu}_2, \dots, \overleftarrow{\mu}_n$ is an $\{e^1, e^2\}$ -proof of l from aa . As $\{e^1, e^2\} \vdash \langle l.l, a, a \rangle$, we get $\{e^1, e^2\} \vdash \langle a, l \rangle$, a contradiction.

Lemma 13. If $a \in W_\Gamma$, then $\{e^1, e^2\} \vdash \langle a, aa \rangle$ does not hold.

Proof by induction on a . It is evident if $a \in X$. Let $a = a_1 a_2$ and suppose $\{e^1, e^2\} \vdash \langle a, aa \rangle$. Let $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_n$ be an arbitrary $\{e^1, e^2\}$ -proof of a from aa .

Suppose that $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_n$ has a leap. Denote by k its last leap. If it were $\mu_{k+1} = cc$ for some $c \in W_\Gamma$, then we would get $\{e^1, e^2\} \vdash \langle a_1, c \rangle$; as $\{e^1, e^2\} \vdash \langle aa, cc \rangle$, Lemma 12 gives $\{e^1, e^2\} \vdash \langle a, c \rangle$; hence, $\{e^1, e^2\} \vdash \langle a_1, a \rangle$, so that $\{e^1, e^2\} \vdash \langle a_1, a_1 a_1 \rangle$, a contradiction with the induction hypothesis. This proves $\mu_k = cc$ for some c and either $\mu_{k+1} = cc.c$ or $\mu_{k+1} = c.cc$. Again, from $\{e^1, e^2\} \vdash \langle aa, cc \rangle$ follows by Lemma 12 $\{e^1, e^2\} \vdash \langle a, c \rangle$. In case $\mu_{k+1} = cc.c$ we have $\{e^1, e^2\} \vdash \langle c, a_2 \rangle$, so that $\{e^1, e^2\} \vdash \langle a, a_2 \rangle$ and consequently $\{e^1, e^2\} \vdash \langle a_2, a_2 a_2 \rangle$, a contradiction

with the induction hypothesis; in case $\mu_{n+1} = c.c.c$ similarly $\{e^1, e^2\} \vdash \langle a_1, a_1 a_1 \rangle$, a contradiction again.

We have proved that $\lceil \mu_1, \dots, \mu_n \rceil$ has no leaps. $\lceil \overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_n \rceil$ is an $\{e^1, e^2\}$ -proof of a_1 from a , so that $\{e^1, e^2\} \vdash \langle a_1, a_1 a_1 \rangle$, a contradiction with the induction hypothesis.

Lemma 14. $Cn(e^1) \vee_{\Gamma} Cn(e^2) = \cup_{\Gamma}$.

Proof. We shall prove by induction on a the following: whenever $e^1 \vdash \langle a, b \rangle$ and $e^2 \vdash \langle a, b \rangle$, then $a = b$. This is evident if $a \in X$. Let $a = a_1 a_2$, $e^1 \vdash \langle a, b \rangle$, $e^2 \vdash \langle a, b \rangle$ and $a \neq b$. Evidently, $b \notin X$, put $b = b_1 b_2$. We have $e^1 \vdash \langle a_2, b_2 \rangle$ and $e^2 \vdash \langle a_1, b_1 \rangle$; it is sufficient to prove $e^1 \vdash \langle a_1, b_1 \rangle$ and $e^2 \vdash \langle a_2, b_2 \rangle$. Suppose on the contrary e.g. that $e^1 \vdash \langle a_1, b_1 \rangle$ does not hold. We have either $e^1 \vdash \langle b_1, b_2 \rangle$ & $e^1 \vdash \langle a_1, b_1 b_2 \rangle$ or $e^1 \vdash \langle a_1, a_2 \rangle$ & $e^1 \vdash \langle b_1, a_1 a_1 \rangle$ by Lemma 10. Evidently, $\{e^1, e^2\} \vdash \langle a_1, a_1 a_1 \rangle$ in both cases, a contradiction with Lemma 13.

Lemma 15. Let x and y be two different variables. Then every minimal $\langle x x . y, x . y x \rangle$ -proof is regular.

Proof. Put $e = \langle x x . y, x . y x \rangle$. We shall prove by induction on n that every minimal e -proof $\lceil \mu_1, \dots, \mu_n \rceil$ is regular. This is evident if $n = 1$. Let $n > 1$. Suppose that $\lceil \mu_1, \dots, \mu_n \rceil$ is not regular, so that it has two neighbouring leaps i and j ($i < j$) such that one of the following two cases takes place:

(1) $\mu_i = a . b a$ & $\mu_{i+1} = a a . b$ & $\mu_j = c c . d$ & $\mu_{j+1} = c . d c$ for some $a, b, c, d \in W_{\Gamma}$. We have $e \vdash \langle a a, c c \rangle$, so that $l(a a) = l(c c)$ and thus $l(a) = l(c)$. The e -proof

$\overleftarrow{\mu}_{i+1}, \dots, \overleftarrow{\mu}_j$ of cc from aa is minimal if we leave out its members $\overleftarrow{\mu}_m$ such that $\overleftarrow{\mu}_m = \overleftarrow{\mu}_{m-1}$; by the induction assumption it follows easily from $l(a) = l(c)$ that $\overleftarrow{\mu}_{i+1}, \dots, \overleftarrow{\mu}_j$ has no leaps. Consequently, $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_i, \overleftarrow{\mu}_{i+2}, (\overleftarrow{\mu}_{i+2}, \overleftarrow{\mu}_{i+2}), \dots, \overleftarrow{\mu}_j, (\overleftarrow{\mu}_j, \overleftarrow{\mu}_j), \mu_{j+2}, \dots, \mu_m$ is an e-proof of μ_m from μ_1 , a contradiction with the minimality of $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_m$.

(2) $\mu_i = a.a.l$ & $\mu_{i+1} = a.l.a$ & $\mu_j = c.d.c$ & $\mu_{j+1} = c.c.d$ for some $a, l, c, d \in W_\Gamma$. We have $e \vdash \langle a, c \rangle$ and $e \vdash \langle l.a, d.c \rangle$, so that $l(a) = l(c)$ and $l(l.a) = l(d.c)$; we infer $l(l) = l(d)$. Similarly as in the previous case, $\overleftarrow{\mu}_{i+1}, \dots, \overleftarrow{\mu}_j$ has no leaps and $\overleftarrow{\mu}_1, \dots, \overleftarrow{\mu}_i, (\overleftarrow{\mu}_{i+2}, \overleftarrow{\mu}_{i+2}), \overleftarrow{\mu}_{i+2}, \dots, (\overleftarrow{\mu}_j, \overleftarrow{\mu}_j), \overleftarrow{\mu}_j, \mu_{j+2}, \dots, \mu_m$ is a shorter proof of μ_m from μ_1 , a contradiction.

Lemma 16. Let x and y be two different variables.

Then

$$Cm(\langle x.x.y, x.y.x \rangle) \vee_\Gamma Cm(\langle x.(x.x.x), (x.x.x).x \rangle) = L_\Gamma.$$

Proof. Put $e = \langle x.x.y, x.y.x \rangle$ and $\bar{e} = \langle x.(x.x.x), (x.x.x).x \rangle$. Let $a, b \in W_\Gamma$, $e \vdash \langle a, b \rangle$ and $\bar{e} \vdash \langle a, b \rangle$. Suppose that a minimal e-proof of b from a has leaps. Using Lemma 15, there exists a natural number $n \geq 1$ such that either $l(\bar{a}) = 2^n \cdot l(\bar{b})$ or $l(\bar{b}) = 2^n \cdot l(\bar{a})$. By Lemma 1 of [6], a minimal \bar{e} -proof of b from a has at most one leap. If it has a leap, we have either $l(\bar{a}) = 3 \cdot l(\bar{b})$ or $l(\bar{b}) = 3 \cdot l(\bar{a})$; if it has not, we have $l(\bar{a}) = l(\bar{b})$. This gives a contradiction in each case, as neither $2^n = 3$ nor $2^n = \frac{1}{3}$ nor $2^n = 1$.

We have proved that a minimal e -proof of b from a has no leaps. This implies $l(\vec{a}) = l(\vec{b})$ and a minimal \bar{e} -proof of b from a has no leaps, too. If we had proved the equality by induction on a , we should get $\vec{a} = \vec{b}$ and $\vec{a} = \vec{b}$, so that $a = b$.

Lemma 17. Let x and y be two different variables; put $e = \langle x.yx, x.y.x \rangle$. Then every minimal e -proof has at most one leap.

Proof. We shall prove by induction on n that every minimal e -proof $\ulcorner \mu_1, \dots, \mu_n \urcorner$ has at most one leap. This is evident if $n = 1$. Let $n > 1$ and suppose that a minimal e -proof $\ulcorner \mu_1, \dots, \mu_n \urcorner$ has at least two leaps. It has two neighbouring leaps i and j ($i < j$); one of the following four cases takes place:

(1) $\mu_i = a.bra$ & $\mu_{i+1} = a.br.a$ & $\mu_j = c.dc$ & $\mu_{j+1} = cd.c$
 for some $a, b, c, d \in W_T$. We have $e \vdash \langle a.br, c \rangle$ and $e \vdash \langle a, dc \rangle$, so that $l(abr) = l(c)$ and $l(a) = l(dc)$ and consequently $l(abr) < l(a)$, which is impossible.

(2) $\mu_i = a.br.a$ & $\mu_{i+1} = a.br.a$ & $\mu_j = cd.c$ & $\mu_{j+1} = c.dc$;
 a contradiction can be derived similarly.

(3) $\mu_i = a.bra$ & $\mu_{i+1} = a.br.a$ & $\mu_j = cd.c$ & $\mu_{j+1} = c.dc$.
 We have $l(abr) = l(cd)$ and $l(a) = l(c)$ and consequently $l(b) = l(d)$, too. By the induction hypothesis, this implies that $\ulcorner \mu_{i+1}, \dots, \mu_j \urcorner$ has no leaps, so that $\ulcorner \mu_1, \dots, \mu_i, \overleftarrow{\mu_{i+1}}, \overleftarrow{\mu_{i+2}}, \dots, \overleftarrow{\mu_j}, (\overleftarrow{\mu_j}, \overleftarrow{\mu_j}), \mu_{j+2}, \dots, \mu_n \urcorner$ is a shorter e -proof of μ_n from μ_1 , a contradiction.

(4) $\mu_i = a.br.a$ & $\mu_{i+1} = a.br.a$ & $\mu_j = c.dc$ & $\mu_{j+1} = cd.c$;
 we can get a contradiction similarly.

Lemma 18. Let x and y be two different variables.

Then

$$Cn(\langle x.yx, xy.x \rangle) \vee_{\tau} Cn(\langle x.(xx.xx), (xx.xx).x \rangle) = \zeta_{\tau}.$$

Proof. Put $e = \langle x.yx, xy.x \rangle$ and $\bar{e} = \langle x.(xx.xx), (xx.xx).x \rangle$. We prove the following by induction on

a : whenever $e \vdash \langle a, b \rangle$ and $\bar{e} \vdash \langle a, b \rangle$, then $a = b$. This is evident if $a \in X$. Let $a = a_1 a_2$, $e \vdash \langle a, b \rangle$ and $\bar{e} \vdash \langle a, b \rangle$. Evidently, $b \notin X$; put $b = b_1 b_2$. Let $\ulcorner \mu_1, \dots, \mu_m \urcorner$ be a minimal \bar{e} -proof of b from a . By Lemma 1 of [6] it has at most one leap.

Suppose that $\ulcorner \mu_1, \dots, \mu_m \urcorner$ has exactly one leap i .

It is sufficient to consider only the case

$\mu_i = \alpha.(xx.xx)$ & $\mu_{i+1} = (xx.xx).\alpha$ for some $\alpha \in W_{\tau}$. As $l(\alpha x) = l(x\alpha)$, the \bar{e} -proof $\ulcorner \bar{\mu}_{i+1}, \dots, \bar{\mu}_m \urcorner$ has no leaps. Hence, $l(b_1) = 4.l(a_1)$, $b_1 \notin X$ and $l(\bar{b}_1) = 2.l(a_1) = l(\bar{a}_1)$. Let $\ulcorner \nu_1, \dots, \nu_m \urcorner$ be a minimal e -proof of b from a . As $l(a_1) < l(b_1)$, $\ulcorner \nu_1, \dots, \nu_m \urcorner$ has leaps; by Lemma 17, it has exactly one leap j ; evidently, there exist $\beta, \gamma \in W_{\tau}$ such that $\nu_j = \beta.\gamma\beta$ & $\nu_{j+1} = \beta\gamma.\beta$. As $\ulcorner \bar{\nu}_{j+1}, \dots, \bar{\nu}_m \urcorner$ is (after leaving its members $\bar{\nu}_k$ such that $\bar{\nu}_k = \bar{\nu}_{k-1}$) a minimal e -proof, it has at most one leap; as $l(\beta) = l(a_1)$ and $l(\bar{b}_1) = 2.l(a_1)$, it has exactly one leap k and there exist ϵ and σ such that $\bar{\nu}_k = \sigma.\epsilon\sigma$ & $\bar{\nu}_{k+1} = \sigma\epsilon.\sigma$. We get $l(\bar{b}_1) = l(\sigma) = l(\beta) = l(a_1)$, a contradiction with $l(\bar{b}_1) = 2.l(a_1)$.

We have proved that $\ulcorner \mu_1, \dots, \mu_m \urcorner$ has no leaps and consequently $\bar{e} \vdash \langle a_1, b_1 \rangle$ and $\bar{e} \vdash \langle a_2, b_2 \rangle$. As

$l(a_1) = l(b_1)$, a minimal e -proof of b from a has no leaps, too, so that $e \vdash \langle a_1, b_1 \rangle$ and $e \vdash \langle a_2, b_2 \rangle$. The induction assumption gives $a_1 = b_1$ and $a_2 = b_2$, so that $a = b$.

Lemma 19. Let x, y and z be three different variables; put $e = \langle ((x \cdot x y) z) x, x(x(y z \cdot z)) \rangle$. Then every minimal e -proof has at most one leap.

Proof. We prove by induction on n for every minimal e -proof $\ulcorner u_1, \dots, u_n \urcorner$ that it has at most one leap. The case $n = 1$ is evident; let $n > 1$ and suppose that $\ulcorner u_1, \dots, u_n \urcorner$ has at least two leaps. It has two neighbouring leaps i and j ($i < j$); one of the following four cases takes place:

(1) $u_i = ((a \cdot a b) c) c$ & $u_{i+1} = a(a(bc \cdot c))$ & $u_j = ((p \cdot p q) \kappa) \kappa$ & $u_{j+1} = p(p(q \kappa \cdot \kappa))$ for some $a, b, c, p, q, \kappa \in W_\Gamma$. We have

$l(a) = l((p \cdot p q) \kappa) > l(\kappa) = l(a(bc \cdot c)) > l(a)$, a contradiction.

(2) $u_i = a(a(bc \cdot c))$ & $u_{i+1} = ((a \cdot a b) c) c$ & $u_j = p(p(q \kappa \cdot \kappa))$ & $u_{j+1} = ((p \cdot p q) \kappa) \kappa$; we get a contradiction similarly.

(3) $u_i = ((a \cdot a b) c) c$ & $u_{i+1} = a(a(bc \cdot c))$ & $u_j = p(p(q \kappa \cdot \kappa))$ & $u_{j+1} = ((p \cdot p q) \kappa) \kappa$. We have $l(a) = l(p)$ and $l(a(bc \cdot c)) = l(p(q \kappa \cdot \kappa))$, so that $l(bc \cdot c) = l(q \kappa \cdot \kappa)$. As the e -proof $\ulcorner \vec{u}_{i+1}, \dots, \vec{u}_j \urcorner$ is (after leaving out some members) minimal, it has no leaps by the induction assumption. Hence, $\ulcorner \vec{u}_{i+1}, \dots, \vec{u}_j \urcorner$ is an e -proof and it is minimal if we leave out some members; as $l(bc) > l(c)$ and $l(q \kappa) > l(\kappa)$, it has no leaps.

We get $l(bc) = l(q\kappa)$ and $l(c) = l(\kappa)$, so that $l(b) = l(q)$, too. Again, the e -proof

$\ulcorner \overline{\mu}_{i+1}, \dots, \overline{\mu}_j \urcorner$ has no leaps. Evidently,

$$\ulcorner \mu_1, \dots, \mu_i, ((\overline{\mu}_{i+2} \cdot (\overline{\mu}_{i+2} \cdot \overline{\mu}_{i+2})) \cdot \overline{\mu}_{i+2}) \cdot \overline{\mu}_{i+2}, \dots, ((\overline{\mu}_j \cdot (\overline{\mu}_j \cdot \overline{\mu}_j)) \cdot \overline{\mu}_j) \cdot \overline{\mu}_j, \mu_{j+2}, \dots, \mu_n \urcorner$$

is a shorter e -proof of μ_n from μ_1 , a contradiction.

(4) The last case is similar to the previous one.

Lemma 20. Let x, y and z be three different variables. Then

$$Cn(\langle\langle\langle(x \cdot xy)x\rangle x, x(x(yx \cdot z))\rangle\rangle \vee_p Cn(\langle\langle x \cdot xx, xx \cdot x \rangle\rangle) = \mathcal{L}_p.$$

Proof. Put $e = \langle\langle\langle(x \cdot xy)x\rangle x, x(x(yz \cdot z))\rangle\rangle$

and $\overline{e} = \langle x \cdot xx, xx \cdot x \rangle$. We prove by induction on a : whenever $e \vdash \langle a, b \rangle$ and $\overline{e} \vdash \langle a, b \rangle$, then $a = b$. This is evident if $a \in X$. Let $a = a_1 a_2$, $e \vdash \langle a, b \rangle$ and $\overline{e} \vdash \langle a, b \rangle$. Evidently, $b \notin X$; put $b = b_1 b_2$. Let

$\ulcorner \mu_1, \dots, \mu_n \urcorner$ be a minimal \overline{e} -proof of b from a . By Lemma 1 of [6], it has at most one leap. Suppose that it has exactly one leap i . It is sufficient to derive a contradiction in the case $\mu_i = \alpha \cdot \alpha \alpha$ & $\mu_{i+1} = \alpha \alpha \cdot \alpha$ for some $\alpha \in W_p$. We have $l(b_1) = 2 \cdot l(a_1)$. Hence, using Lemma 19, a minimal e -proof of b from a has exactly one leap, too, and for some $\beta, \gamma, \sigma \in W_p$

$$l(b_1) = l(\langle\langle\beta \cdot \beta \gamma\rangle \sigma\rangle) > 2 \cdot l(\beta) = 2 \cdot l(a_1), \text{ a contradiction.}$$

We have proved that $\ulcorner \mu_1, \dots, \mu_n \urcorner$ has no leaps. We get $\overline{e} \vdash \langle a_1, b_1 \rangle$ and $\overline{e} \vdash \langle a_2, b_2 \rangle$, so that $l(a_1) = l(b_1)$ and a minimal e -proof of b from a has no leaps, too. This implies $e \vdash \langle a_1, b_1 \rangle$ and

$e \vdash \langle a_2, b_2 \rangle$; by the induction assumption $a_1 = b_1$ and $a_2 = b_2$, so that $a = b$.

Lemma 21. Let x, y and z be three different variables; let e be any of the following eight equations:

$$\begin{aligned} & \langle (xx.x)y, xy \rangle ; \langle y(x.xx), yx \rangle ; \langle xx.x, xx \rangle ; \\ & \langle x.xx, xx \rangle ; \langle xx.y, x.yx \rangle ; \langle y.xx, xy.x \rangle ; \langle x.yx, xy.x \rangle ; \\ & \langle ((x.xy)x, x(x(yx.x))) \rangle . \end{aligned}$$

Then $Cn(e)$ is an upper semicomplement in \mathcal{L}_τ .

Proof follows from Lemmas 7, 14, 16, 18 and 20 and their duals.

§ 2. The infimum of the set of all upper semicomplements in \mathcal{L}_τ

Lemma 22. Let $x \in X$, $w \in W_\tau$ and $w \neq x$. Then $Cn(\langle x, w \rangle)$ is not an upper semicomplement in \mathcal{L}_τ .

Proof. Suppose on the contrary that there exists a non-trivial equation $\langle a, b \rangle$ such that $Cn(\langle x, w \rangle) \vee_\tau \vee_\tau Cn(\langle a, b \rangle) = \iota_\tau$. By Theorem 2 of [6], x is the only variable that is a subword of w ; i.e. $w \in T_\tau(x)$. As $w \neq x$, there exist $u, v \in T_\tau(x)$ such that $w = uv$. For every two elements κ, ρ of W_τ define $\kappa[\rho]$ by $\kappa[\rho] = \varphi(\kappa)$ where φ is the endomorphism of W_τ , assigning ρ to each variable. The equation $e = \langle u[w[a]], v[w[b]], w[u[a].v[b]] \rangle$ is evidently non-trivial and we have both $\langle x, w \rangle \vdash e$ and $\langle a, b \rangle \vdash e$, a contradiction.

Lemma 23. Let x, y and z be three different variables. If $a, b \in W_\tau$, then

$$\langle a, b \rangle \in \text{Cn}(\{ \langle xx.y, xy \rangle, \langle xy, yx \rangle, \langle xy.z, x.yz \rangle \})$$

if and only if $X \cap S(a) = X \cap S(b)$ and either $a = b$ or $a \notin X$ & $b \notin X$.

Proof is easy.

Theorem. The infimum in \mathcal{L}_Γ of all upper semicomplements in \mathcal{L}_Γ is just $\text{Cn}(\{ \langle xx.y, xy \rangle, \langle xy, yx \rangle, \langle xy.z, x.yz \rangle \})$ (where x, y and z are three different variables).

Proof. Denote the infimum by E . (E is a fully invariant congruence relation of W_Γ .) By Lemma 21 we have $\text{Cn}(\{ \langle xx.y, xy \rangle, \langle xy, yx \rangle, \langle xy.z, x.yz \rangle \}) \subseteq E$. The converse inclusion follows easily (some care is necessary) from Theorem 2 of [6] and Lemmas 22 and 23.

Denote by \mathcal{G} the variety of all groupoids. We reformulate the theorem two times:

Corollary 1. For every groupoid A , the following two conditions are equivalent:

- (i) $A \in \mathcal{U} \cap \mathcal{L}$ for every two proper subvarieties \mathcal{U}, \mathcal{L} of \mathcal{G} such that \mathcal{G} is the only variety containing both \mathcal{U} and \mathcal{L} ;
- (ii) A is a commutative semigroup satisfying $xx.y = xy$.

Corollary 2. Denote by E the set of all Γ -equations e such that $\text{Cn}(e)$ is an upper semicomplement in \mathcal{L}_Γ . Then

$$\text{Cn}(E) = \text{Cn}(\{ \langle xx.y, xy \rangle, \langle xy, yx \rangle, \langle xy.z, x.yz \rangle \})$$

Let L be an arbitrary lattice. An element $a \in L$

is called definable in L if there exists a formula φ of the first-order predicate calculus such that

- (i) φ contains only logical symbols, variables and the two function symbols \wedge and \vee ;
- (ii) φ has exactly one free variable;
- (iii) a satisfies φ in L and no other element of L satisfies φ .

Any lattice L has at most countably many definable elements. The set of all definable elements of L is a sublattice of L . Every definable element is a fix-point of any automorphism of L .

If L has the greatest and the smallest element, then they are evidently both definable in L . A less trivial example is the supremum of all atoms in a complete atomic lattice L . Hence, the variety of all semigroups satisfying $x y z w = x z y w$ (see [3]) is a definable element in the lattice of all semigroup varieties. Unfortunately, the supremum of the set of all atoms in \mathcal{L}_τ is just the greatest element of \mathcal{L}_τ (see [1] or [5]). However, the theorem gives us

Corollary 3. \mathcal{L}_τ has definable elements different from the greatest and the smallest elements.

$Cm(\langle\langle x x . y, x y \rangle, \langle x y, y x \rangle, \langle x y . z, x . y z \rangle\rangle)$ is a definable element.

The infimum of the set of all upper semicomplements is a definable element. It follows from Theorems 1 and 2 of [6] that if Δ is an arbitrary type containing at least one at least binary function symbol, then the infimum is a definable element in \mathcal{L}_Δ , different from the

extreme elements. It could be interesting to find this variety.

Problem. Find and describe other varieties of groupoids that are definable elements of \mathcal{L}_τ . Are the important varieties (the variety of semigroups, commutative groupoids, commutative semigroups, idempotent groupoids, semilattices,...) definable in \mathcal{L}_τ ? Denote by Δ the type consisting of one binary, one unary and one nullary function symbol. Is the variety of groups definable in \mathcal{L}_Δ ?

The problem stated in [6] remains open.

R e f e r e n c e s

- [1] A.D. BOL'BOT: O mnogoobrazijach Ω -algebr, Algebra i logika 9(1970),406-414.
- [2] R.A. DEAN and T. EVANS: A remark on varieties of lattices and semigroups, Proc.Amer.Math.Soc. 21(1969),394-396.
- [3] T. EVANS: The lattice of semigroup varieties, Semigroup Forum 2(1971),1-43.
- [4] J. JEŽEK: Principal dual ideals in lattices of primitive classes, Comment.Math.Univ.Carolinae 9(1968),533-545.
- [5] J. JEŽEK: On atoms in lattices of primitive classes, Comment.Math.Univ.Carolinae 11(1970),515-532.
- [6] J. JEŽEK: The existence of upper semicomplements in lattices of primitive classes, Comment.

Math.Univ.Carolinae 12(1971),519-532.

Matematicko-fyzikální fakulta

Karlova universita

Sokolovská 83, Praha 8

Československo

(Oblatum 18.3.1971)