

Roger Emanuel Kaufman

A nonlinear operator in potential theory

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 17 (1976), No. 4, 731--735

Persistent URL: <http://dml.cz/dmlcz/105733>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1976

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## A NONLINEAR OPERATOR IN POTENTIAL THEORY

R. KAUFMAN, Urbana

Abstract: A property of the first eigenvalue of the operator  $\Delta$  leads to the solvability of a nonlinear equation whose main part is a singular linear equation.

Key words: First eigenvalue, Hölder-continuity, fixed point.

AMS: 35J05, 47H15, 46E15      Ref. Ž.: 7.955.81

1. Let  $T$  be the nonlinear operator defined by  $T(u) = (\Delta + c^2)u + p(u)$ , where  $\Delta$  is the Laplacean in the unit disk  $D: x^2 + y^2 < 1$ ,  $p$  is continuous on  $(-\infty, \infty)$ , and  $p(u) = o(|u|)$ ,  $\lim_{|u| \rightarrow \infty} |p(u)| = +\infty$ , as  $|u| \rightarrow \infty$ . The domain of the operator  $\Delta$  is the space of all  $u$ , continuous on  $D$ , vanishing on  $\partial D$ , whose Laplacean (in the sense of distributions) belongs to  $L^2(D)$ ; Green's formula confirms that these functions  $u$  are Hölder-continuous. Moreover  $-c^2$  is the smallest eigenvalue of this operator, and  $\Delta$  is a closed, negative-definite operator.

Theorem. For each  $r > 2$  and each  $M$ , the set defined by the inequality  $\|T(u)\|_r \leq M$  is compact in the Banach space  $C^1(D^-)$ , and the range of  $T$  meets  $L^r(D)$  in a closed subset of  $L^r(D)$ .

The range contains  $L^r(D)$  if and only if

$p(+\infty) \cdot p(-\infty) < 0$ .

This theorem was suggested by a remarkable paper of Ambrosetti and Prodi [1] in which a similar use is made of the first eigenvalue of the operator  $\Delta$ .

2. The operator  $\Delta + a^2$  is singular precisely when  $a > 0$  is a zero of some Bessel function  $J_k$ , and the eigenfunction for  $c^2$  is  $f_0 = J_0(cr)$ ;  $f_0 > 0$  within  $D$ , and the normal derivative of  $f_0$  is negative on  $\partial D$ . (See [2, p. 373].) (Tables show that  $c \cong 2.40$  and the next zero is  $\cong 3.83$ .)

Green's formula, with zero boundary data,

$$f(z) = (2\pi)^{-1} \iint (\Delta f)(z') G(z, z') dx' dy'$$

shows that if  $\Delta f \in L^r$ ,  $2 < r < \infty$ , then  $f \in C^1(D^-)$ , and the first partial derivatives of  $f$  are Hölder-continuous in exponent  $1 - 2/r$ ; this is proved by means of Hölder's inequality and the potential-theoretic lemmas presented in [3, p. 198]. When  $1 < r < 2$  similar consideration yields Hölder-continuity of  $f$ .

To prove the first statement in the theorem we take a sequence  $u_n$ , in the domain of  $\Delta$ , such that  $\|T(u_n)\|_r \leq M$ . Supposing that  $m_n = \|u_n\|_\infty$  tends to infinity, we proceed to obtain a contradiction. We write  $u_n = a_n f_0 + v_n$ , where  $v_n$  is orthogonal to  $f_0$  in  $L^2(D)$ , and  $a_n$  is a real number. Since  $p(u) = o(|u|)$  as  $|u| \rightarrow +\infty$ , we see that  $(\Delta + c^2)v_n = o(m_n)$  uniformly, and therefore

in  $L^2$ . By the discreteness of the spectrum of  $\Delta$ , we see that  $(\Delta + c^2)v_n$  and  $\Delta v_n$  are of the same magnitude in  $L^2$ , whence  $v_n = o(m_n)$  uniformly. We now observe the identities

$$\Delta v_n = (\Delta + c^2)v_n - c^2v_n = T(u_n) - p(u_n) + c^2v_n,$$

and deduce that  $\Delta v_n = o(m_n)$  in  $L^r(D^2)$ . Therefore  $v_n = o(m_n)$  in the Banach space  $C^1(D^-)$ , whence  $v_n(z) = o(m_n)(1 - |z|)$ . We have also  $a_n \approx \frac{1}{2} m_n$ , so that  $u_n = a_n f_0 + v_n$  has no zeroes for large  $n$ , in view of the inequality  $f_0(z) \geq a(1 - |z|)$  valid for some  $a > 0$ .

If, for example,  $a_n > 0$  and  $p(+\infty) = +\infty$ , then  $p(u_n)$  tends everywhere to  $+\infty$ , while  $p(u_n) \geq -C$ . But  $(\Delta + c^2)u_n$  is orthogonal to  $f_0$ , so  $\iint p(u_n)f_0(z)dx dy = 0(1)$ , while  $f_0 > 0$ . This contradiction shows that  $m_n$  must remain bounded.

Now, by steps similar to the above, we find that  $a_n = 0(1)$ , so  $\|\Delta v_n\|_r = 0(1)$ , and then the functions  $u_n$  are bounded, with uniformly Hölder-continuous partial derivatives, in exponent  $1 - 2/r$ .

To prove the closure of the range of  $T$  in  $L^r$ , suppose  $\lim T(u_n) = g$  in  $L^r$ ; we can then select a subsequence  $u_j$ , converging to  $u_0$  in  $C^1(D^-)$ . Now  $\Delta u_j = T(u_j) - c^2u_j - p(u_j)$  and Green's formula shows that  $\Delta u_0 = g - c^2u_0 - p(u_0)$ , or  $T(u_0) = g$ .

3. Suppose now that  $p(u) \geq -C$  for all  $u$ ; then  $(T(u), f_0) \geq -C'$ , so that the range of  $T$  contains  $\lambda f_0$  only

when  $\lambda \leq \lambda_0$ .

To complete the proof, we suppose that  $p(+\infty) = +\infty$  and  $p(-\infty) = -\infty$  and prove that  $T(u) = g$  is solvable for every  $g$  in  $L^r$ ,  $r > 2$ . First we solve a perturbed equation  $T(u) + \varepsilon u = g$ , for small  $\varepsilon > 0$ . We write this in the form

$$(\Delta + c^2 + \varepsilon)u = g - p(u)$$

and observe that  $\Delta + c^2 + \varepsilon$  admits a bounded completely continuous inverse in  $L^2$ , for small  $\varepsilon$ . Let us define

$$A_\varepsilon(u) = (\Delta + c^2 + \varepsilon)^{-1} (g - p(u)).$$

$A_\varepsilon$  is continuous because  $g \in L^2$  and  $p(u) = o(|u|)$ , and compact, because  $(\Delta + c^2 + \varepsilon)^{-1}$  is compact. On the ball  $\|u\|_2 \leq N$ , we have  $\|A_\varepsilon(u)\|_2 = o(N)$  so that  $A_\varepsilon$  is a compact mapping of some ball into itself and admits a fixpoint by Schauder's theorem, i.e. a solution of the perturbed equation. To obtain a solution to the original equation, we prove that the solutions of the equations

$(\Delta + c^2 + \varepsilon)u + p(u) = g$  remain bounded as  $\varepsilon \rightarrow 0+$ . We write  $u = a_\varepsilon f_0 + v_\varepsilon$ , and suppose that  $\|u\|_\infty$  becomes unbounded. Then  $\|v_\varepsilon\|_2 = o(1)\|u\|_\infty$ , and we observe that

$$\Delta v_\varepsilon = g - p(u) - c^2 v_\varepsilon - \varepsilon a_\varepsilon f_0.$$

Thus  $\|v_\varepsilon\|_\infty = o(1)\|u\|_\infty$ , and finally  $v_\varepsilon = o(1)\|u\|_\infty$  in  $C^1(D^-)$ . Hence  $u$  maintains the same sign, and  $\varepsilon u + p(u)$  tends to  $+\infty$  (or  $-\infty$ ) remaining bounded below (above), so that the inner produce  $(T_\varepsilon u, f_0)$  becomes infinite. This completes the proof in the case  $p(+\infty) > 0$ ,  $p(-\infty) < 0$ . In

the event that  $p(+\infty) < 0$ ,  $p(-\infty) > 0$  we employ the perturbed operator  $T(u) - \varepsilon u$ .

4. An extension. The main theorem remains true in part for each  $r > 1$ , but to verify this we must consider the inverse of the operator  $\Delta + c^2$  on the appropriate subspace of  $L^r$ . It seems likely that an existence theorem remains true when  $r = 1$ , provided  $p'$  is bounded; the analysis would be difficult since the solutions  $u$  are unbounded.

#### R e f e r e n c e s

- [1] A. AMBROSETTI and G. PRODI: On the inversion of some differentiable mappings with singularities between Banach spaces, *Annali di Mat. Pura ed Appl.* (4) 93 (1972), 231-246.
- [2] P.W. BERG and J.L. MCGREGOR: *Elementary Partial Differential Equations*, San Francisco, 1966.
- [3] G. HELLMIG: *Partial Differential Equations*, New York, 1964.

University of Illinois  
at Urbana-Champaign  
Urbana, Illinois 61801  
U.S.A.

(Oblatum 3.8. 1976)