

Tomáš Kepka

Distributive groupoids and preradicals. II.

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 24 (1983), No. 2, 199--209

Persistent URL: <http://dml.cz/dmlcz/106219>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1983

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

DISTRIBUTIVE GROUPOIDS AND PRERADICALS II

T.KEPKA

Abstract: One-sided ideals and the corresponding preradicals of distributive groupoids are studied.

Key words: Groupoid, preradical.

Classification: 20L10

---

This note is an immediate continuation of [2]. The theory of preradicals developed in [2] is applied to some special cases. Two preradicals derived from left and right ideals are defined and their rôle in the structure theory of distributive groupoids is studied.

9. Ideals. Let  $A$  denote the class of distributive idempotent groupoids.

9.1. Lemma. Let  $G \in A$ .

- (i) If  $I$  is an ideal of  $G$  and  $K$  a left (right) ideal of  $I$  then  $K$  is a left (right) ideal of  $G$ .
- (ii) If  $I$  is a left (right) ideal of  $G$  and  $K$  an ideal of  $I$  then  $K$  is a left (right) ideal of  $G$ .
- (iii) If  $I$  is an ideal of  $G$  and  $K$  an ideal of  $I$  then  $\hat{K}$  is an ideal of  $G$ .

Proof. (i) We have  $ab \in I$  and  $ab = ab.ab = (ab.a)(ab.b)$  for all  $a \in G$  and  $b \in K$ . Since  $I$  is an ideal of  $G$  and  $K$  a left ideal of  $I$ ,  $ab.a \in I$  and  $ab.b \in K$ . Consequently,  $ab \in K$ .

(ii) We have  $ab \in I$  and  $ab = ab.ab = (a.ab)(b.ab)$  for all  $a \in G$  and  $b \in K$ . Since  $I$  is a left ideal of  $G$  and  $K$  an ideal of  $I$ ,  $a.ab \in I$  and  $b.ab \in K$ . Consequently,  $ab \in K$ .

(iii) Use (i) or (ii).

9.2. Lemma. Let  $I, K$  be left ideals of a groupoid  $G \in A$ . Then  $IK$  is a left ideal of  $G$  and  $IK \subseteq K$ .

9.3. Lemma. Let  $I$  be a left and  $K$  a right ideal of a groupoid  $G \in A$ . Then  $KI = I \cap K$  is a left (right) ideal of  $K(I)$ . Moreover,  $KI$  is an ideal of  $G$ , provided  $IK \subseteq KI$ .

9.4. Lemma. Let  $I$  and  $K$  be ideals of a groupoid  $G \in A$ . Then  $IK = I \cap K = KI$  is an ideal of  $G$ .

9.5. Lemma. Let  $I$  be an ideal of a groupoid  $G \in A$  and let  $a, b, c \in G$ . Then  $ab \in I$  iff  $ba \in I$  and  $a.bc \in I$  iff  $ab.e \in I$ .

Proof. We have  $ba = ba.ba = (ba.b)(ba.a) = (b.ab)(ba.a)$  and  $ab.e = ac.bc = (a.bc)(c.bc)$ ,  $a.bc = ab.ac = (ab.a)(ab.c)$ .

For  $G \in A$ , define a relation  $w(G)$  by  $(a, b) \in w(G)$  iff the elements  $a$  and  $b$  generate the same ideal of  $G$ .

9.6. Lemma. Let  $G \in A$ . Then:

(i)  $(a, b) \in w(G)$  iff  $a = f(b)$  and  $b = g(a)$  for some  $f, g \in \text{Mul}(G)$ .

(ii)  $w(G)$  is a congruence of  $G$ ,  $H = G/w(G)$  is a semilattice and  $w(H) = \text{id}_H$ .

(iii) Every block of  $w(G)$  is ideal-free.

(iv) If  $I$  is an ideal of  $G$  then  $w(I) = w(G) \cap (I \times I)$ .

Proof. (i) This is clear.

(ii) Apply (i) and 9.5.

(iii) Let  $H$  be a block of  $w(G)$  and  $a, b \in H$ . There are positive integers  $n, m, S_1, \dots, S_n, T_1, \dots, T_m \in \{L, R\}$  and  $a_1, \dots, a_n, b_1, \dots, b_m \in G$  such that  $a = S_{1, a_1} \dots S_{n, a_n}(b)$  and  $b = T_{1, b_1} \dots T_{m, b_m}(a)$ . Then  $a = aa = S_{1, aa_1} \dots S_{n, aa_n}(a)$  and  $b = T_{1, bb_1} \dots T_{m, bb_m}(ba)$ . From this, it is easy to see that  $aa_j, bb_j \in H$ , and so  $(a, b) \in w(H)$ . Thus  $w(H) = H \times H$  and  $H$  is ideal-free.

9.7. Corollary.  $w$  is an idempotent radical.

9.8. Lemma. Let  $I$  be an ideal of a groupoid  $G \in \mathcal{A}$ . Then  $G$  is isomorphic to a subgroupoid of the product of  $G/I$  and a set of copies of  $I$ .

Proof. For every  $a \in I$ , both  $L_a$  and  $R_a$  are homomorphisms of  $G$  into  $I$  and  $r \cap \ker(L_a) \cap \ker(R_a) = \text{id}$ , where  $r = (I \times I) \cup \text{id}$ .

9.9. Corollary. Let  $I$  be an ideal of a groupoid  $G \in \mathcal{A}$ . Then the groupoids  $G$  and  $I \times G/I$  generate the same groupoid variety.

9.10. Proposition. Let  $G \in \mathcal{A}$ . Then  $w(G)$  is just the least congruence of  $G$  such that the corresponding factorgroupoid is a semilattice.

Proof. Denote by  $r$  the congruence. By 9.6(ii),  $r \subseteq w(G)$ . However, if  $H$  is a block of  $w(G)$  then  $r|_H = H \times H$  as it follows from 9.6(iii). Hence  $w(G) \subseteq r$ .

9.11. Corollary.  $w = \pi_S$ ,  $S$  being the class of semilattices (see 6.2).

**10. Left and right ideals.** Let  $A$  denote the class of distributive idempotent groupoids. A groupoid  $G$  is said to be left (right) permutable if it satisfies the identity  $x.yz = y.xz$  ( $xy.z = xz.y$ ).

**10.1. Lemma.** Let  $G \in A$  be left (right) permutable. Then  $G$  is medial.

**Proof.** For  $a, b, c, d \in G$ ,  $ab.cd = c(ab.d) = c(ad.bd) = c(b(ad.d)) = b(c(ad.d)) = b(ad.cd) = b(ac.d) = ac.bd$ .

**10.2. Lemma.** Let  $G \in A$  be both left and right permutable. Then  $G$  is a semilattice.

**Proof.** For  $a, b, c \in G$ ,  $a.bc = ab.ac = (a.ac)b = (ab)(ac.b) = (ac)(ab.b) = (a(ab.b))c = (ab.ab)c = ab.c$  and  $ab = ab.a = a.ba = ba$ .

**10.3. Lemma.** Let  $I$  be a left ideal of a groupoid  $G \in A$  and let  $a, b, c \in G$ . Then  $a.bc \in I$  iff  $b.ac \in I$ .

**Proof.**  $a.bc = ab.ac = (a.ac)(b.ac)$ .

For  $G \in A$ , define a relation  $u(G)$  (resp.  $v(G)$ ) by  $(a, b) \in u(G)$  (resp.  $v(G)$ ) iff the elements  $a$  and  $b$  generate the same left (resp. right) ideal of  $G$ .

**10.4. Lemma.** Let  $G \in A$ . Then:

- (i)  $(a, b) \in u(G)$  iff  $a = f(b)$  and  $b = g(a)$  for some  $f, g \in \text{Mull}(G)$ .
- (ii)  $u(G)$  is a congruence of  $G$ ,  $H = G/u(G)$  is left permutable and  $u(H) = \text{id}_H$ .
- (iii) If  $K$  is a block of  $u(G)$  then  $K/u(K)$  is a semigroup of right zeros.

**Proof.** (i) This is clear.

(ii)  $u(G)$  is a congruence and  $H$  is left permutable by (i) and

10.3. Denote by  $f$  the natural projection of  $G$  onto  $H$ . Let

$a, b \in G$  be such that  $(f(a), f(b)) \in u(H)$ . Then there are positive integers  $n, m$  and  $a_1, \dots, a_n, b_1, \dots, b_m \in G$  with  $(a_1(a_2(\dots(a_n a)))) , b) \in u(G)$  and  $(b_1(b_2(\dots(b_m b)))) , a) \in u(G)$ . From this, it is easy to see that  $(a, b) \in u(G)$ .

(iii) Let  $a, b \in K$ . There are positive integers  $n, m$  and  $a_1, \dots, a_n, b_1, \dots, b_m \in G$  with  $a = b_1(\dots(b_m b))$  and  $b = a_1(\dots(a_n a))$ . Then  $a = b_1(\dots(b_m(a_1(\dots(a_n a)))))$  and  $a = b_1(\dots(b_{i-1}((b_i b_{i+1})(\dots((b_i b_m)((b_i a_1)(\dots(b_i a_n \cdot b_i a)))))))$  for every  $1 \leq i \leq m$  and we see that  $b_i a \in K$ . On the other hand,  $a = (b_1(\dots(b_m b)))a = (b_1 a)(\dots(b_m a \cdot ba))$ , and therefore  $(a, ba) \in u(K)$ .

10.5. Corollary. Both  $u$  and  $v$  are radicals.

10.6. Corollary. Both  $\bar{u}$  and  $\bar{v}$  are idempotent radicals.

10.7. Lemma. Let  $G \in A, n \geq 2$  and  $a_1, \dots, a_n \in G$ . Then there are  $b_1, \dots, b_{n-2} \in G$  such that  $((a_1 a_2) \dots) a_n = b_1(\dots(b_{n-2} \cdot R_{a_n}^{n-1}(a_1)))$ .

Proof. By induction on  $n$ . For  $n = 2$ , there is nothing to prove. For  $n \geq 3$ ,  $((a_1 a_2) \dots) a_n = b_1(\dots(b_{n-3} c))$ ,  $c = R_{a_n}^{n-2}(a_1 a_2)$ . But,  $c = R_{a_n}^{n-2}(a_1) \cdot R_{a_n}^{n-2}(a_2) = R_{a_n}^{n-2}(a_1)(R_{a_n}^{n-3}(a_2) \cdot a_n) = (R_{a_n}^{n-2}(a_1) \cdot R_{a_n}^{n-3}(a_2)) \cdot R_{a_n}^{n-1}(a_1)$ .

10.7. Lemma. Let  $G \in A, n \geq 1, a, b, a_1, \dots, a_n \in G$ ,  $a = ((ba_1) \dots) a_n$  and let  $H$  be the block of  $u(G)$  containing  $a$ . Then there are  $m \geq 1$  and  $b_1, \dots, b_m \in H$  such that  $a = ((bb_1) \dots) b_m$ .

Proof. Let  $1 \leq i \leq n$ . We have  $a = c(((a_1 a_{i+1}) \dots) a_n)$ ,  $c = (((((ba_1) \dots) a_{i-1}) a_{i+1}) \dots) a_n)$ . From this,  $a = aa = (ca)((((a_1 a_{i+1}) \dots) a_n) a)$ . By 10.6, there are  $c_1, \dots, c_{n-1} \in G$  such that  $a = (ca)(c_1(\dots(c_{n-1} \cdot R^{n-1+1}(a_1))))$ . Obviously,

$R_a^{n-1+1}(a_1) \in H$ . However, then  $d_1 = R_a^n(a_1) \in H$  and  $a = ((R_a^n(b)d_1) \dots) d_n$ .

10.8. Proposition. (i)  $u.v = v.u = u \cap v$ .

(ii)  $\widehat{a\ell} \subseteq \bar{u}$  and  $\widehat{a\bar{r}} \subseteq \bar{v}$  (see 7.2).

(iii)  $m_M \subseteq u \cap v$  (see 7.3).

(iv)  $u+v \subseteq w$ ,  $u:v \subseteq w$  and  $v:u \subseteq w$ .

Proof. (i) By 10.7 and its dual,  $u \cap v \subseteq v.u$ ,  $u \cap v \subseteq u.v$  and the result follows from 4.1(i).

(ii) The inclusion  $a\ell \subseteq u$  is clear directly from the definitions. Since  $u$  is a radical and  $\widehat{a\ell}$  is idempotent,  $\widehat{a\ell} \subseteq \bar{u}$ .

(iii) This follows from 10.1, 10.4(ii) and its dual.

(iv) This is clear.

10.9. Corollary.  $u^n.v^m = v^m.u^n = u^n \cap v^m$  for all positive integers  $n, m$ .

10.10. Corollary.  $u.\bar{v} \subseteq \bar{v}.u$  and  $v.\bar{u} \subseteq \bar{u}.v$ .

10.11. Lemma. Let  $G \subset A$  be left permutable and  $(a, b) \in u(G)$ . Then  $ab = b$  and  $ba = a$ .

10.12. Proposition. Let  $G \subset A$  be left permutable. Then:

(i)  $u(G) \subseteq \text{ar}(G) \subseteq v(G) = w(G) = \bar{v}(G)$ .

(ii) Every block of  $u(G)$  is a semigroup of right zeros.

(iii)  $\bar{u}(G) = u^2(G) = \text{id}_G$  and  $a\ell(G) = \text{id}_G$ .

Proof. (i) By 10.11 and 10.8(ii), (iv),  $u(G) \subseteq \text{ar}(G) \subseteq v(G) \subseteq w(G)$ . Denote by  $f$  the natural projection of  $G$  onto  $H = G/v(G)$ . By 10.2 and the dual of 10.4(ii),  $H$  is a semilattice, and hence  $w(H) = \text{id}_H$ . If  $(a, b) \in w(G)$  then  $(f(a), f(b)) \in w(H)$ ,  $f(a) = f(b)$  and  $(a, b) \in v(G)$ . Thus  $w(G) = v(G)$ .

(ii) This is clear from 10.11.

(iii) Use (ii) and 10.8(ii).

10.13. Proposition.  $\bar{v}:u = \bar{u}:v = u:v = v:u = w$ .

Proof. Let  $G \in A$  and let  $f$  denote the natural projection of  $G$  onto  $H = G/u(G)$ . We have  $w(H) = \bar{v}(H)$  by 10.12(1). Consequently,  $(w:u)(G) = (\bar{v}:u)(G)$ . However,  $w(G) \subseteq (w:u)(G)$ , we have proved  $w \subseteq \bar{v}:u$ , and so  $w = \bar{v}:u$ . Similarly,  $w = \bar{u}:v$ .

10.14. Proposition. Let  $G \in A$  be left permutable. Then  $\widehat{\text{ar}}(G) * w(G) = v(G)$ .

Proof. Put  $H = G/\widehat{\text{ar}}(G)$ . Then  $\text{ar}(H) = \text{id}_H$ . However,  $(a.ab)(ab) = a(ab.b) = ab.ab = ab$  and  $(ab)(a.ab) = a(ab.ab) = a.ab$  for all  $a, b \in H$ . Hence  $ab = a.ab$ . Further,  $ba.ab = a(ba.b) = a(b.ab) = a.ab = ab$  and  $ab.ba = ba$ . From this,  $ab = ba$  and  $H$  is a semilattice. The rest is clear.

10.15. Corollary. Let  $G \in A$  be left permutable and ideal-free. Then  $G$  is  $\widehat{\text{ar}}$ -torsion and right-ideal-free.

10.16. Lemma. Let  $G$  be a groupoid containing a subgroupoid  $H$  such that  $H$  is a semigroup of right zeros,  $G = H \cup \{0\}$ ,  $0 \notin H$ ,  $0.H \subseteq H$  and  $a0 = 0$  for every  $a \in G$ . Then  $G \in A$  is left permutable and  $u(G) \subseteq H \times H \subseteq p(G)$ .

Proof. Obviously,  $G$  is idempotent. Now, we show that  $G$  is medial. For, let  $a, b, c, d \in G$ . If  $a, b, c, d \in H$ , then  $ab.cd = d = ac.bd$ . If  $d = 0$  then  $ab.cd = d = ac.bd$ . If  $c = 0$  and  $a, b, d \in H$  then  $ab.cd = cd = c.bd = ac.bd$ . If  $a = 0$  and  $b, c, d \in H$  then  $ab.cd = d = ac.bd$ . If  $a = 0 = c$  and  $b, d \in H$  then  $ab.cd = cd = c.bd = ac.bd$ . If  $a = b = c = 0$  and  $d \in H$  then  $ab.cd = ac.bd$ . Finally, we show that  $G$  is left permutable. For, let  $a, b, c \in G$ . If  $a, b, c \in H$  then  $a.bc = b.ac$ . If  $c = 0$  then  $a.bc = c = b.ac$ . If  $a = 0$  and  $b, c \in H$  then  $a.bc = ac = b.ac$ . If  $a = 0 = b$  and  $c \in H$  then  $a.bc = b.ac$ .



10.17. Example. Consider the following three-element groupoid  $G = \{a, b, c\}$ ;  $aa = ba = ca = a$ ,  $ab = bc = cc = c$ ,  $ac = bb = cb = b$ . By 10.16,  $G \in \mathcal{A}$  and  $G$  is left permutable. Moreover, it is easy to see that  $p(G) = ar(G) = u(G) = id_G \cup \{(b,c), (c,b)\}$ . Hence  $u(G) \neq id_G$ .

10.18. Lemma. Let  $n$  be a non-negative integer and let  $G \in \mathcal{A}$  be  $u^n$ -torsionfree. Then  $v(G) = w(G) = \bar{v}(G)$ .

Proof. We show by induction on  $n$  that  $\bar{v}(G) = w(G)$ . With respect to 10.12(i), we can assume that  $n \geq 2$ . Denote by  $f$  the natural projection of  $G$  onto  $H = G/u^{n-1}(G)$  and by  $g$  that of  $G$  onto  $K = G/\bar{v}(G)$ . According to 10.4(iii), every block of  $u^{n-1}(G)$  is a semigroup of right zeros, and hence  $u^{n-1}(G) \subseteq \bar{v}(G)$ . Using this, we see that there is a projective homomorphism  $h$  of  $H$  onto  $K$  such that  $g = hf$ . Now, let  $(a,b) \in w(G)$ . Then, by the induction hypothesis,  $(f(a), f(b)) \in \bar{v}(H)$ , and so  $(g(a), g(b)) \in \bar{v}(K)$ . Consequently,  $(a,b) \in (\bar{v}:\bar{v})(G) = \bar{v}(G)$ .

10.19. Proposition.  $\bar{v}:u^n = w = \bar{u}:v^n$  for every positive integer  $n$ .

Proof. Let  $G \in \mathcal{A}$  and let  $f$  denote the natural projection of  $G$  onto  $H = G/u^n(G)$ . Let  $(a,b) \in w(G)$ . Then  $(f(a), f(b)) \in w(H) = \bar{v}(H)$  by 10.18, and hence  $(a,b) \in (\bar{v}:u^n)(G)$ .

10.20. Corollary.  $v^n:u^m = w = u^m:v^n$  for all positive integers  $n, m$ .

11. An application. A congruence  $r$  of a groupoid  $G$  is said to be  $e$ -invariant (resp.  $a$ -invariant) if it is invariant with respect to all endomorphisms (resp. automorphisms) of  $G$ .

The groupoid  $G$  is said to be  $e$ -simple (resp.  $a$ -simple) if it is non-trivial and  $\text{id}_G, G \times G$  are the only  $e$ -invariant (resp.  $a$ -invariant) congruences of  $G$ .

11.1. Proposition. Let  $A$  be a non-empty abstract class of groupoids and  $r$  a semipreradical (resp. a preradical). If  $G \in A$  is  $a$ -simple (resp.  $e$ -simple) then either  $r(G) = \text{id}_G$  or  $r(G) = G \times G$ .

11.2. Proposition. Every  $e$ -simple distributive groupoid is either idempotent or a semigroup with zero multiplication. Conversely, every non-trivial semigroup with zero multiplication is an  $e$ -simple distributive groupoid.

Proof. Let  $G$  be an  $e$ -simple distributive groupoid. The set  $I$  of all idempotents of  $G$  is an ideal and it is easy to see that  $r = (I \times I) \cup \text{id}_G$  is an  $e$ -invariant congruence of  $G$ . If  $r = G \times G$  then  $I = G$  and  $G$  is idempotent.

Suppose that  $r \neq G \times G$ . Then  $r = \text{id}_G$ ,  $I$  contains only one element and  $G$  is a semigroup nilpotent of class at most 3. Put  $K = GG$  and  $s = (K \times K) \cup \text{id}_G$ . Again,  $K$  is an ideal of  $G$  and  $s$  is an  $e$ -invariant congruence. If  $s = G \times G$  then  $G = GG$  and  $G$  is idempotent, a contradiction. Thus  $s = \text{id}_G$ ,  $K$  contains just one element and  $G$  is a semigroup with zero multiplication.

11.3. Corollary. Every  $a$ -simple distributive groupoid is either idempotent or a two-element semigroup with zero multiplication.

11.4. Proposition. Let  $G$  be an  $e$ -simple distributive idempotent groupoid. Then exactly one of the following four cases takes place:

(1)  $u(G) = G \times G = v(G)$ ,  $G$  is both left and right-ideal free

and  $G$  is cancellative.

(ii)  $u(G) = id_G = v(G)$  and  $G$  is a semilattice.

(iii)  $u(G) = id_G$ ,  $v(G) = G \times G$ ,  $G$  is right-ideal-free and  $G$  is left permutable.

(iv)  $v(G) = id_G$ ,  $u(G) = G \times G$ ,  $G$  is left-ideal-free and  $G$  is right permutable.

*Proof.* By 11.1,  $u(G), v(G) \in \{id_G, G \times G\}$ . If  $u(G) = id_G = v(G)$  then  $G$  is a semilattice by 10.2, 10.4(ii) and its dual. If  $u(G) = id_G$  and  $v(G) = G \times G$  then  $G$  is left permutable by 10.4(ii) and  $G$  is clearly right-ideal-free. Suppose that  $u(G) = G \times G = v(G)$ . Then  $G$  is both left and right-ideal-free and  $G$  is regular (see [1]). However, the regularity of  $G$  implies that  $p(G)$  is an  $e$ -invariant congruence of  $G$ . If  $p(G) = G \times G$  then  $G$  is a semigroup of right zeros, and, since it is left-ideal-free, it is trivial, a contradiction. We have proved that  $p(G) = id_G$ , and hence  $G$  is right cancellative. Similarly,  $G$  is left cancellative.

11.5. Lemma. Every non-trivial semigroup of right zeros is an  $a$ -simple distributive idempotent groupoid.

11.6. Lemma. (i) If  $G$  is a finite  $a$ -simple semilattice then every non-zero element of  $G$  is an atom.

(ii) The three-element chain is an  $e$ -simple semilattice.

11.7. Proposition. Let  $G$  be a finitely generated  $e$ -simple distributive groupoid. Then exactly one of the following five cases takes place:

(i)  $G$  is a finite semigroup with zero multiplication.

(ii)  $G$  is a finite semigroup of left zeros.

(iii)  $G$  is a finite semigroup of right zeros.

(iv)  $G$  is a finite semilattice.

(v)  $G$  is a finite quasigroup.

Proof. With respect to 11.2, we can assume that  $G$  is idempotent. Denote by  $A$ ,  $B$  and  $C$  the classes of left-zero semigroups, right-zero semigroups and semilattices, resp. Then  $m_A(G)$ ,  $m_B(G)$  and  $m_C(G)$  are  $e$ -invariant congruences of  $G$  and we can assume that  $m_A(G) = m_B(G) = m_C(G) = G \times G$ . Since  $G$  is finitely generated,  $G$  possesses a non-trivial simple factor-groupoid  $Q$  and we see that  $Q$  is a finite quasigroup. Denote by  $V$  the variety generated by  $Q$ . Then  $V$  is locally finite and  $G \in V$ . In particular,  $G$  is a finite quasigroup.

#### R e f e r e n c e s

- [1] J. JEŽEK, T. KEPKA, P. NĚMEC: Distributive groupoids, Rozpravy ČSAV 91/3, 1981.
- [2] T. KEPKA: Distributive groupoids and preradicals I, Comment. Math. Univ. Carolinae 24(1983),

Matematicko-fyzikální fakulta, Univerzita Karlova, Sokolovská 83, 18600 Praha 8, Czechoslovakia

(Oblatum 13.10. 1982)