

Heinz-Jürgen Voss

Note on a paper of McMorris and Shier

Commentationes Mathematicae Universitatis Carolinae, Vol. 26 (1985), No. 2, 319--322

Persistent URL: <http://dml.cz/dmlcz/106372>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1985

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

NOTE ON A PAPER OF McMORRIS AND SHIER
Heinz-Jürgen VOSS

Abstract: F. R. McMORRIS and D. R. SHIER [3] proved: A graph G is split iff G can be represented as an intersection graph of a set of distinct subtrees of $K_{1,n}$. They give a method for constructing this intersection graph. Here an improved construction with minimum n is described.

Keywords: Chordal graphs, split graphs, intersection graphs.

Classification: 05C75

Only finite connected simple graphs are to be considered.

For the terminology see [1] and [3].

A graph G is said to be represented on a tree T if and only if G is isomorphic to the intersection graph of a set of distinct subtrees of T .

A graph $G = (V, E)$ is split if and only if there is a partition of the vertex set as $V = I \cup K$, where I is an independent set and K is complete. Furthermore, the partition $V = I \cup K$ can be chosen so that K is a maximum clique [2]. Henceforth we shall assume that K has been chosen in this manner.

Investigating chordal graphs F. R. McMORRIS and D. R. SHIER proved [3]:

Theorem 1. A graph $G = (V, E)$ is split if and only if G can be represented on $K_{1,n}$ for some n .

In their proof F. R. MORRIS and D. R. SHIER [3] constructed a representation on $K_{1,n}$ for a given split graph. They claim that their method of construction provides a representation of G on $K_{1,n}$ using the smallest possible n . This is not true and I shall give the corrected method.

Before describing it we define: if A, B are sets then $P(A) = \{M/M \subseteq A\}$, where $\emptyset \in P(A)$ and $B \cup P(A) = \{X/X = B \cup M, M \in P(A)\}$. The subgraph of G induced by H is denoted by $G[H]$. Let $N(x)$ denote the set of all neighbours of the vertex $x \in V$. If $I \subseteq V$ then $N_I(x) = N(x) \cap I$. For real q let $\lceil q \rceil$ denote the smallest integer $\geq q$.

C o n s t r u c t i o n. Suppose $G = (V, E)$ is split, where $V = I \cup K$ and $I = \{x_1, \dots, x_r\}$. First, label the end vertices (of degree 1) in T by the integers $1, \dots, r$ and the vertex of degree r by 0 . Define the subtree $T(x_1)$, corresponding to vertex x_1 , by $T(x_1) = \{1\}$, for all $1 \leq i \leq r$. Next, let L , initially empty, denote a collection of subsets and A , also initially empty, a set of additional vertices of T . For each $y \in K$, we consult L to see if all members of $N_I(y) \cup P(A)$ are in the list L . If not, choose one of the members M not in L , define subtree $T(y) = T[M \cup \{0\}]$ and add M to the list L . If all members of $N_I(y) \cup P(A)$ are in the list L we add a new end vertex α to the current T (joining it to vertex 0) and define $T(y) = T[N_I(y) \cup \{0, \alpha\}]$. We add α to the list A and $N_I(y) \cup \{\alpha\}$ to the list L . This procedure is repeated for all vertices $y \in K$. Upon completion, the process yields a $K_{1,n}$ and a set of distinct subtrees that represent G . \square

Applying my construction to K_4 and K_6^- (obtained from K_6

by omitting an edge) I have in both cases a $K_{1,n}$ with an n which is smaller than the one of F. R. McMORRIS and D. R. SHIER.

Theorem 2. For every split graph $G = (V, E)$, $V = I \cup K$, the Construction provides a representation of G on $K_{1,n}$ with minimal n . If m denotes the maximum number of vertices of K having the same neighbourhood $N_I(y)$ in I then the minimal $n = |I| + \lceil \log_2 m \rceil$.

For the simple proof we need the following obvious lemma.

Lemma 3. Let S_1, \dots, S_r and T_1, \dots, T_s be the subtrees of $K_{1,n}$ containing precisely 1 vertex or ≥ 2 vertices, respectively. Then

- 1) in the intersection graph G^* the subtrees S_1, \dots, S_r form a K_r and the subgraph of G^* induced by T_1, \dots, T_s is a K_s ;
- ii) if S_i ($1 \leq i \leq r$) consists of the "central" vertex (of degree n) of $K_{1,n}$ then S_i is joined to all T_j ($1 \leq j \leq s$) by edges; i. e. the subgraph of G^* induced by S_i, T_1, \dots, T_s is a K_{s+1} .

Proof of Theorem 2. Let $G = (V, E)$ be split with partition $V = I \cup K$ such that K is of maximum possible order. Let $I = \{x_1, x_2, \dots, x_r\}$ and $K = \{y_1, y_2, \dots, y_s\}$. Let $X_1, \dots, X_r, Y_1, \dots, Y_s$ be a representation of G on $K_{1,n}$ such that $x_i \leftrightarrow X_i$ and $y_j \leftrightarrow Y_j$. By Lemma 3 and the maximality of K each subtree X_i consists of an end vertex of $K_{1,n}$. Let the vertices of $K_{1,n}$ be denoted by $0, 1, \dots, n$ so that 0 is the "central" vertex of $K_{1,n}$ and $X_i = \{i\}$ for $1 \leq i \leq r$.

The subtree Y_j contains the vertex i of $K_{1,n}$ ($1 \leq i \leq r$) iff $(x_i, y_j) \in E$. Thus the subtree $Y_j[\{0, 1, \dots, r\}]$ of Y_j

induced by $\{0, 1, \dots, r\}$ is uniquely determined.

Let m be an integer defined as follows: there are m vertices $y^1, \dots, y^m \in K$ having the same neighbourhood in I and there are no $m+1$ such vertices in K . Let Y^1, \dots, Y^m denote the corresponding subtrees. Then $Y^1[\{0, \dots, r\}] = \dots$
 $\dots = Y^m[\{0, \dots, r\}]$.

Since Y^1, \dots, Y^m are pairwise distinct subtrees they contain some of the vertices $r+1, \dots, n$. With these vertices a set $N_I(y_1) \cup \{r+1, \dots, n\}$ of 2^{n-r} subtrees of $K_{1,n}$ with fixed $N_I(y_1)$ can be formed. Consequently, the minimal n has to be chosen $n = r + \lceil \log_2 m \rceil$. \square

In a further paper I shall investigate intersection graphs of a set S of distinct subtrees of a tree T , where no element of S is contained in an other element of S .

References

- [1] J. A. BONDY and U. S. R. MURTY: Graph Theory with Applications, American Elsevier, New York (1977)
- [2] M. C. GOLUMBIC: Algorithmic Graph Theory and Perfect Graphs, Academic Press, New York (1980)
- [3] F. R. McMORRIS and D. R. SHIER: Representing chordal graphs on $K_{1,n}$. Comm. Math. Univ. Carolinae 24, 3 (1983); 489 - 494

Sektion Mathematik
Pädagogische Hochschule "Karl Friedrich Wilhelm Wacker" Dresden
DDR - 8060 Dresden
Wigardstr. 17
German Democratic Republic

(Oblatum 29.10. 1984)