

Abdelkader Stouti

A fuzzy version of Tarski's fixpoint theorem

*Archivum Mathematicum*, Vol. 40 (2004), No. 3, 273--279

Persistent URL: <http://dml.cz/dmlcz/116923>

## Terms of use:

© Masaryk University, 2004

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## A FUZZY VERSION OF TARSKI'S FIXPOINT THEOREM

ABDELKADER STOUTI

ABSTRACT. A fuzzy version of Tarski's fixpoint Theorem for fuzzy monotone maps on nonempty fuzzy complete lattice is given.

### 1. INTRODUCTION

Let  $X$  be a nonempty set. A fuzzy set in  $X$  is a function  $\mu : X \rightarrow [0, 1]$ . Fuzzy set theory is a powerful tool for modelling uncertainty and for processing vague or subjective information in mathematical models. In [9], Zadeh introduced the notion of fuzzy order and similarity. Recently, several authors studied the existence of fixed point in fuzzy setting, Heilpern [7], Hadzic [6], Fang [5] and Beg [1, 2, 3]. In fuzzy ordered sets, I. Beg [1] proved the existence of maximal fixed point of fuzzy monotone maps. The aim of this note is to give the following fuzzy version of Tarski's fixpoint Theorem [8]: suppose that  $(X, r)$  is a nonempty  $r$ -fuzzy complete lattice and  $f : X \rightarrow X$  is a  $r$ -fuzzy monotone map. Then the set  $\text{Fix}(f)$  of all fixed points of  $f$  is a nonempty  $r$ -fuzzy complete lattice.

### 2. PRELIMINARIES

In this note we shall use the following definition of order due to Claude Ponsard (see [4]).

**Definition 2.1.** Let  $X$  be a crisp set. A fuzzy order relation on  $X$  is a fuzzy subset  $R$  of  $X \times X$  satisfying the following three properties

- (i) for all  $x \in X$ ,  $r(x, x) \in [0, 1]$  (f-reflexivity);
- (ii) for all  $x, y \in X$ ,  $r(x, y) + r(y, x) > 1$  implies  $x = y$  (f-antisymmetry);
- (iii) for all  $(x, y, z) \in X^3$ ,  $[r(x, y) \geq r(y, x) \text{ and } r(y, z) \geq r(z, y)]$  implies  $r(x, z) \geq r(z, x)$  (f-transitivity).

A nonempty set  $X$  with fuzzy order  $r$  defined on it, is called  $r$ -fuzzy ordered set. We denote it by  $(X, r)$ . A  $r$ -fuzzy order is said to be total if for all  $x \neq y$  we

---

2000 *Mathematics Subject Classification*: 03E72, 06B23, 47H10.

*Key words and phrases*: fuzzy set, fuzzy order relation, complete lattices, monotone map, fixed point.

Received August 2, 2002.

have either  $r(x, y) > r(y, x)$  or  $r(y, x) > r(x, y)$ . A  $r$ -fuzzy ordered set on which the  $r$ -fuzzy order is total is called  $r$ -fuzzy chain.

Let  $A$  be a nonempty subset of  $X$ . We say that  $x \in X$  is a  $r$ -upper bound of  $A$  if  $r(y, x) \geq r(x, y)$  for all  $y \in A$ . A  $r$ -upper bound  $x$  of  $A$  with  $x \in A$  is called a greatest element of  $A$ . An  $x \in A$  is called a maximal element of  $A$  if there is no  $y \neq x$  in  $A$  for which  $r(x, y) \geq r(y, x)$ . Similarly, we can define  $r$ -lower bound, minimal and least element of  $A$ . As usual,  $\sup_r(A)$ = the unique least element of  $r$ -upper bound of  $A$  (if it exists),

- $\max_r(A)$ = the unique greatest element of  $A$  (if it exists),
- $\inf_r(A)$ = the unique greatest element of  $r$ -lower bound of  $A$  (if it exists),
- $\min_r(A)$ = the unique least element of  $A$  (if it exists).

**Definition 2.2.** Let  $(X, r)$  be a nonempty  $r$ -fuzzy ordered set. The inverse fuzzy relation  $s$  of  $r$  is defined by  $s(x, y) = r(y, x)$ , for all  $x, y \in X$ .

**Definition 2.3.** Let  $(X, r)$  be a nonempty  $r$ -fuzzy ordered set. We say that  $(X, r)$  is a  $r$ -fuzzy complete lattice if every nonempty subset of  $X$  has a  $r$ -infimum and a  $r$ -supremum.

Let  $X$  be a  $r$ -fuzzy ordered set and let  $f : X \rightarrow X$  be a map. We say that  $f$  is  $r$ -fuzzy monotone if for all  $x, y \in X$  with  $r(x, y) \geq r(y, x)$ , then  $r(f(x), f(y)) \geq r(f(y), f(x))$ .

We denote the set of all fixed points of  $f$  by  $\text{Fix}(f)$ .

### 3. THE RESULTS

In this section, we establish a fuzzy version of Tarski’s fixpoint Theorem [8]. More precisely, we show the following:

**Theorem 3.1.** *Let  $(X, r)$  be a nonempty  $r$ -fuzzy complete lattice and let  $f : X \rightarrow X$  be a  $r$ -fuzzy monotone map. Then the set  $\text{Fix}(f)$  of all fixed points of  $f$  is a nonempty  $r$ -fuzzy complete lattice.*

In this section, we shall we need the three following technical lemmas which their proofs will be given in the Appendix.

**Lemma 3.2.** *Let  $X$  be a nonempty  $r$ -fuzzy ordered set and let  $E$  be a nonempty fuzzy ordered subset of  $X$ . If  $\sup_r(E) = s$ , then we have*

$$\{x \in X : r(s, x) = r(x, s)\} = \{s\}.$$

**Lemma 3.3.** *Let  $(X, r)$  be a nonempty  $r$ -fuzzy ordered set and let  $s$  be the inverse fuzzy relation of  $r$ . Then,*

- (i) *The fuzzy relation  $s$  is a fuzzy order on  $X$ .*
- (ii) *Every  $r$ -fuzzy monotone map  $f : X \rightarrow X$  is also  $s$ -fuzzy monotone.*
- (iii) *If a nonempty subset  $A$  of  $X$  has a  $r$ -infimum, then  $A$  has a  $s$ -supremum and  $\inf_r(A) = \sup_s(A)$ .*
- (iv) *If a nonempty subset  $A$  of  $X$  has a  $r$ -supremum, then  $A$  has a  $s$ -infimum and  $\inf_s(A) = \sup_r(A)$ .*

(v) If  $(X, r)$  is a fuzzy complete lattice, then  $(X, s)$  is also a fuzzy complete lattice.

For starting the third Lemma, we have to introduce the following subset  $E$  of  $X$  by  $x \in E$  if and only if  $r(x, f(x)) \geq r(f(x), x)$  and  $r(f(x), y) \geq r(y, f(x))$  for all  $y \in A$ , where  $A$  is a subset of  $\text{Fix}(f)$ .

**Lemma 3.4.** *Let  $(X, r)$  be a nonempty  $r$ -fuzzy complete lattice and let  $f : X \rightarrow X$  be a  $r$ -fuzzy monotone map. Let us suppose that  $E$  is defined as above and  $t = \sup_r(E)$ . Then  $t$  is a fixed point of  $f$ .*

In order to prove Theorem 3.1, we need the following proposition:

**Proposition 3.5.** *Let  $(X, r)$  be a nonempty  $r$ -fuzzy complete lattice and let  $f : X \rightarrow X$  be a  $r$ -fuzzy monotone map. Then  $f$  has a greatest and least fixed points. Furthermore,*

$$\max_r(\text{Fix}(f)) = \sup_r \{x \in X : r(x, f(x)) \geq r(f(x), x)\} ,$$

and

$$\min_r(\text{Fix}(f)) = \inf_r \{x \in X : r(f(x), x) \geq r(x, f(x))\} .$$

**Proof of Proposition 3.2.** Let  $D$  be the fuzzy ordered subset defined by

$$D = \{x \in X : r(x, f(x)) \geq r(f(x), x)\} .$$

Since  $\min_r(X) \in D$ , so  $D$  is nonempty. Let  $d$  be the  $r$ -supremum of  $D$ .

*Claim 1.* The element  $d$  is the greatest fixed point of  $f$ . Indeed, as  $d = \sup_r(D)$ , then  $r(x, d) \geq r(d, x)$  for all  $x \in D$ . Since  $f$  is  $r$ -fuzzy monotone, so  $r(f(x), f(d)) \geq r(f(d), f(x))$ , for all  $x \in D$ . We know that  $r(x, f(x)) \geq r(f(x), x)$ , for every  $x \in D$ . Then by fuzzy transitivity, we obtain  $r(x, f(d)) \geq r(f(d), x)$ , for all  $x \in D$ . Thus,  $f(d)$  is a  $r$ -upper bound of  $D$ . On the other hand,  $d$  is the least  $r$ -upper bound of  $D$ . So,

$$(3.1) \quad r(d, f(d)) \geq r(f(d), d) .$$

From this and fuzzy monotonicity of  $f$ , we get

$$(3.2) \quad r(f(d), f(f(d))) \geq r(f(f(d)), f(d)) .$$

Hence, we get  $f(d) \in D$ . From this and as  $d = \sup_r(D)$ , then

$$(3.3) \quad r(f(d), d) \geq r(d, f(d)) .$$

By combining (3.1) and (3.3), we get  $r(d, f(d)) = r(f(d), d)$ . From Lemma 3.2, we conclude that we have  $f(d) = d$ . Now let  $x \in \text{Fix}(f)$ . Then  $x \in D$ . So  $\text{Fix}(f) \subset D$ . From this and as  $d$  is the  $r$ -supremum of  $D$ , then we deduce that  $d$  is a  $r$ -upper bound of  $\text{Fix}(f)$ . Since  $d \in \text{Fix}(f)$ . Therefore  $d$  is the greatest element of  $\text{Fix}(f)$ .

*Claim 2.* The map  $f$  has a least fixed point. Let  $s$  be the fuzzy inverse order relation of  $r$  and let  $B$  be the following ordered subset of  $X$  defined by

$$B = \{x \in X : r(f(x), x) \geq r(x, f(x))\} .$$

Since  $\min_r(X) \in B$ , then  $B \neq \emptyset$ . On the other hand, by the definition of inverse fuzzy relation, we have

$$B = \{x \in X : s(x, f(x)) \geq s(f(x), x)\} .$$

By hypothesis,  $(X, r)$  is a nonempty fuzzy complete lattice, then from Lemma 3.3,  $(X, s)$  is also a nonempty fuzzy complete lattice. Furthermore,  $f$  is  $s$ -fuzzy monotone. Then by Claim 1,  $f$  has a greatest fixed point  $l$  in  $(X, s)$  with

$$l = \sup_s \{x \in X : s(x, f(x)) \geq s(f(x), x)\} .$$

Thus  $l$  is a least fixed point of  $f$  in  $(X, r)$ . By Lemma 3.3, we get

$$l = \inf_r \{x \in X : r(f(x), x) \geq r(x, f(x))\} . \quad \square$$

Now we are able to give the proof of Theorem 3.1.

**Proof of Theorem 3.1.** Let  $X$  be a nonempty  $r$ -fuzzy complete lattice and  $f : X \rightarrow X$  be a  $r$ -fuzzy monotone map.

*First Step.* We shall prove that every nonempty subset  $A$  of  $\text{Fix}(f)$  has a  $r$ -infimum in  $(\text{Fix}(f), r)$ . Let  $E$  and  $F$  be the two following subsets of  $X$  defined by  $x \in E$  if and only if

$$r(x, f(x)) \geq r(f(x), x) \quad \text{and} \quad r(f(x), y) \geq r(y, f(x))$$

for all  $y \in A$ , and

$$F = \{x \in \text{Fix}(f) : r(x, y) \geq r(y, x) \quad \text{for all } y \in A\} .$$

By Proposition 2.5,  $\min_r(\text{Fix}(f))$  exists in  $(X, r)$ . Since  $\min_r(\text{Fix}(f)) \in F$ , then  $F \neq \emptyset$ . Let  $m = \sup_r(F)$  and  $t = \sup_r(E)$ . We claim that the element  $m$  is the  $r$ -infimum of  $A$  in  $(\text{Fix}(f), r)$ . Indeed, Since  $F \subset E$ , then  $r(\sup_r(F), \sup_r(E)) \geq r(\sup_r(E), \sup_r(F))$ . Thus  $r(m, t) \geq r(t, m)$ . On the other hand  $t \in F$ , hence  $r(t, m) \geq r(t, m)$ . It follows that we have  $r(t, m) = r(m, t)$ . From Lemma 3.2, we get  $m = t$ . By Lemma 3.4,  $t$  is a fixed point of  $f$ . Therefore  $A$  has a  $r$ -infimum in  $\text{Fix}(f)$ .

*Second Step.* We shall prove that every nonempty subset  $A$  of  $\text{Fix}(f)$  has a  $r$ -supremum in  $(\text{Fix}(f), r)$ . Let  $G$  be the following ordered subset of  $X$  defined by  $x \in G$  if and only if

$$r(y, f(x)) \geq r(f(x), y)$$

for all  $y \in A$ , and

$$r(f(x), x) \geq r(x, f(x)) .$$

By Proposition 3.5,  $\max_r(\text{Fix}(f))$  exists in  $(X, r)$ . As  $\max_r(\text{Fix}(f)) \in G$ , then  $G \neq \emptyset$  and  $p = \inf_r(G)$  exists in  $(X, r)$ . Let  $s$  be the fuzzy inverse order relation of  $r$ . Then we get,  $x \in G$  if and only if

$$s(f(x), y) \geq s(y, f(x))$$

for all  $y \in A$  and

$$s(x, f(x)) \geq s(f(x), x).$$

We know by Lemma 3.3 that  $(X, s)$  is a nonempty fuzzy complete lattice. Moreover,  $f$  is  $s$ -fuzzy monotone and  $p = \sup_s(G)$ . From Lemma 3.4, we get  $f(p) = p$ . On the other hand, by the first step above,  $p$  is the  $s$ -supremum of  $A$ . Therefore, we deduce by Lemma 3.3 that the element  $p$  is the  $r$ -infimum of  $A$  in  $(\text{Fix}(f), r)$ .  $\square$

#### 4. APPENDIX

In this section, we give the proofs of Lemmas 3.2, 3.3 and 3.4.

**Proof of Lemma 3.2.** Let  $s = \sup_r(E)$  and let  $x \in X$  such that  $r(s, x) = r(x, s)$ .

*Claim 1.* The element  $x$  is a  $r$ -upper bound of  $E$ . Indeed, if  $a \in E$ , then  $r(a, s) \geq r(s, a)$ . Since  $r(s, x) = r(x, s)$ , then by fuzzy transitivity we get  $r(a, s) \geq r(s, a)$  for all  $a \in E$  and our claim is proved.

*Claim 2.* The element  $x$  is a least  $r$ -upper bound of  $E$ . Indeed, if  $b$  is a  $r$ -upper bound of  $E$ , then  $r(s, b) \geq r(b, s)$ . As  $r(s, x) = r(x, s)$ , then  $r(x, b) \geq r(b, x)$ . It follows that  $x$  is a least  $r$ -upper bound of  $E$ . Hence  $x$  is a  $r$ -supremum of  $E$ .

By Claims 1 and 2, we deduce that the element  $x$  is a  $r$ -supremum of  $A$ . From hypothesis, the  $r$ -supremum of  $A$  is unique, therefore  $x = s$ .  $\square$

**Proof of Lemma 3.3.** (i) For all  $x \in X$ , we have  $s(x, x) = r(x, x) \in [0, 1]$ . Let  $x, y \in X$  such that  $s(x, y) + s(y, x) > 1$ . Since  $r(x, y) + r(y, x) = s(x, y) + s(y, x) > 1$ , so  $r(x, y) + r(y, x) > 1$ . By  $r$ -fuzzy antisymmetry, we deduce that we have  $x = y$ . Let  $x, y, z \in X$  with  $s(x, y) \geq s(y, x)$  and  $s(y, z) \geq s(z, y)$ . Then we have  $r(z, y) \geq r(y, z)$  and  $r(y, x) \geq r(x, y)$ . By  $r$ -fuzzy transitivity, we obtain  $r(z, x) \geq r(x, z)$ . Therefore we get  $s(x, z) \geq s(z, x)$ . Thus the fuzzy relation  $s$  is a fuzzy order on  $X$ .

(ii) Let  $x, y \in X$  with  $s(x, y) \geq s(y, x)$ . Then we get  $r(y, x) \geq r(x, y)$ . Since  $f$  is  $r$ -fuzzy monotone, hence  $r(f(y), f(x)) \geq r(f(x), f(y))$ . Therefore  $s(f(x), f(y)) \geq s(f(y), f(x))$ . Thus the map  $f$  is  $s$ -fuzzy monotone.

(iii) Let  $m = \sup_r(A)$ . Then  $r(x, m) \geq r(m, x)$ , for all  $x \in A$ . So  $s(m, x) \geq s(x, m)$ , for all  $x \in A$ . Thus  $m$  is a  $s$ -lower bound of  $A$ . Now let  $t$  be another  $s$ -lower bound of  $A$ . Hence  $s(t, x) \geq s(x, t)$ , for all  $x \in A$ . Then  $r(x, t) \geq r(t, x)$ . Thus  $t$  is a  $r$ -upper bound of  $A$ . From this and as  $m = \sup_r(A)$ , we deduce that we have  $r(m, t) \geq r(t, m)$ . So  $s(t, m) \geq s(m, t)$ . Thus  $m$  is a greatest  $s$ -lower bound of  $A$ . Suppose that  $p$  is another greatest  $s$ -lower bound of  $A$ . By using a similar proof as above we deduce that  $p$  is a least  $r$ -upper bound of  $A$ . By hypothesis, the  $r$ -supremum of  $A$  is unique. Therefore, we conclude that  $p = m$ . Thus  $m = \inf_s(A)$ .

(iv) Since  $s$  is the inverse fuzzy relation of  $r$ , then  $r$  is the inverse fuzzy relation of  $s$ . By (iii), we get  $\inf_s(A) = \sup_r(A)$ .

(v) Let  $A$  be a nonempty set in  $X$ . Then  $A$  has a  $r$ -infimum and a  $r$ -supremum. From (iii) and (iv), we deduce that  $A$  has a  $s$ -infimum and a  $s$ -supremum. Thus  $(X, s)$  is a nonempty fuzzy complete lattice.  $\square$

**Proof of Lemma 3.4.** Let  $E$  be the subset of  $X$  defined by  $x \in E$  if and only if

$$r(x, f(x)) \geq r(f(x), x) \quad \text{and} \quad r(f(x), y) \geq r(y, f(x))$$

for all  $y \in A$ .

By Proposition 2.5,  $\min_r(\text{Fix}(f))$  exists in  $(X, r)$ . As  $\min_r(\text{Fix}(f)) \in E$ , then  $E \neq \emptyset$  and  $t = \sup_r(E)$  exists in  $X$ . We claim that we have:  $t = f(t)$ . Indeed, since for all  $x \in E$ , we have  $r(x, t) \geq r(t, x)$  and as  $f$  is  $r$ -fuzzy monotone, then

$$(4.1) \quad r(f(x), f(t)) \geq r(f(t), f(x)), \quad \text{for all } x \in E.$$

By definition, we have

$$(4.2) \quad r(x, f(x)) \geq r(f(x), x), \quad \text{for all } x \in E.$$

From (4.1) and (4.2) and fuzzy-transitivity, we get  $r(x, f(t)) \geq r(f(t), x)$  for all  $x \in E$ . Thus  $f(t)$  is a  $r$ -upper bound of  $E$ . From this and as  $t = \sup_r(E)$  so

$$(4.3) \quad r(t, f(t)) \geq r(t, f(t)).$$

From (4.3) and fuzzy monotonicity of  $f$ , we obtain

$$(4.4) \quad r(f(t), f(f(t))) \geq r(f(f(t)), f(t)).$$

Now let  $y \in A$ . Then for all  $x \in E$ , we have  $r(f(x), y) \geq r(y, f(x))$ . By using (4.2) and  $r$ -fuzzy transitivity, we obtain  $r(x, y) \geq r(y, x)$  for all  $x \in E$ . Thus every element of  $A$  is a  $r$ -upper bound of  $E$ . Since  $t$  is the least  $r$ -upper bound of  $E$ , then we get  $r(t, y) \geq r(y, t)$ , for all  $y \in A$ . Then by fuzzy monotonicity of  $f$ , we deduce that we have

$$(4.5) \quad r(f(t), y) \geq r(y, t), \quad \text{for all } y \in A.$$

Combining (4.4) and (4.5) we get  $f(t) \in E$ . On the other hand the element  $t$  is the  $r$ -supremum of  $E$ , then we deduce that we have

$$(4.6) \quad r(f(t), t) \geq r(t, f(t)).$$

By using (4.3) and (4.6) we deduce that we have  $r(f(t), t) = r(t, f(t))$ . Therefore by Lemma 3.2, we conclude that we have  $f(t) = t$ .  $\square$

## REFERENCES

- [1] Beg, I., *Fixed points of fuzzy monotone maps*, Arch. Math. (Brno) **35** (1999), 141–144.
- [2] Beg, I., *Fixed points of fuzzy multivalued mappings with values in fuzzy orders sets*, J. Fuzzy Math. **6**(1) (1998), 127–131.
- [3] Beg, I., *Fixed points of expansive mapping on fuzzy preordered sets*, J. Fuzzy Math. **7**(2) (1999), 746–749.
- [4] Billot, A., *Economic theory of fuzzy equilibria*, Lecture Notes in Econom. and Math. Systems **373**, Springer-Verlag, Berlin 1992.

- [5] Fang, J. X., *On fixed point theorems in fuzzy metric spaces*, Fuzzy Sets and Systems **46** (1992), 107–113.
- [6] Hadzic, O., *Fixed point theorems for multivalued mapping in some classes of fuzzy metric spaces*, Fuzzy Sets and Systems **29** (1989), 115–125.
- [7] Heilpern, S., *Fuzzy mapping and fixed point theorem*, J. Math. Anal. Appl. **83** (1981), 566–569.
- [8] Tarski, A., *A lattice-theoretical fixpoint theorem and its applications*, Pacific J. Math. **5** (1955), 285–309.
- [9] Zadeh, L. A., *Fuzzy sets*, Inform. and Control **8** (1965), 338–353.

UNIVERSITÉ CADI AYYAD, DÉPARTEMENT DE MATHÉMATIQUES  
FACULTÉ DES SCIENCES ET TECHNIQUES  
B. P. 523, BENI - MELLAL, MOROCCO  
*E-mail*: stouti@yahoo.com