

Časopis pro pěstování matematiky

Shigeyoshi Owa

On certain classes of p -valent functions

Časopis pro pěstování matematiky, Vol. 113 (1988), No. 4, 342--350

Persistent URL: <http://dml.cz/dmlcz/118352>

Terms of use:

© Institute of Mathematics AS CR, 1988

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON CERTAIN CLASSES OF p -VALENT FUNCTIONS

SHIGEYOSHI OWA, Osaka

(Received September 10, 1984)

Summary. We introduce the classes $T^*(n+p-1, \alpha)$ of analytic p -valent functions with negative coefficients by using the symbol $D^{n+p-1} f(z)$ defined by $\{z^n/(1-z)^{n+p}\} * f(z)$. The object of the present paper is to show distortion theorems and some closure theorems for functions $f(z)$ belonging to the class $T^*(n+p-1, \alpha)$. Further, we consider the modified Hadamard product of functions $f(z)$ in $T^*(n+p-1, \alpha)$.

Keywords: analytic function, p -valent function, distortion theorem, closure theorem, Hadamard product.

AMS (MOS) subject classification (1980): 30C45.

1. INTRODUCTION

Let $A(p)$ denote the class of functions $f(z)$ of the form

$$(1.1) \quad f(z) = z^p + \sum_{k=1}^{\infty} a_{p+k} z^{p+k} \quad (p \in N = \{1, 2, 3, \dots\})$$

which are analytic in the unit disk $U = \{z: |z| < 1\}$. Further, let $f(z) \in A(p)$ and $g(z) \in A(p)$. Then we denote by $f * g(z)$ the Hadamard product of $f(z)$ and $g(z)$, that is, if $f(z)$ is given by (1.1) and $g(z)$ is given by

$$(1.2) \quad g(z) = z^p + \sum_{k=1}^{\infty} b_{p+k} z^{p+k} \quad (p \in N),$$

then

$$(1.3) \quad f * g(z) = z^p + \sum_{k=1}^{\infty} a_{p+k} b_{p+k} z^{p+k}.$$

With the aid of the Hadamard product, we introduce the symbol

$$(1.4) \quad D^{n+p-1} f(z) = \left\{ \frac{z^n}{(1-z)^{n+p}} \right\} * f(z)$$

for n , where n is any integer greater than $-p$. Then we can observe that

$$(1.5) \quad D^{n+p-1} f(z) = \frac{z^p (z^{n-1} f(z))^{(n+p-1)}}{(n+p-1)!}.$$

In particular, the symbol $D^n f(z)$ was introduced by Ruscheweyh [7] and was named the n th order Ruscheweyh derivative of $f(z)$ by Al-Amiri [1].

Now, we introduce the following classes by using the symbol $D^{n+p-1} f(z)$.

Definition. We say that $f(z)$ is in the class $T^*(n + p - 1, \alpha)$, if $f(z)$ defined by

$$(1.6) \quad f(z) = z^p - \sum_{k=1}^{\infty} a_{p+k} z^{p+k} \quad (a_{p+k} \geq 0)$$

satisfies the condition

$$(1.7) \quad \operatorname{Re} \left\{ \frac{D^{n+p} f(z)}{D^{n+p-1} f(z)} \right\} > \alpha \quad (z \in U)$$

for $p \in N$, $n > -p$ and $0 \leq \alpha \leq \frac{1}{2}$.

Particularly, $T^*(n + p - 1, \alpha)$ is a subclass of $T(n, \alpha)$ which was studied by Goel and Sohi [3]. Further, by using the symbol $D^{n+p-1} f(z)$, other classes were studied by Goel and Sohi [4], [5], Sohi [8], Fukui and Sakaguchi [2] and Owa [6].

2. DISTORTION THEOREMS

Theorem 1. Let the function $f(z)$ be defined by (1.6). Then $f(z)$ is in the class $T^*(n + p - 1, \alpha)$ if and only if

$$(2.1) \quad \sum_{k=1}^{\infty} \frac{(n + p + k - 1)! \{(n + p)(1 - \alpha) + k\}}{k!} a_{p+k} \leq (n + p)! (1 - \alpha).$$

Equality holds for the function $f(z)$ given by

$$(2.2) \quad f(z) = z^p - \frac{1 - \alpha}{(n + p)(1 - \alpha) + 1} z^{p+1}.$$

Proof. Suppose that the inequality (2.1) holds and let $|z| = 1$. Then we obtain

$$\begin{aligned} (2.3) \quad & \left| \frac{D^{n+p} f(z)}{D^{n+p-1} f(z)} - 1 \right| = \\ & = \left| \frac{\sum_{k=1}^{\infty} (n + p + k - 1)(n + p + k - 2) \dots k a_{p+k} z^k}{(n + p)! - (n + p) \sum_{k=1}^{\infty} (n + p + k - 1)(n + p + k - 2) \dots (k + 1) a_{p+k} z^k} \right| \leq \\ & \leq \left| \frac{\sum_{k=1}^{\infty} (n + p + k - 1)(n + p + k - 2) \dots k a_{p+k} |z|^k}{(n + p)! - (n + p) \sum_{k=1}^{\infty} (n + p + k - 1)(n + p + k - 2) \dots (k + 1) a_{p+k} |z|^k} \right| \leq \end{aligned}$$

$$\leq \frac{\sum_{k=1}^{\infty} (n+p+k-1)(n+p+k-2)\dots k a_{p+k}}{(n+p)! - (n+p) \sum_{k=1}^{\infty} (n+p+k-1)(n+p+k-2)\dots (k+1) a_{p+k}} \leq \\ \leq 1 - \alpha.$$

Hence we can see that the values of $D^{n+p} f(z)/D^{n+p-1} f(z)$ lie in a circle centered at $w = 1$ whose radius is $(1 - \alpha)$. Consequently, we can observe that the function $f(z)$ satisfies the condition (1.7), hence $f(z)$ belongs to the class $T^*(n + p - 1, \alpha)$.

For the converse, suppose that the function $f(z)$ belongs to the class $T^*(n + p - 1, \alpha)$. Then we get

$$(2.4) \quad Re \left\{ \frac{D^{n+p} f(z)}{D^{n+p-1} f(z)} \right\} = \\ = Re \left\{ \frac{(n+p)! - \sum_{k=1}^{\infty} (n+p+k)(n+p+k-1)\dots(k+1) a_{p+k} z^k}{(n+p)! - (n+p) \sum_{k=1}^{\infty} (n+p+k-1)(n+p+k-2)\dots(k+1) a_{p+k} z^k} \right\} \\ > \alpha$$

for $z \in U$. Choose values of z on the real axis so that $D^{n+p} f(z)/D^{n+p-1} f(z)$ is real. Clearing the denominator in (2.4) and letting $z \rightarrow 1^-$ through real values, we have

$$(2.5) \quad (n+p)! - \sum_{k=1}^{\infty} (n+p+k)(n+p+k-1)\dots(k+1) a_{p+k} \geq \\ \geq \alpha \{ (n+p)! - (n+p) \sum_{k=1}^{\infty} (n+p+k-1)(n+p+k-2)\dots(k+1) a_{p+k} \}$$

which implies (2.1).

Finally, we can see that the function $f(z)$ given by (2.2) is an extreme one for the theorem. This completes the proof of the theorem.

Corollary 1. *Let the function $f(z)$ defined by (1.6) be in the class $T^*(n + p - 1, \alpha)$. Then*

$$(2.6) \quad a_{p+k} \leq \frac{k! (n+p)! (1-\alpha)}{(n+p+k-1)! \{ (n+p)(1-\alpha) + k \}}$$

for $k \geq 1$. The equality holds for the function $f(z)$ of the form

$$(2.7) \quad f(z) = z^p - \frac{k! (n+p)! (1-\alpha)}{(n+p+k-1)! \{ (n+p)(1-\alpha) + k \}} z^{p+k}.$$

Theorem 2. *Let the function $f(z)$ defined by (1.6) be in the class $T^*(n + p - 1, \alpha)$.*

Then

$$(2.8) \quad |f(z)| \geq |z|^p - \frac{1-\alpha}{(n+p)(1-\alpha)+1} |z|^{p+1}$$

and

$$|f(z)| \leq |z|^p + \frac{1-\alpha}{(n+p)(1-\alpha)+1} |z|^{p+1}$$

for $z \in U$. The results are sharp.

Proof. Since $f(z)$ belongs to the class $T^*(n+p-1, \alpha)$, in view of Theorem 1 we obtain

$$\begin{aligned} (2.10) \quad & (n+p)! \{(n+p)(1-\alpha)+1\} \sum_{k=1}^{\infty} a_{p+k} \leq \\ & \leq \sum_{k=1}^{\infty} \frac{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}}{k!} a_{p+k} \leq \\ & \leq (n+p)! (1-\alpha) \end{aligned}$$

which gives

$$(2.11) \quad \sum_{k=1}^{\infty} a_{p+k} \leq \frac{1-\alpha}{(n+p)(1-\alpha)+1}.$$

Consequently, we see that

$$\begin{aligned} (2.12) \quad & |f(z)| \geq |z|^p - |z|^{p+1} \sum_{k=1}^{\infty} a_{p+k} \geq \\ & \geq |z|^p - \frac{1-\alpha}{(n+p)(1-\alpha)+1} |z|^{p+1} \end{aligned}$$

and

$$\begin{aligned} (2.13) \quad & |f(z)| \leq |z|^p + |z|^{p+1} \sum_{k=1}^{\infty} a_{p+k} \leq \\ & \leq |z|^p + \frac{1-\alpha}{(n+p)(1-\alpha)+1} |z|^{p+1} \end{aligned}$$

for $z \in U$.

Further, by taking the function $f(z)$ given by

$$(2.14) \quad f(z) = z^p - \frac{1-\alpha}{(n+p)(1-\alpha)+1} z^{p+1}$$

we can observe that the results of the theorem are sharp.

Corollary 2. Let the function $f(z)$ defined by (1.6) be in the class $T^*(n+p-1, \alpha)$.

Then $f(z)$ is included in a disk with its center at the origin and radius r given by

$$(2.15) \quad r = \frac{(n+p+1)(1-\alpha)+1}{(n+p)(1-\alpha)+1}.$$

Theorem 3. Let the function $f(z)$ defined by (1.6) be in the class $T^*(n+p-1, \alpha)$. Then

$$(2.16) \quad |f'(z)| \geq p|z|^{p-1} - \frac{(p+1)(1-\alpha)}{(n+p)(1-\alpha)+1} |z|^p$$

and

$$(2.17) \quad |f'(z)| \leq p|z|^{p-1} + \frac{(p+1)(1-\alpha)}{(n+p)(1-\alpha)+1} |z|^p$$

for $z \in U$. The results are sharp.

Proof. Since $f(z)$ is in the class $T^*(n+p-1, \alpha)$, by virtue of Theorem 1 we have

$$\begin{aligned} (2.18) \quad & \frac{(n+p)! \{(n+p)(1-\alpha)+1\}}{p+1} \sum_{k=1}^{\infty} (p+k) a_{p+k} \leq \\ & \leq \sum_{k=1}^{\infty} \frac{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}}{k!} a_{p+k} \leq \\ & \leq (n+p)! (1-\alpha) \end{aligned}$$

which implies

$$(2.19) \quad \sum_{k=1}^{\infty} (p+k) a_{p+k} \leq \frac{(p+1)(1-\alpha)}{(n+p)(1-\alpha)+1}.$$

Hence, by using (2.19), we obtain

$$\begin{aligned} (2.20) \quad & |f'(z)| \geq p|z|^{p-1} - |z|^p \sum_{k=1}^{\infty} (p+k) a_{p+k} \geq \\ & \geq p|z|^{p-1} - \frac{(p+1)(1-\alpha)}{(n+p)(1-\alpha)+1} |z|^p \end{aligned}$$

and

$$\begin{aligned} (2.21) \quad & |f'(z)| \leq p|z|^{p-1} + |z|^p \sum_{k=1}^{\infty} (p+k) a_{p+k} \leq \\ & \leq p|z|^{p-1} + \frac{(p+1)(1-\alpha)}{(n+p)(1-\alpha)+1} |z|^p \end{aligned}$$

for $z \in U$.

Further, we can see that the results of the theorem are sharp for the function $f(z)$ given by (2.14).

Corollary 3. Let the function $f(z)$ defined by (1.6) be in the class $T^*(n + p - 1, \alpha)$. Then $f'(z)$ is included in a disk with its center at the origin and radius R given by

$$(2.22) \quad R = \frac{(np + p^2 + p + 1)(1 - \alpha) + p}{(n + p)(1 - \alpha) + 1}.$$

3. CLOSURE THEOREMS

Theorem 4. Let the functions

$$(3.1) \quad f_i(z) = z^p - \sum_{k=1}^{\infty} a_{p+k,i} z^{p+k} \quad (a_{p+k,i} \geq 0)$$

be in the class $T^*(n + p - 1, \alpha)$ for every $i = 1, 2, 3, \dots, m$. Then the function $h(z)$ defined by

$$(3.2) \quad h(z) = \sum_{i=1}^m c_i f_i(z) \quad (c_i \geq 0)$$

is also in the same class $T^*(n + p - 1, \alpha)$, where

$$(3.3) \quad \sum_{i=1}^m c_i = 1.$$

Proof. By means of the definition of $h(z)$, we can write

$$(3.4) \quad h(z) = z^p - \sum_{k=1}^{\infty} \left(\sum_{i=1}^m c_i a_{p+k,i} \right) z^{p+k}.$$

Now, since $f_i(z) \in T^*(n + p - 1, \alpha)$ for every $i = 1, 2, 3, \dots, m$, we obtain

$$(3.5) \quad \sum_{k=1}^{\infty} \frac{(n + p + k - 1)! (n + p)(1 - \alpha) + k}{k!} a_{p+k,i} \leq (n + p)! (1 - \alpha)$$

for every $i = 1, 2, 3, \dots, m$, by virtue of Theorem 1. Consequently, with the aid of (3.5) we can see that

$$\begin{aligned} (3.6) \quad & \sum_{k=1}^{\infty} \frac{(n + p + k - 1)! \{(n + p)(1 - \alpha) + k\}}{k!} \left(\sum_{i=1}^m c_i a_{p+k,i} \right) = \\ & = \sum_{i=1}^m c_i \left\{ \sum_{k=1}^{\infty} \frac{(n + p + k - 1)! \{(n + p)(1 - \alpha) + k\}}{k!} a_{p+k,i} \right\} \leq \\ & \leq \left(\sum_{i=1}^m c_i \right) (n + p)! (1 - \alpha) = (n + p)! (1 - \alpha). \end{aligned}$$

This proves that the function $h(z)$ belongs to the class $T^*(n + p - 1, \alpha)$.

Theorem 5. Let

$$(3.7) \quad f_p(z) = z^p$$

and

$$(3.8) \quad f_{p+k}(z) = z^p - \frac{k! (n+p)! (1-\alpha)}{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}} z^{p+k}$$

for $p \in N$, $n > -p$, $k \in N$ and $0 \leq \alpha \leq 1/2$. Then $f(z)$ is in the class $T^*(n+p-1, \alpha)$ if and only if it can be expressed in the form

$$(3.9) \quad f(z) = \sum_{k=0}^{\infty} \lambda_{p+k} f_{p+k}(z),$$

where $\lambda_{p+k} \geq 0$ and

$$(3.10) \quad \sum_{k=0}^{\infty} \lambda_{p+k} = 1.$$

Proof. Assume that

$$(3.11) \quad \begin{aligned} f(z) &= \sum_{k=0}^{\infty} \lambda_{p+k} f_{p+k}(z) = \\ &= z^p - \sum_{k=1}^{\infty} \frac{k! (n+p)! (1-\alpha)}{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}} \lambda_{p+k} z^{p+k}. \end{aligned}$$

Then we get

$$(3.12) \quad \begin{aligned} &\sum_{k=1}^{\infty} \left\{ \frac{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}}{k!} \right. \\ &\left. X \frac{k! (n+p)! (1-\alpha)}{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}} \lambda_{p+k} \right\} \leq (n+p)! (1-\alpha). \end{aligned}$$

By virtue of Theorem 1 this shows that $f(z)$ is in the class $T^*(n+p-1, \alpha)$.

Conversely, assume that $f(z)$ belongs to the class $T^*(n+p-1, \alpha)$. Again, by virtue of Theorem 1, we have

$$(3.13) \quad a_{p+k} \leq \frac{k! (n+p)! (1-\alpha)}{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}}$$

for $p \in N$, $n > -p$, $k \in N$ and $0 \leq \alpha \leq 1/2$. Next, setting

$$(3.14) \quad \lambda_{p+k} = \frac{(n+p+k-1)! \{(n+p)(1-\alpha)+k\}}{k! (n+p)! (1-\alpha)} a_{p+k}$$

for $p \in N$, $n > -p$, $k \in N$ and $0 \leq \alpha \leq 1/2$, and

$$(3.15) \quad \lambda_p = 1 - \sum_{k=1}^{\infty} \lambda_{p+k},$$

we have the representation (3.9). This completes the proof of the theorem.

4. MODIFIED HADAMARD PRODUCT

Let $f(z)$ be defined by (1.6) and let $g(z)$ be defined by

$$(4.1) \quad g(z) = z^p - \sum_{k=1}^{\infty} b_{p+k} z^{p+k} \quad (b_{p+k} \geq 0).$$

Further, let $f * g(z)$ denote the modified Hadamard product of $f(z)$ and $g(z)$, that is,

$$(4.2) \quad f * g(z) = z^p - \sum_{k=1}^{\infty} a_{p+k} b_{p+k} z^{p+k}.$$

Theorem 6. *Let the functions $f_i(z)$ defined by (3.1) be in the classes $T^*(n_i + p - 1, \alpha)$ for each $i = 1, 2, 3, \dots, m$, respectively. Then the modified Hadamard product $f_1 * f_2 * \dots * f_m(z)$ is in the class $T^*(n + p - 1)$, where $n = \min_{1 \leq i \leq m} \{n_i\}$.*

Proof. Since $f_i(z) \in T^*(n_i + p - 1, \alpha)$ for each $i = 1, 2, 3, \dots, m$, by using Theorem 1 we get

$$(4.3) \quad a_{p+k,i} \leq \frac{1 - \alpha}{(n_i + p)(1 - \alpha) + 1}$$

for each $i = 1, 2, 3, \dots, m$. Hence we can show that

$$\begin{aligned} (4.4) \quad & \sum_{k=1}^{\infty} \frac{(n + p + k - 1)! \{(n + p)(1 - \alpha) + k\}}{k!} \left(\prod_{i=1}^m a_{p+k,i} \right) \leq \\ & \leq (n + p)! (1 - \alpha) \frac{(n + p)(1 - \alpha) + 1}{1 - \alpha} \prod_{i=1}^m \left(\frac{1 - \alpha}{(n_i + p)(1 - \alpha) + 1} \right) \leq \\ & \leq (n + p)! (1 - \alpha) \end{aligned}$$

with the aid of (4.3) and Theorem 1. This shows that the modified Hadamard product $f_1 * f_2 * \dots * f_m(z)$ is in the class $T^*(n + p - 1, \alpha)$.

Corollary 4. *Let the functions $f_i(z)$ defined by (3.1) be in the same class $T^*(n + p - 1, \alpha)$ for every $i = 1, 2, 3, \dots, m$. Then the modified Hadamard product $f_1 * f_2 * \dots * f_m(z)$ is also in the class $T^*(n + p - 1, \alpha)$.*

References

- [1] H. S. Al-Amiri: On Ruscheweyh derivative. Ann. Polon. Math. 38 (1980), 87–94.
- [2] S. Fukui and K. Sakaguchi: An extension of a theorem of S. Ruscheweyh. Bull. Fac. Edu. Wakayama Univ. Nat. Sci. 29 (1980), 1–3.
- [3] R. M. Goel and N. S. Sohi: A new criterion for univalence and its applications. Glas. Mat. 16 (1981), 39–49.
- [4] R. M. Goel and N. S. Sohi: A new criterion for p -valent functions. Proc. Amer. Math. Soc. 78 (1980), 353–357.

- [5] R. M. Goel and N. S. Sohi: New criteria for p -valence. Indian J. Pure Appl. Math. 11 (1980), 1356–1360.
- [6] S. Owa: On new criteria for p -valent functions. Indian J. Pure Appl. Math. 13 (1982), 920–930.
- [7] St. Ruscheweyh: New criteria for univalent functions. Proc. Amer. Math. Soc. 49 (1975), 109–115.
- [8] N. S. Sohi: On a subclass of p -valent functions. Indian J. Pure Appl. Math. 11 (1980), 1504–1508.

Souhrn

O JISTÝCH TŘÍDÁCH p -ZNAČNÝCH FUNKCÍ

SHIGEYOSHI OWA

Autor zavádí třídy $T^*(n+p-1, \alpha)$ analytických p -značných funkcí s negativními koeficienty s použitím symbolu $D^{n+p-1} f(z)$ definovaného pomocí $\{z^p/(1-z)^{n+p}\} * f(z)$. Cílem práce je dokázat jisté věty o distorsi a o uzávěru pro funkce $f(z)$ patřící do třídy $T^*(n+p-1, \alpha)$. Je studován rovněž modifikovaný Hadamardův součin funkcí $f(z)$ v $T^*(n+p-1, \alpha)$.

Резюме

О НЕКОТОРЫХ КЛАССАХ p -ЗНАЧНЫХ ФУНКЦИЙ

SHIGEYOSHI OWA

Автор, пользуясь символом $D^{n+p-1} f(z)$, определённым при помощи $\{z^n/(1-z)^{n+p}\} * f(z)$, вводит классы $T^*(n+p-1, \alpha)$ p -значных аналитических функций с отрицательными коэффициентами. Целью статьи является доказательство некоторых теорем о дисторсии и замыкания для функций $f(z)$ из класса $T^*(n+p-1, \alpha)$. Кроме того изучается модифицированное произведение Адамарда функций из $T^*(n+p-1, \alpha)$.

Author's address: Department of Mathematics, Kinki University, Higashi-Osaka, Osaka 577, Japan.