

Sławomir Turek

A note on universal minimal dynamical systems

Commentationes Mathematicae Universitatis Carolinae, Vol. 32 (1991), No. 4, 781--783

Persistent URL: <http://dml.cz/dmlcz/118459>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1991

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A note on universal minimal dynamical systems

SŁAWOMIR TUREK

Abstract. Let $M(G)$ denote the phase space of the universal minimal dynamical system for a group G . Our aim is to show that $M(G)$ is homeomorphic to the absolute of $D^{2^{\omega}}$, whenever G is a countable Abelian group.

Keywords: dynamical system, universal minimal dynamical system, Abelian group, absolute

Classification: 54H20

Let G be a group and X a compact space. We call an **action** of the group G on space X a homomorphism Φ from G into the group of homeomorphisms of X . The pair (X, G) is called a **dynamical system** and the space X a **phase space** of the system (X, G) ; we will write gx for $\Phi(g)(x)$, $g \in G$, $x \in X$.

A dynamical system (X, G) is called **minimal** if the set $\{gx : g \in G\}$ is dense in X for each $x \in X$. The system (X, G) is minimal iff for each non-empty open set $U \subseteq X$ there are $g_1, \dots, g_n \in G$ such that $g_1U \cup \dots \cup g_nU = X$.

Let (X, G) and (Y, G) be dynamical systems and let $\varphi : X \rightarrow Y$ be a continuous map. If $\varphi \circ g = g \circ \varphi$ for any $g \in G$ then φ is called a **homomorphism** of the system (X, G) into the system (Y, G) . If in addition φ is a homeomorphism of spaces, then φ is called an **isomorphism** of dynamical systems.

The dynamical system (X, G) is called a **universal minimal dynamical system** for a group G if the following conditions hold:

- (a) (X, G) is a minimal dynamical system,
- (b) if (Y, G) is a minimal dynamical system then there exists a homomorphism $\varphi : (X, G) \rightarrow (Y, G)$.

The well-known results of Ellis [4; 7.13, 7.16] say that for every group G there is a universal minimal dynamical system which is unique up to an isomorphism. The phase space of this system is homeomorphic to a closed subspace of the Čech–Stone compactification of the discrete space G . Let $M(G)$ denote the phase space of the universal minimal dynamical system for a group G .

It was proved by van Douwen [3] that for every infinite Abelian group G $\pi w(M(G)) > |G|$, where $\pi w(X)$ denotes π -weight of X . Balcar and Błaszczyk [1] have shown that if (X, G) is a minimal dynamical system and X is an extremally disconnected space and G is a countable group then X is homeomorphic to the absolute of the Cantor cube $D^{\pi w(X)}$. On the other hand, it is known (cf. van Douwen [3]) that $M(G)$ has to be extremally disconnected. We will show that if G

is an Abelian group then $\pi w(M(G)) = 2^{|G|}$. Therefore $M(G)$ is homeomorphic to the absolute of D^{2^ω} , for every countable Abelian group G .

A continuous map $\varphi : X \rightarrow Y$ is **semi-open** if $\text{int } \varphi(U) \neq \emptyset$ for every non-empty open set $U \subseteq X$.

The following lemma is known; see e.g. [6]. We will include its proof for completeness.

Lemma 1. *Homomorphisms of minimal dynamical systems are semi-open and “onto”.*

PROOF: Let φ be a homomorphism of a minimal system (X, G) into a minimal system (Y, G) . If $x \in X$ then $\{g\varphi(x) : g \in G\}$ is dense in Y . Hence $\varphi(X) = \varphi(\text{cl}\{gx : g \in G\}) = \text{cl}\varphi(\{gx : g \in G\}) = \text{cl}\{\varphi(gx) : g \in G\} = \text{cl}\{g\varphi(x) : g \in G\} = Y$.

Let U be a non-empty open subset of X . Let us choose a non-empty open set V so that $\text{cl } V \subseteq U$. Since (X, G) is minimal, then there exist g_1, \dots, g_n from G such that $g_1V \cup \dots \cup g_nV = X$. Thus

$$\begin{aligned} Y &= \varphi(X) = \varphi(g_1V \cup \dots \cup g_nV) = \varphi(g_1V) \cup \dots \cup \varphi(g_nV) = \\ &= g_1\varphi(V) \cup \dots \cup g_n\varphi(V) \end{aligned}$$

and hence

$$\emptyset \neq \text{int } \varphi(\text{cl } V) \subseteq \text{int } \varphi(U).$$

□

Lemma 2. *If there exists a semi-open map of X onto Y , then $\pi w(Y) \leq \pi w(X)$.*

The proof of the above lemma is clear. □

Let G be an Abelian group and let $\text{Hom}(G, \mathbf{T})$ denote the group of all homomorphisms from G into the circle group $\mathbf{T} = \{z \in \mathbf{C} : |z| = 1\}$. It is well known that $\text{Hom}(G, \mathbf{T})$ is point-separating and the power of $\text{Hom}(G, \mathbf{T})$ equals $2^{|G|}$; see [5; 22.17, 24.47].

Let the homomorphism $e : G \rightarrow \mathbf{T}^{\text{Hom}(G, \mathbf{T})}$ be defined by the formula:

$$e(g)(h) = h(g), \quad \text{for } g \in G, \quad h \in \text{Hom}(G, \mathbf{T}).$$

The range $e(G)$ is a subgroup of the compact topological group $\mathbf{T}^{\text{Hom}(G, \mathbf{T})}$, where $\mathbf{T}^{\text{Hom}(G, \mathbf{T})}$ is regarded with the Tichonoff topology. Hence $bG = \text{cl}(e(G))$ is a compact topological group. The group bG is the so-called Bohr compactification of the (discrete) Abelian group G .

It is not hard to see that G acts on bG in the following way:

$$\Phi(g)(x) = e(g) \cdot x; \quad g \in G, \quad x \in bG.$$

Then (bG, G) forms a minimal dynamical system. Indeed, if $x \in bG$ then $f_x : bG \rightarrow bG$ defined by $f_x(y) = x \cdot y$, is a homeomorphism. Hence $\{gx : g \in G\} = f_x(e(G))$ is dense in bG .

The following lemma is known; see e.g. [2; 3.6. (ii)].

Lemma 3. *If X is a topological group then $w(X) = \pi w(X)$.*

For any locally compact Abelian group K , let \hat{K} denote the group of all the continuous homomorphisms of K into \mathbf{T} , endowed with a compact-open topology. It is known that $w(K) = w(\hat{K})$ and $bK = (\hat{K})_d$, where X_d denotes a space X with a discrete topology; see [5; 24.14, 26.12].

Theorem. *If G is an Abelian group then $\pi w(M(G)) = 2^{|G|}$.*

PROOF: Since (bG, G) is a minimal dynamical system, then there exists a homomorphism $\varphi : (M(G), G) \rightarrow (bG, G)$. Lemmas 1, 2 and 3 imply

$$w(bG) = \pi w(bG) \leq \pi w(M(G)).$$

From the above remarks, we get

$$w(bG) = w((\hat{G})_d) = w((\hat{G})_d) = |\hat{G}| = |\text{Hom}(G, \mathbf{T})| = 2^{|G|},$$

because G is a discrete space.

The inequality $\pi w(M(G)) \leq 2^{|G|}$ follows from the fact that $M(G)$ is homeomorphic to a closed subset of βG . □

The result of [1] leads to the following:

Corollary. *If G is a countable Abelian group then $M(G)$ is homeomorphic to the absolute of D^{2^ω} .*

REFERENCES

- [1] Balcar B., Błaszczyk A., *On minimal dynamical systems on Boolean algebras*, Comment. Math. Univ. Carolinae **31** (1990), 7–11.
- [2] Comfort W.W., *Topological Groups*, Handbook of set-theoretic topology, North-Holland, 1984, 1143–1260.
- [3] van Douwen E.K., *The maximal totally bounded group topology on G and the biggest minimal G -space, for Abelian groups G* , Topology and its Appl. **34** (1990), 69–91.
- [4] Ellis R., *Lectures on Topological Dynamics*, Benjamin, New York, 1969.
- [5] Hewitt E., Ross K.A., *Abstract Harmonic Analysis I*, Springer, Berlin, 1963.
- [6] van der Woude J., *Topological Dynamix*, Mathematisch Centrum, Amsterdam, 1982.

INSTYTUT MATEMATYKI UNIwersytetu ŚLĄskiego, ul. BANKOWA 14, KATOWICE, POLAND

(Received August 23, 1991)