

Wolfgang Rother

Bifurcation for some semilinear elliptic equations when the linearization has no eigenvalues

Commentationes Mathematicae Universitatis Carolinae, Vol. 34 (1993), No. 1, 125--138

Persistent URL: <http://dml.cz/dmlcz/118562>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1993

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Bifurcation for some semilinear elliptic equations when the linearization has no eigenvalues

WOLFGANG ROTHER

Abstract. We prove existence and bifurcation results for a semilinear eigenvalue problem in \mathbb{R}^N ($N \geq 2$), where the linearization $-\Delta$ has no eigenvalues. In particular, we show that under rather weak assumptions on the coefficients $\lambda = 0$ is a bifurcation point for this problem in H^1, H^2 and L^p ($2 \leq p \leq \infty$).

Keywords: bifurcation point, variational method, eigenvalues, exponential decay, standing waves

Classification: 35P30, 35A30

1. Introduction and presentation of the results.

In the present paper, we consider the nonlinear eigenvalue problem

$$(1.1) \quad -\Delta u - q(x)|u|^{\sigma_1}u + r(x)|u|^{\sigma_2}u = \lambda u \quad \text{in } \mathbb{R}^N,$$

where $N \geq 2$ and σ_1 and σ_2 are positive constants such that $\sigma_1 < 4/N$. In particular, we are interested in the question if $\lambda = 0$ is a bifurcation point for the equation (1.1).

Since the problem (1.1) is considered in \mathbb{R}^N , the linearization $-\Delta$ has no eigenvalues and $\lambda = 0$ is the infimum of the spectrum of $-\Delta$. In case that $r \equiv 0$, this problem has been studied by many authors. See for instance [5]–[7], [9], [13]–[18] and the literature quoted therein. In case that $r \not\equiv 0$, we only know some existence results for the equation (1.1) (see [1], [2], [8] and [12]), but no bifurcation results. In the following, we will close this gap by presenting some bifurcation results for the general case.

We always assume that the functions q and r satisfy the subsequent conditions:

(A) The functions $q, r : \mathbb{R}^N \rightarrow \mathbb{R}$ are measurable and r fulfills $r(x) \geq 0$ for almost all $x \in \mathbb{R}^N$.

(B) There exist a constant $0 < a \leq 2 - (\sigma_1 N/2)$ and an open ball $B \subset \mathbb{R}^N$, satisfying $B \neq \emptyset$ and $0 \notin \bar{B}$ (\bar{B} is the closure of B), such that $q(x) \geq f(x)|x|^{-a}$ holds for almost all $x \in \zeta$, where $\zeta = \{tx; t \geq 1, x \in B\}$ and $f : \zeta \rightarrow [0, \infty)$ is a measurable function satisfying $f(x) \rightarrow \infty$ as $|x| \rightarrow \infty$.

Moreover, we assume that there exists a constant \mathcal{K} such that

$$r(x) \leq \mathcal{K}|x|^b \quad \text{holds for almost all } x \in \zeta,$$

where b is defined by $b = (2 - a)(\sigma_2/\sigma_1) - 2$.

(C) The functions r and $q_- = \min(q, 0)$ are locally integrable.

(D) The function $q_+ = \max(q, 0)$ can be written as $q_+ = q_1 + q_2$, where

(D1) the function q_1 satisfies $0 \leq q_1 \in L^\infty$, and $q_1(x)$ tends uniformly to zero as $|x| \rightarrow \infty$,

(D2) and the function q_2 satisfies $0 \leq q_2 \in L^{p_0}$ for some constant

$$2N/(4 - \sigma_1 N) < p_0 < \infty.$$

We want to point out that the above assumptions allow the function q to decay exponentially to $-\infty$ or faster in some direction, and allow the function r to increase exponentially to $+\infty$ or faster in some direction.

Theorem 1.1. *Suppose that the functions q and r satisfy the assumptions (A)–(D) and that the constant a is defined as in condition (B). Then, there exists a constant $\mu_a \in (0, \infty]$, depending on a , such that for each $\mu \in (0, \mu_a)$ there exists a nonpositive constant $\lambda(\mu)$ and a nontrivial nonnegative function $u_\mu \in H^1 \cap L^\infty$ which solves equation (1.1) in the sense of distributions. In case that $a = 2 - (\sigma_1 N/2)$, we have $\mu_a = \infty$. Moreover, it follows that $\lambda(\mu) \rightarrow 0$, $\|u_\mu\|_{H^1} \rightarrow 0$ and, if $p \in [2, \infty]$, that $\|u_\mu\|_p \rightarrow 0$ as $\mu \rightarrow 0$. Hence, $\lambda = 0$ is a bifurcation point for equation (1.1) in H^1 and in L^p for $p \in [2, \infty]$.*

Corollary 1.2. (a) *If $q_-, r \in L^p_{\text{loc}}$ holds for some constant $p > N/2$, then u_μ is positive and locally Hölder continuous.*

(b) *If q and r are locally Hölder continuous, then we have $u_\mu \in C^2$ and the equation (1.1) holds in the classical sense.*

Corollary 1.3. *Suppose in addition to (A)–(D) that $p_0 \geq 2$ and that $q, r \in L^\infty + L^2$. Then, it follows that $u_\mu \in H^2$ and that $\|u_\mu\|_{H^2} \rightarrow 0$ as $\mu \rightarrow 0$. Thus, $\lambda = 0$ is a bifurcation point for (1.1) in H^2 .*

Remark 1.4. In case that $r \equiv 0$, Corollary 1.3 improves Theorem 2.6 (c) in [13]. In [13] it is assumed that q is nonnegative, that $q = q_+$ satisfies condition (D) and that $p_0 \geq 2$. Moreover, it is assumed

(i) that there exist constants $A > 0$ and $0 \leq t < 2 - (\sigma_1 N/2)$ such that $q(x) \geq A(1 + |x|)^{-t}$ holds a.e. in \mathbb{R}^N . In case that $N \geq 3$ the author requires additionally

(ii) that $\sigma_1 < 2/(N - 2)$ and $p_0 > 2N/(2 - \sigma_1(N - 2))$. Hence, Corollary 1.3 shows that the condition (i) can be weakened considerably and that condition (ii) is superfluous.

The solutions of the equation (1.1) supply standing waves for nonlinear Klein-Gordon and Schrödinger equations. So, from the standpoint of physics it is an interesting question if the solutions of (1.1) decay exponentially to 0 at infinity.

For the proof of the exponential decay to 0 we need an additional assumption:

(E) There exists a constant $R_0 > 0$ such that q_2 satisfies

$$q_2(x) = 0 \text{ for almost all } |x| \geq R_0.$$

Theorem 1.5. *Suppose that $\sigma_2 \leq \sigma_1$ and that the functions q and r satisfy the assumptions (A)–(E). Then, for each $\mu \in (0, \mu_a)$ the function u_μ decays exponentially to 0 at infinity.*

Theorem 1.6. *Suppose that $\sigma_1 < \sigma_2$ and that the functions q and r satisfy the assumptions (A)–(E). Then, there exists a decreasing sequence $(\mu_n) \subset (0, \mu_a)$ such that $\lim_{n \rightarrow \infty} \mu_n = 0$ and u_{μ_n} decays exponentially to 0 at infinity.*

The proofs for Theorem 1.5–1.6 can be found in § 4.

2. Some preliminaries.

For $p \in [1, \infty]$, $L^p = L^p(\mathbb{R}^N)$ and $L^p_{loc} = L^p_{loc}(\mathbb{R}^N)$ are the usual Lebesgue spaces and $\|\cdot\|_p$ is the norm on L^p . If $1 < p < \infty$, then the dual index p' of p is defined by $p' = p/(p - 1)$. Furthermore, H^k ($k = 1, 2$) is the Hilbert space $H^k(\mathbb{R}^N) = W^{k,2}(\mathbb{R}^N)$. The norm on H^1 is given by $\|u\|_{H^1} = (\|\nabla u\|_2^2 + \|u\|_2^2)^{1/2}$ and the norm on H^2 by $\|u\|_{H^2} = (\|\Delta u\|_2^2 + \|\nabla u\|_2^2 + \|u\|_2^2)^{1/2}$. Finally, $C_0^\infty = C_0^\infty(\mathbb{R}^N)$ denotes the set of all functions which have compact support and derivatives of any order.

If $N = 2$, then it follows from the Sobolev imbedding theorem that for each $p \in [2, \infty)$ there exists a constant A_p such that

$$(2.1) \quad \|u\|_p \leq A_p \|u\|_{H^1} \quad \text{holds for all } u \in H^1.$$

In case that $N \geq 3$, we define $2^* = 2N/(N - 2)$. Then, there exists a constant C_0 such that

$$(2.2) \quad \|u\|_{2^*} \leq C_0 \|\nabla u\|_2 \quad \text{holds for all } u \in H^1.$$

In particular we see that for each $p \in [2, 2^*]$ there exists a constant B_p such that

$$(2.3) \quad \|u\|_p \leq B_p \|u\|_{H^1} \quad \text{holds for all } u \in H^1.$$

Let F be one of the Banach spaces H^1 , H^2 or L^p . Then a real number λ is called a bifurcation point for the equation (1.1) in F if and only if there exists a sequence $(\lambda_n, u_n) \subset \mathbb{R} \times F$ such that $u_n \not\equiv 0$, $\lambda_n \rightarrow \lambda$, $\|u_n\|_F \rightarrow 0$ ($n \rightarrow \infty$) and

$$\int \nabla u_n \nabla \varphi \, dx - \int q|u_n|^{\sigma_1} u_n \varphi \, dx + \int r|u_n|^{\sigma_2} u_n \varphi \, dx = \lambda_n \int u_n \varphi \, dx$$

holds for all $\varphi \in C_0^\infty$ and $n \in \mathbb{N}$.

When the domain of integration is not indicated, it is understood to be \mathbb{R}^N .

Lemma 2.1. *Let $v \in H^1$ be a nonnegative function. Then, there exists a sequence (φ_n) of nonnegative functions $\varphi_n \in C_0^\infty$ such that*

$$\varphi_n \rightarrow v \quad \text{in } H^1.$$

PROOF: The functions η_n ($n \in \mathbb{N}$) may be chosen such that $\eta_n \in C_0^\infty$, $0 \leq \eta_n \leq 1$, $\eta_n(x) = 1$ holds for $|x| \leq n$, $\eta_n(x) = 0$ if $|x| \geq n + 1$ and $\|\nabla \eta_n\|_\infty \leq C$, where the constant C is independent of n . Then $\eta_n v \rightarrow v$ in H^1 .

For a function $u \in L^1_{loc}$, the regularization u_ε may be defined as in [3, p. 147]. Then, we can find a sequence (ε_n) of positive numbers ε_n , satisfying $\varepsilon_n \rightarrow 0$, such that $\varphi_n = (\eta_n v)_{\varepsilon_n} \rightarrow v$ in H^1 . □

Lemma 2.2. *Let $v \in H^1$ be a nonnegative function and, for $t > 0$, v_t may be defined by $v_t = \min(v, t)$. Then it follows that $v_t \in H^1$, $\partial_i v_t = \partial_i v$ holds almost everywhere in $\{x; v(x) \leq t\}$ and $\partial_i v_t = 0$ holds almost everywhere in $\{x; v(x) > t\}$. Moreover, for each $s \in [1, \infty)$, we have $0 \leq v_t^s \in H^1 \cap L^\infty$ and $\partial_i v_t^s = s v_t^{s-1} \partial_i v_t$ ($i = 1, \dots, N$).*

PROOF: The first part of the lemma follows from Lemma 1.1 in [10] and Theorem 7.8 in [3]. The functions η_n and the regularizations u_ε may be defined as in the proof of Lemma 2.1. Then, there exists a sequence of positive numbers (ε_n) such that $\varepsilon_n \rightarrow 0$ and

$$\varphi_n = (\eta_n v_t)_{\varepsilon_n} \longrightarrow v_t \text{ in } H^1.$$

Here, the functions φ_n satisfy $\varphi_n \in C_0^\infty$ and $0 \leq \varphi_n \leq t$. Since $\varphi_n \rightarrow v_t$ in L^2 , we can find a subsequence $(\varphi_{n(k)})$ of (φ_n) such that $\varphi_{n(k)}(x) \rightarrow v_t(x)$ for almost all $x \in \mathbb{R}^N$.

Now, suppose that $s > 1$. Then it follows that $\varphi_{n(k)}^s \in C_0^1$ and that

$$\partial_i \varphi_{n(k)}^s = s \varphi_{n(k)}^{s-1} \partial_i \varphi_{n(k)}.$$

Moreover, since $|v_t^s - \varphi_{n(k)}^s| \leq s |v_t - \varphi_{n(k)}| t^{s-1}$, we see that $\varphi_{n(k)}^s \rightarrow v_t^s$ in L^2 . Hence, we obtain: $\partial_i v_t^s = s v_t^{s-1} \partial_i v_t$. □

The following lemma can be found in [11, p. 93].

Lemma 2.3. *Suppose that $\varphi(t)$ ($t \in [t_0, \infty)$) is a nonnegative and nonincreasing function such that $\varphi(h) \leq C(h - t)^{-\gamma} \varphi(t)^\delta$ holds for all $h > t \geq t_0$. The constants γ and C are assumed to be positive and δ may satisfy $\delta > 1$. Then, for $d = C^{1/\gamma} \varphi(t_0)^{(\delta-1)/\gamma} 2^{\delta/(\delta-1)}$ it follows that $\varphi(t_0 + d) = 0$.*

3. Proof of the main results.

In the present paragraph, we will prove Theorem 1.1 and Corollary 1.2–1.3. We start with

Lemma 3.1. *There exist positive constants α and β , and for each $\varepsilon > 0$ a constant $K_\varepsilon > 0$, such that*

$$(2 + \sigma_1)^{-1} \int q_+ |u|^{2+\sigma_1} dx \leq \varepsilon \|\nabla u\|_2^2 + K_\varepsilon \left(\|u\|_2^{2+\alpha} + \|u\|_2^{2+\beta} \right)$$

holds for all $u \in H^1$.

PROOF: For $\varepsilon = \frac{1}{4}$, the proof can be found in [5, pp. 568–569]. For general $\varepsilon > 0$, the proof proceeds quite similarly. □

The nonlinear functional ξ may be defined by

$$\begin{aligned} \xi(u) = \frac{1}{2} \int |\nabla u|^2 dx - (2 + \sigma_1)^{-1} \int q |u|^{2+\sigma_2} dx \\ + (2 + \sigma_2)^{-1} \int r |u|^{2+\sigma_2} dx. \end{aligned}$$

By D , we denote the set

$$D = \{u \in H^1; \int |q_-||u|^{2+\sigma_1} dx < \infty \text{ and } \int r|u|^{2+\sigma_2} dx < \infty\}.$$

Moreover, for $\mu \geq 0$, we define $D_\mu = \{u \in D; \|u\|_2 \leq \mu\}$. Then, according to Lemma 3.1, we see that $I(\mu) = \inf_{u \in D_\mu} \xi(u)$ is a well defined real number.

Lemma 3.2. (a) *Suppose that the constant a in condition (B) satisfies $a = 2 - (\sigma_1 N/2)$. Then it follows that $I(\mu) < 0$ holds for all $\mu > 0$.*

(b) *Suppose that $a < 2 - (\sigma_1 N/2)$. Then, there exists a constant $\mu_a > 0$ such that $I(\mu) < 0$ holds for all $\mu \in (0, \mu_a)$.*

Remark 3.3. In the following, we define $\mu_a = \infty$ if $a = 2 - (\sigma_1 N/2)$.

PROOF OF LEMMA 3.2: The ball B may be defined as in condition (B) and ν may be a positive constant. Then, the function $\varphi_0 \in C_0^\infty$ may be chosen such that $\text{supp } \varphi_0 \subset B$ and $\|\varphi_0\|_2 = \nu$. Moreover, for each $t \geq 1$, we define $\varphi_t(x) = t^k \varphi_0(t^{-1}x)$, where $k = (a - 2)/\sigma_1$. Since $\|\varphi_t\|_2 = \nu t^{k+(N/2)}$, we see that $\varphi_t \in D_{\nu t^{k+(N/2)}}$ and that

$$\begin{aligned} I\left(\nu t^{k+(N/2)}\right) &\leq \xi(\varphi_t) = t^{2k+N-2} \left(\frac{1}{2} \int |\nabla \varphi_0(x)|^2 dx \right. \\ &\quad - t^{2+k\sigma_1} (2 + \sigma_1)^{-1} \int_B q(tx) |\varphi_0(x)|^{2+\sigma_1} dx \\ &\quad \left. + t^{2+k\sigma_2} (2 + \sigma_2)^{-1} \int_B r(tx) |\varphi_0(x)|^{2+\sigma_2} dx \right) \\ &\leq t^{2k+N-2} \left(\frac{1}{2} \int |\nabla \varphi_0(x)|^2 dx \right. \\ &\quad - \inf_{x \in B} f(tx) (2 + \sigma_1)^{-1} \int_B |x|^{-a} |\varphi_0(x)|^{2+\sigma_1} dx \\ &\quad \left. + \mathcal{K} (2 + \sigma_2)^{-1} \int_B |x|^b |\varphi_0(x)|^{2+\sigma_2} dx \right). \end{aligned}$$

Since $\inf_{x \in B} f(tx) \rightarrow \infty$ as $t \rightarrow \infty$, we can find a constant $t_0 \geq 1$ such that

$$(3.1) \quad I\left(\nu t^{k+(N/2)}\right) < 0 \text{ holds for all } t > t_0.$$

Now, suppose that $a = 2 - (\sigma_1 N/2)$. Then, we have $k + (N/2) = 0$. Hence, the part (a) of the lemma follows from (3.1) for $\nu = \mu$. In case that $a < 2 - (\sigma_1 N/2)$, we have $k + (N/2) < 0$. Then, the assertion of the part (b) follows from (3.1) if we define $\nu = 1$, $\mu_a = t_0^{k+(N/2)}$ and $\mu = t^{k+(N/2)}$. \square

Lemma 3.4. *For each $\mu \in (0, \mu_a)$ there exists a function $u_\mu \in D_\mu$ such that $u_\mu \geq 0$, $\|u_\mu\|_2 > 0$ and $\xi(u_\mu) = I(\mu)$.*

PROOF: Let $\mu \in (0, \mu_a)$, and $(v_n) \subset D$ may be a sequence such that $\xi(v_n) \rightarrow I(\mu)$. Then, we may assume without restriction that $\xi(v_n) \leq 0$ and that $v_n \geq 0$ holds for all n . Hence, we obtain from Lemma 3.1:

$$(3.2) \quad \begin{aligned} \frac{1}{4} \|\nabla v_n\|_2^2 + (2 + \sigma_1)^{-1} \int |q_-| |v_n|^{2+\sigma_1} dx \\ + (2 + \sigma_2)^{-1} \int r |v_n|^{2+\sigma_1} dx \leq K_{1/4} (\mu^{2+\alpha} + \mu^{2+\beta}). \end{aligned}$$

Since (v_n) is bounded in H^1 , we can find a subsequence of (v_n) , still denoted by (v_n) , and a $u_\mu \in H^1$ such that $v_n \xrightarrow{w} u_\mu$ in H^1 and $v_n(x) \rightarrow u_\mu(x)$ for almost all $x \in \mathbb{R}^N$. Then, it follows from the uniform boundedness principle, (3.2) and Fatou's lemma that $\|u_\mu\|_2 \leq \mu$, $\|\nabla u_\mu\|_2 \leq \liminf \|\nabla v_n\|_2$,

$$\int |q_-| |u_\mu|^{2+\sigma_1} dx \leq \liminf \int |q_-| |v_n|^{2+\sigma_1} dx < \infty$$

and

$$\int r |u_\mu|^{2+\sigma_2} dx \leq \liminf \int r |v_n|^{2+\sigma_2} dx < \infty.$$

Moreover, we see that $u_\mu \geq 0$. Since the imbedding $H^1(G) \rightarrow L^{(2+\sigma_1)p'_0}(G)$ is compact for all bounded balls G and $q_1(x) \rightarrow 0$ as $|x| \rightarrow \infty$, it follows that

$$\int q_+ |v_n|^{2+\sigma_1} dx \longrightarrow \int q_+ |u_\mu|^{2+\sigma_1} dx \quad (\text{see [5, p. 570]}).$$

Moreover, we obtain

$$I(\mu) \leq \xi(u_\mu) \leq \liminf \xi(v_n) = I(\mu) < 0$$

and consequently that $\xi(u_\mu) = I(\mu)$ and $\|u_\mu\|_2 > 0$. □

Lemma 3.5. *For $\mu \in (0, \mu_a)$, the function u_μ may be chosen as in Lemma 3.4. Then, it follows that*

$$\int \nabla u_\mu \nabla \varphi dx - \int q |u_\mu|^{\sigma_1} u_\mu \varphi dx + \int r |u_\mu|^{\sigma_2} u_\mu \varphi dx = \lambda(\mu) \int u_\mu \varphi dx$$

holds for all functions $\varphi \in C_0^\infty$, where

$$\lambda(\mu) = \|u_\mu\|_2^{-2} \left(\|\nabla u_\mu\|_2^2 - \int q |u_\mu|^{2+\sigma_1} dx + \int r |u_\mu|^{2+\sigma_2} dx \right).$$

PROOF: Let $\varphi \in C_0^\infty$. Then $d\xi(\|u_\mu\|_2 \|u_\mu + \varepsilon \varphi\|_2^{-1} (u_\mu + \varepsilon \varphi)) / d\varepsilon |_{\varepsilon=0} = 0$ implies the assertion. □

Lemma 3.6. *The constant $\lambda(\mu)$ may be defined as in Lemma 3.5. Then, we have $\lambda(\mu) \leq 0$.*

PROOF: For all $t \in (0, 1]$, we have

$$\xi(u_\mu) = I(\mu) \leq I(t\mu) \leq \xi(tu_\mu).$$

Hence $\lambda(\mu) = \|u_\mu\|_2^{-2} d\xi(tu_\mu)/dt |_{t=1} \leq 0$ implies the assertion. □

Proposition 3.7. *The constants α and β may be chosen as in Lemma 3.1. Then, there exists a constant C such that*

$$|\lambda(\mu)| \leq C(\mu^\alpha + \mu^\beta) \quad \text{and} \quad \|\nabla u_\mu\|_2^2 \leq C(\mu^{2+\alpha} + \mu^{2+\beta})$$

holds for all $\mu \in (0, \mu_a)$. Hence, $\lambda = 0$ is a bifurcation point for the equation (1.1) in H^1 .

PROOF: Since $\xi(u_\mu) < 0$, we obtain from Lemma 3.1 that

$$(3.3) \quad \|\nabla u_\mu\|_2^2 \leq 4K_{1/4}(\|u_\mu\|_2^{2+\alpha} + \|u_\mu\|_2^{2+\beta}) \leq 4K_{1/4}(\mu^{2+\alpha} + \mu^{2+\beta}).$$

Moreover, since $\lambda(\mu) \leq 0$, it follows from (3.3) and Lemma 3.1 that

$$\begin{aligned} |\lambda(\mu)| &= -\lambda(\mu) \leq \|u_\mu\|_2^{-2} \int q_+ |u_\mu|^{2+\sigma_1} dx \\ &\leq (2 + \sigma_1)(4K_{1/4} + K_1) \left(\|u_\mu\|_2^\alpha + \|u_\mu\|_2^\beta \right) \leq C(\mu^\alpha + \mu^\beta). \end{aligned}$$

□

Lemma 3.8. *For all nonnegative functions $v \in H^1$ we obtain*

$$(3.4) \quad \int \nabla u_\mu \nabla v \, dx \leq \lambda(\mu) \int u_\mu v \, dx + \int q_+ u_\mu^{1+\sigma_1} v \, dx$$

and, according to Lemma 3.6, that

$$(3.5) \quad \int \nabla u_\mu \nabla v \, dx \leq \int q_+ u_\mu^{1+\sigma_1} v \, dx.$$

PROOF: Clearly, the assertion holds for all nonnegative functions $v \in C_0^\infty$. Hence, the result follows from Lemma 2.1. □

Lemma 3.9. *Suppose that $N \geq 3$ and that $\int q_+ u_\mu^{1+\sigma_1+s} \, dx < \infty$ holds for some constant $s > 1$. Then, it follows that $u_\mu \in L^{2^*(s+1)/2}$.*

PROOF: For $t > 0$, the function v_t may be defined by $v_t = \min(u_\mu, t)$. Then, according to Lemma 2.2, we see that $0 \leq v_t^s \in H^1$. Inserting v_t^s in (3.5) shows that

$$4s(s+1)^{-2} \int |\nabla v_t^{(s+1)/2}|^2 \, dx \leq \int q_+ u_\mu^{1+\sigma_1+s} \, dx.$$

Hence, using (2.2) and letting $t \rightarrow \infty$, we obtain the assertion by Fatou's lemma. □

Lemma 3.10. *For each $p \in [2, \infty)$, we have $u_\mu \in L^p$.*

PROOF: For $N = 2$ and for $p \in [2, 2^*]$, if $N \geq 3$, the assertion follows from the Sobolev imbedding theorem. Now, suppose that $N \geq 3$ and that the constants r_n and s_n are defined by $r_n = 2^*(1 + \varepsilon_0)^n$ and $s_n = (r_n/p'_0) - 1 - \sigma_1$, where $\varepsilon_0 = (2^*/2p'_0) - (\sigma_1/2) - 1$. Here, the constant p_0 is defined as in condition (D2). Since $p_0 > 2N/(4 - \sigma_1N + 2\sigma_1)$ and $r_n \geq 2^*$, it follows that $\varepsilon_0 > 0$ and $s_n > 1$.

Now, assume that $u_\mu \in L^{r_n}$ holds for some $n \in \mathbb{N}_0$. Then $2 \leq 1 + \sigma_1 + s_n < (1 + \sigma_1 + s_n)p'_0 = r_n$ implies that

$$\int q_+ u_\mu^{1+\sigma_1+s_n} dx < \infty.$$

Hence, we obtain from Lemma 3.9 that $u_\mu \in L^{2^*(s_n+1)/2}$. But

$$\begin{aligned} (2^*/2)(s_n + 1) &= (2^*/2)((r_n/p'_0) - \sigma_1) \\ &\geq (2^*/2)(r_n/p'_0) - (r_n/2)\sigma_1 \\ &= r_n(1 + \varepsilon_0) = r_{n+1} \end{aligned}$$

implies that $u_\mu \in L^{r_{n+1}}$. Hence, we see that $u_\mu \in L^p$ holds for all $p \in [2^*, \infty)$. \square

Lemma 3.11. *For each $\mu \in (0, \mu_a)$, we have $u_\mu \in L^\infty$.*

PROOF: For $t > 0$, we define the function U_t by $U_t = (u_\mu - t)_+$ and the set $A(t)$ by $A(t) = \{x; u_\mu(x) \geq t\}$. Then, we obtain from (3.5) that

$$(3.6) \quad \int \nabla u_\mu \nabla U_t dx \leq \int_{A(t)} q_+ u_\mu^{2+\sigma_1} dx.$$

The constant p_1 may be defined by $p_1 = 2N/(4 - \sigma_1N)$. Since $p_0 > p_1$, we can find a constant $p_2 \in (1, \infty)$ such that $1/p'_0 \cdot 1/p'_2 = 1/p'_1$. Then, the inequality (3.6) implies

$$(3.7) \quad \int |\nabla U_t|^2 dx \leq C(\mu)(\text{meas } A(t))^{1/p'_1}$$

for all $t > 0$, where $C(\mu)$ is defined by

$$(3.8) \quad \begin{aligned} C(\mu) &= \|q_1\|_\infty \left(\int u_\mu^{(2+\sigma_1)p_1} dx \right)^{1/p_1} \\ &\quad + \|q_2\|_{p_0} \left(\int u_\mu^{(2+\sigma_1)p'_0 p_2} dx \right)^{1/(p'_0 p_2)}. \end{aligned}$$

Now, let us assume that $N \geq 3$. Then, it follows from (2.2) and (3.7) that

$$(3.9) \quad \left(\int_{A(t)} (u_\mu - t)^{2^*} dx \right)^{2/2^*} \leq C_0^2 C(\mu) (\text{meas } A(t))^{1/p'_1}.$$

Moreover, for each $h > t$, we have

$$(3.10) \quad \begin{aligned} \left(\int_{A(t)} (u_\mu - t)^{2^*} dx \right)^{2/2^*} &\geq \left(\int_{A(h)} (u_\mu - t)^{2^*} dx \right)^{2/2^*} \\ &\geq (h - t)^2 (\text{meas } A(h))^{2/2^*}. \end{aligned}$$

Combining (3.9) and (3.10) yields

$$\text{meas } A(h) \leq (C_0^2 C(\mu))^{2^*/2} (h - t)^{-2^*} (\text{meas } A(t))^{2^*/2p'_1}$$

for all $h > t > 0$. Since $2^*/(2p'_1) = 1 + (\sigma_1 N)/2(N - 2) > 1$, it follows from Lemma 2.3 that u_μ is essentially bounded. Moreover, for each $t_0 > 0$, we have

$$\|u_\mu\|_\infty \leq d + t_0,$$

where $d = C_0 C(\mu)^{1/2} (\text{meas } A(t_0))^{\sigma_1/4} 2^{1+(2(N-2)/\sigma_1 N)}$. For $t_0 = \|u_\mu\|_2$, it follows that

$$\text{meas } A(t_0) \leq \|u_\mu\|_2^{-2} \int_{A(t_0)} u_\mu^2 dx \leq 1.$$

Hence, we obtain that

$$(3.11) \quad \|u_\mu\|_\infty \leq C_0 C(\mu)^{1/2} 2^{1+(2(N-2)/\sigma_1 N)} + \mu.$$

Finally, we consider the case that $N = 2$. Here, we obtain for all $t > 0$:

$$(3.12) \quad \begin{aligned} \int U_t^2 dx &\leq \int_{A(t)} u_\mu^2 dx \\ &\leq \left(\int_{A(t)} u_\mu^{2p_1} dx \right)^{1/p_1} (\text{meas } A(t))^{1/p'_1}. \end{aligned}$$

Combining (3.7) and (3.12) yields

$$\|U_t\|_{H^1}^2 \leq C^*(\mu) (\text{meas } A(t))^{1/p'_1}$$

for all $t > 0$, where

$$(3.13) \quad C^*(\mu) = C(\mu) + \left(\int u_\mu^{2p_1} dx \right)^{1/p_1}.$$

Hence, (2.1) implies

$$\left(\int_{A(t)} (u_\mu - t)^p dx \right)^{2/p} \leq C_p^2 C^*(\mu) (\text{meas } A(t))^{1/p'_1}$$

for all $t > 0$ and $p \in [2, \infty)$. Then, proceeding as in the case that $N \geq 3$, one can show that

$$\text{meas } A(h) \leq C_p^p C^*(\mu)^{p/2} (h - t)^{-p} (\text{meas } A(t))^{p/(2p'_1)}$$

holds for all $h > t > 0$ and $p \in [2, \infty)$. Hence, according to Lemma 2.3, we see that u is essentially bounded and that

$$(3.14) \quad \|u_\mu\|_\infty \leq C_p C^*(\mu)^{1/2} 2^{(p/(2p'_1))((p/2p'_1)-1)} + \mu$$

if $p > 2p'_1$. □

Lemma 3.12. *For all $p \in [2, \infty)$ we have $\|u_\mu\|_p \rightarrow 0$ as $\mu \rightarrow 0$.*

PROOF: We start with the case that $N = 2$. Then, according to (2.1), we obtain:

$$\|u_\mu\|_p \leq C_p \|u_\mu\|_{H^1} \quad \text{for all } \mu \in (0, \mu_a).$$

Hence, the assertion follows from Proposition 3.7. In case that $N \geq 3$ and $p \in [2, 2^*]$, the assertion is obtained by (2.3) and Proposition 3.7. Now, assume that $N \geq 3$ and that $p \in (2^*, \infty)$. Then, we can find a constant $t > 0$ such that $p = (1 + (t/2))2^*$. Thus, by the Sobolev inequality (2.2), we see that

$$\begin{aligned} (3.15) \quad \|u_\mu\|_p^{2+t} &= \|u_\mu^{1+(t/2)}\|_{2^*}^2 \leq C_0^2 \|\nabla u_\mu^{1+(t/2)}\|_2^2 \\ &= C_0^2 (1 + (t/2))^2 (1+t)^{-1} \int \nabla u_\mu \nabla u_\mu^{1+t} dx. \end{aligned}$$

The right hand side of (3.15) is well defined since u_μ is bounded. From (3.5), we conclude that

$$\begin{aligned} (3.16) \quad \int \nabla u_\mu \nabla u_\mu^{1+t} dx &\leq \int q_+ u_\mu^{2+\sigma_1+t} dx \\ &\leq \|q_1\|_\infty \int u_\mu^{2+\sigma_1+t} dx + \|q_2\|_{p_0} \left(\int u_\mu^{(2+\sigma_1+t)p'_0} dx \right)^{1/p'_0}. \end{aligned}$$

Since

$$\begin{aligned} p'_0 &< 2N/(2(N-2) + \sigma_1 N) < 2N/(2(N-2) + \sigma_1(N-2)) \\ &\leq (2N + tN)/((2 + \sigma_1)(N-2) + t(N-2)) \\ &= (2 + \sigma_1 + t)^{-1} \cdot (2N + tN)/(N-2) \\ &= (2 + \sigma_1 + t)^{-1} p, \end{aligned}$$

we see that there is a constant $\tau \in (0, 1)$ such that

$$(2 + \sigma_1 + t)p'_0 = \tau p + (1 - \tau)2.$$

Hence, by Hölder's inequality, we obtain

$$\left(\int u_\mu^{(2+\sigma_1+t)p'_0} dx \right)^{1/p'_0} \leq \|u_\mu\|_p^{\tau p/p'_0} \|u_\mu\|_2^{2(1-\tau)/p'_0}.$$

Then, using again the fact that $p'_0 < 2N/(2(N-2) + \sigma_1 N)$, it is not difficult to show that $p\tau/p'_0 < 2 + t$.

Quite similarly, one can prove that there exist constants $c_1 \in (0, 2+t)$ and $c_2 > 0$ such that $\int u_\mu^{2+\sigma_1+t} dx \leq \|u_\mu\|_p^{c_1} \|u_\mu\|_2^{c_2}$. Hence, we conclude from (3.15), (3.16) and Young's inequality that $\|u_\mu\|_p \rightarrow 0$ as $\mu \rightarrow 0$. \square

Lemma 3.13. *We have $\|u_\mu\|_\infty \rightarrow 0$ as $\mu \rightarrow 0$.*

PROOF: The constants $C(\mu)$ and $C^*(\mu)$ may be defined as in (3.8) and (3.13). Then, according to Lemma 3.12, it follows that $C(\mu) \rightarrow 0$ and $C^*(\mu) \rightarrow 0$ as $\mu \rightarrow 0$. Hence, the assertion follows from (3.11) and (3.14). \square

PROOF OF COROLLARY 1.2: Suppose that the assumptions of part (a) are fulfilled. Then, according to Lemma 3.5, we see that

$$-\Delta u_\mu + c(x)u_\mu = 0 \text{ holds in } \mathcal{D}'(\mathbb{R}^N),$$

where $c(x) = -q(x)u_\mu^{\sigma_1}(x) + r(x)u_\mu^{\sigma_2}(x) - \lambda(\mu)$. Since $p_0 > N/2$ and $u_\mu \in L^\infty$, we see that $c \in L_{loc}^{p_1}$, where $p_1 = \min(p_0, p)$ satisfies $p_1 > N/2$. Now, the assertion follows from Theorem 7.1 and Corollary 8.1 in [10].

Next, we suppose that the assumptions of the part (b) are fulfilled. Then, it follows from part (a) that u is locally Hölder continuous. Hence, the distribution Δu_μ can be represented by a locally Hölder continuous function. Thus, the assertion of the part (b) follows by a well known result from the regularity theory of elliptic differential equations. \square

PROOF OF COROLLARY 1.3: According to Lemma 3.5, we see that

$$(3.17) \quad -\Delta u_\mu = \lambda(\mu)u_\mu + qu_\mu^{1+\sigma_1} - ru_\mu^{1+\sigma_2} \text{ holds in } \mathcal{D}'(\mathbb{R}^N).$$

Then, it follows from the assumptions and from Lemma 3.10 – Lemma 3.13 that the right hand side of (3.17) defines a function $F_\mu \in L^2$ such that $\|F_\mu\|_2 \rightarrow 0$ as $\mu \rightarrow 0$. Consequently, we see that $u_\mu \in H^2$ and that $\|u_\mu\|_{H^2} \rightarrow 0$ as $\mu \rightarrow 0$. \square

4. Exponential decay.

Lemma 4.1. *Suppose that the functions q and r satisfy the assumptions (A)–(E) and that for $\mu \in (0, \mu_a)$ the function u_μ and the constant $\lambda(\mu)$ are defined as in Lemma 3.4 resp. Lemma 3.5. Moreover, we assume that $\lambda(\mu) < 0$ holds for some $\mu \in (0, \mu_a)$. Then, for each $c \in (0, -\lambda(\mu))$ there exists a constant A_c such that*

$$u_\mu(x) \leq A_c \exp(-(-\lambda(\mu) - c)^{1/2}|x|)$$

holds for almost all $x \in \mathbb{R}^N$.

PROOF: Using the fact that u_μ is bounded, we conclude from (D1) and (E) that there exists a constant $R_c > R_0$ such that

$$(4.1) \quad q_+(x)u_\mu^{\sigma_1}(x) \leq c \text{ holds for almost all } x \in \{y; |y| > R_c\}.$$

The function ψ may be defined by

$$\psi(x) = A_c \exp(-(-\lambda(\mu) - c)^{1/2}|x|) \quad (x \in \mathbb{R}^N).$$

Here, the constant A_c may be chosen such that

$$(4.2) \quad \psi(x) \geq u_\mu(x) \text{ holds for almost all } x \in \{y; |y| \leq R_c\}.$$

Then it follows that $\psi \in H^1$ and that

$$(4.3) \quad \int \nabla \psi \nabla v \, dx \geq (\lambda(\mu) + c) \int \psi v \, dx$$

holds for all nonnegative functions $v \in H^1$.

Inequality (4.2) shows that $(u_\mu - \psi)_+$ is a nonnegative function on H^1 satisfying $(u_\mu - \psi)_+(x) = 0$ for almost all $x \in \{y; |y| \leq R_c\}$. Hence, we obtain from (3.4), (4.1) and (4.3) that

$$\begin{aligned} \|\nabla(u_\mu - \psi)_+\|_2^2 &= \int \nabla(u_\mu - \psi) \nabla(u_\mu - \psi)_+ \, dx \\ &\leq \lambda(\mu) \int u_\mu (u_\mu - \psi)_+ \, dx + c \int u_\mu (u_\mu - \psi)_+ \, dx \\ &\quad - (\lambda(\mu) + c) \int \psi (u_\mu - \psi)_+ \, dx \\ &= (\lambda(\mu) + c) \|(u_\mu - \psi)_+\|_2^2 \leq 0 \end{aligned}$$

and consequently that $u_\mu \leq \psi$. □

Lemma 4.2. *Let q and r satisfy the assumptions (A)–(D) and suppose that $\sigma_2 \leq \sigma_1$. Then $\lambda(\mu) < 0$ holds for all $\mu \in (0, \mu_a)$.*

PROOF: Since $\xi(u_\mu) < 0$, we see that

$$\int r |u_\mu|^{2+\sigma_2} \, dx < -((2 + \sigma_2)/2) \|\nabla u_\mu\|_2^2 + ((2 + \sigma_2)/(2 + \sigma_1)) \int q |u_\mu|^{2+\sigma_1} \, dx$$

and that

$$\lambda(\mu) < \|u_\mu\|_2^{-2} \left(-(\sigma_2/2) \|\nabla u_\mu\|_2^2 + ((\sigma_2 - \sigma_1)/(2 + \sigma_1)) \int q |u_\mu|^{2+\sigma_1} \, dx \right).$$

Then using the fact that

$$\int q |u_\mu|^{2+\sigma_1} \, dx > -(2 + \sigma_1) \xi(u_\mu) > 0,$$

we obtain the assertion. □

Now, we consider the case that $\sigma_1 < \sigma_2$. Since $I(\cdot)$ is a monotone decreasing function on $[0, \mu_a)$, we can find a measurable subset \mathcal{M} of $[0, \mu_a)$ such that $[0, \mu_a) \setminus \mathcal{M}$ has measure zero and $I(\cdot)$ is differentiable on \mathcal{M} (see [4, Theorem 17.12]). Then, we see that

$$(4.4) \quad I'(\mu) \leq 0 \text{ holds for all } \mu \in \mathcal{M}.$$

Lemma 4.3. *The function $I(\cdot)$ is Lipschitz continuous on $[0, \mu_a)$ and for all $\mu \in \mathcal{M}$ we have $I'(\mu) \geq \mu^{-1} \|u_\mu\|_2^2 \lambda(\mu)$.*

PROOF: Let $0 \leq \nu < \mu < \mu_a$. Then, we obtain

$$I(\nu) \leq \xi((\nu/\mu)u_\mu)$$

and therefore that

$$(4.5) \quad \begin{aligned} I(\nu) - I(\mu) &\leq \frac{1}{2}((\nu/\mu)^2 - 1) \int |\nabla u_\mu|^2 dx \\ &\quad - (2 + \sigma_1)^{-1}((\nu/\mu)^{2+\sigma_1} - 1) \int q|u_\mu|^{2+\sigma_1} dx \\ &\quad + (2 + \sigma_2)^{-1}((\nu/\mu)^{2+\sigma_2} - 1) \int r|u_\mu|^{2+\sigma_2} dx. \end{aligned}$$

Thus, (4.5) implies for $\mu \in \mathcal{M}$: $I'(\mu) \geq \mu^{-1} \|u_\mu\|_2^2 \lambda(\mu)$. Moreover, we obtain

$$\begin{aligned} |I(\mu) - I(\nu)| |\mu - \nu|^{-1} &= (I(\nu) - I(\mu))(\mu - \nu)^{-1} \\ &\leq (2 + \sigma_1)^{-1} (1 - (\nu/\mu)^{2+\sigma_1}) (\mu - \nu)^{-1} \int q_+ |u_\mu|^{2+\sigma_1} dx \\ &\leq (1 - (\nu/\mu)) (\mu - \nu)^{-1} \int q_+ |u_\mu|^{2+\sigma_1} dx \\ &= \mu^{-1} \int q_+ |u_\mu|^{2+\sigma_1} dx. \end{aligned}$$

Hence, Lemma 3.1 and Proposition 3.7 show that

$$|I(\mu) - I(\nu)| \leq C(\mu^{1+\alpha} + \mu^{1+\beta}) |\mu - \nu|.$$

□

Lemma 4.4. *There exists a monotone decreasing sequence $(\mu_n) \subset (0, \mu_a)$ such that $\lim_{n \rightarrow \infty} \mu_n = 0$ and $\lambda(\mu_n) < 0$ holds for all n .*

PROOF: Suppose that $\lambda(\mu) \geq 0$ holds for all $\mu \in (0, \mu_a)$. Then, according to Lemma 3.6, we see that $\lambda(\mu) = 0$ holds for all $\mu \in (0, \mu_a)$. Furthermore, (4.4) and Lemma 4.3 would imply that $I'(\mu) = 0$ for all $\mu \in \mathcal{M}$ and consequently that $I(\cdot)$ is constant on $[0, \mu_a)$ (see [4, Theorem 18.15]). In particular, we would obtain that

$$0 = I(0) = I(\min((\mu_a/2), 1)) < 0.$$

Hence, there exists a constant $\mu_1 \in (0, \mu_a)$ such that $\lambda(\mu_1) < 0$. Now, repeating this procedure, we can find a $\mu_2 \in (0, \min(\mu_1, 1/2))$ such that $\lambda(\mu_2) < 0$. Moreover, by induction we can show that for each n there is a constant $\mu_n \in (0, \min(\mu_{n-1}, 1/n))$ so that $\lambda(\mu_n) < 0$. □

Finally, we see that Lemma 4.1 and Lemma 4.2 imply Theorem 1.5 and that Theorem 1.6 is obtained by Lemma 4.1 and Lemma 4.4.

REFERENCES

- [1] Anderson D., *Stability of time — dependent particle solutions in nonlinear field theories II*, J. Math. Phys. **12** (1971), 945–952.
- [2] Berestycki H., Lions P.L., *Nonlinear scalar field equations I: Existence of a ground state*, Arch. Rat. Mech. Anal. **82** (1983), 313–345.
- [3] Gilbarg D., Trudinger N.S., *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, Heidelberg, New York, 1983.
- [4] Hewitt E., Stromberg K., *Real and Abstract Analysis*, Springer-Verlag, Berlin, Heidelberg, New York, 1975.
- [5] Rother W., *Bifurcation of nonlinear elliptic equations on \mathbb{R}^N* , Bull. London Math. Soc. **21** (1989), 567–572.
- [6] ———, *Bifurcation of nonlinear elliptic equations on \mathbb{R}^N with radially symmetric coefficients*, Manuscripta Math. **65** (1989), 413–426.
- [7] ———, *The existence of infinitely many solutions all bifurcating from $\lambda = 0$* , Proc. Royal Soc. Edinburgh **118A** (1991), 295–303.
- [8] ———, *Nonlinear Scalar Field Equations*, Differential and Integral Equations, to appear.
- [9] Ruppen J.-H., *The existence of infinitely bifurcation branches*, Proc. Royal Soc. Edinburgh **101A** (1985), 307–320.
- [10] Stampacchia G., *Le problème de Dirichlet pour les équations elliptique du second ordre à coefficients discontinues*, Annls Inst. Fourier Univ. Grenoble **15** (1965), 189–257.
- [11] ———, *Équations elliptiques du second ordre à coefficients discontinues*, Séminaire de Mathématiques Supérieures, No. 16, Montreal, 1965.
- [12] Strauss W.A., *Existence of solitary waves in higher dimensions*, Commun. Math. Phys. **55** (1977), 149–162.
- [13] Stuart C.A., *Bifurcation from the continuous spectrum in the L^2 — theory of elliptic equations on \mathbb{R}^N* , Recent Methods in Nonlinear Analysis and Applications, Proc. SAFA IV, Liguori, Napoli, 1981, pp. 231–300.
- [14] ———, *Bifurcation for Dirichlet problems without eigenvalues*, Proc. London Math. Soc. (3) **45** (1982), 169–192.
- [15] ———, *Bifurcation from the essential spectrum*, Lecture Notes in Math. **1017** (1983), 575–596.
- [16] ———, *Bifurcation in $L^p(\mathbb{R}^N)$ for a semilinear elliptic equation*, Proc. London Math. Soc. (3) **57** (1988), 511–541.
- [17] ———, *Bifurcation from the essential spectrum for some non-compact non-linearities*, Math. Methods Appl. Sci. **11** (1989), 525–542.
- [18] Zhou H.-S., Zhu X.P., *Bifurcation from the essential spectrum of superlinear elliptic equations*, Appl. Analysis **28** (1988), 51–61.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BAYREUTH, P.O.B. 101251,
W-8580 BAYREUTH, GERMANY

(Received June 12, 1992)