

Oleg Okunev

On analyticity in cosmic spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 34 (1993), No. 1, 185--190

Persistent URL: <http://dml.cz/dmlcz/118568>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1993

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On analyticity in cosmic spaces

OLEG OKUNEV

Abstract. We prove that a cosmic space (= a Tychonoff space with a countable network) is analytic if it is an image of a K -analytic space under a measurable mapping. We also obtain characterizations of analyticity and σ -compactness in cosmic spaces in terms of metrizable continuous images. As an application, we show that if X is a separable metrizable space and Y is its dense subspace then the space of restricted continuous functions $C_p(X | Y)$ is analytic iff it is a $K_{\sigma\delta}$ -space iff X is σ -compact.

Keywords: measurable mapping, cosmic space, analyticity, topology of pointwise convergence

Classification: 54C50, 54C35

All spaces below are assumed to be Tychonoff (= completely regular Hausdorff). We denote by \mathbf{P} the space $\mathbf{N}^{\mathbf{N}}$ where \mathbf{N} is the space of naturals (endowed with the discrete topology). *Analytic* spaces are images of \mathbf{P} under continuous mappings. A space is called *K -analytic* if it is a continuous image of a $K_{\sigma\delta}$ -space (see [9]), that is, a space representable as the intersection of a sequence of σ -compact subspaces of some embracing space. A space X is called *cosmic* if it has a countable network, that is, a countable family of subsets F such that any open set in X is a union of elements of F . Note that in a regular space the family of closures of elements of a network is again a network. A space X is analytic iff it is cosmic and K -analytic (see [9]).

The family of *Baire sets* in a space X is defined as the minimal family of sets that contains all zero-sets of continuous real functions and is closed with respect to countable unions and complements. Of course, in perfectly normal spaces (as all cosmic spaces are), a set is Baire iff it is Borel. A mapping $f : X \rightarrow Y$ is called *measurable* if the preimage of any Baire set in Y is a Baire set in X . One readily checks that if Y is cosmic (in fact, the hereditary Lindelöf property is used) then a mapping $f : X \rightarrow Y$ is measurable as soon as $f^{-1}(U)$ are Baire sets for all elements U of an open subbase for Y .

Clearly, a continuous image of an analytic space is analytic. In 1938 Kuratowski proved the following beautiful theorem (see [6]).

Theorem 0.1. *Let $f : X \rightarrow Y$ be a measurable mapping of a separable metric space X onto a separable metric space Y . If X is analytic, then so is Y .*

In [5] Frolík generalized this theorem to

I wish to express my gratitude to R. Hansell, P. Holický and W. Marciszewski for their interest in this work and helpful comments

Theorem 0.2. *Let $f : X \rightarrow Y$ be a measurable mapping of a K -analytic space X onto a separable metrizable space Y . Then Y is analytic.*

Surprisingly enough, no generalization of this theorem to nonmetrizable spaces Y has been hitherto known (or, at least, no such version was found by the author). On the other hand, analytic spaces need not be metrizable, but are always cosmic, so the class of cosmic spaces looks the most natural domain for considering analyticity. In this paper we generalize Theorem 0.2 to cosmic spaces and infer some corollaries. Some results of this paper were announced in [8].

1. The following lemma is crucial for our proofs. The construction is well-known, only the observation that the ensuing topologies are measurably isomorphic is new.

Lemma 1.1. *Let X be a cosmic space. Then there exist separable metrizable spaces X_u and X_l and continuous bijections $i_u : X_u \rightarrow X$ and $i_l : X \rightarrow X_l$ such that the inverse mappings i_u^{-1} and i_l^{-1} are measurable.*

PROOF: Let $\mathcal{F} = \{F_n : n \in \mathbf{N}\}$ be a countable network for X . Without loss of generality, \mathcal{F} is closed with respect to finite intersections and all elements of \mathcal{F} are closed in X .

Define a new topology, \mathcal{T}_u on the same set X by declaring that the elements of \mathcal{F} are open. Clearly, \mathcal{F} is an open base for this topology, and \mathcal{T}_u is stronger than the original topology of X . Furthermore, \mathcal{F} is a countable base for \mathcal{T}_u consisting of clopen sets, whence \mathcal{T}_u is a completely regular second countable topology. Let X_u be the set X equipped with the topology \mathcal{T}_u ; the identity mapping $i_u : X_u \rightarrow X$ is a continuous bijection. To check the measurability of i_u^{-1} , note that any open set in X_u is the union of a subfamily of \mathcal{F} , hence an F_σ -set in X .

To construct X_l , choose for every $n \in \mathbf{N}$ a continuous function $f_n : X \rightarrow \mathbf{R}$ so that $F_n = f_n^{-1}(0)$ and define the topology \mathcal{T}_l on X as the weakest making all functions $f_n, n \in \mathbf{N}$, continuous. Clearly, \mathcal{T}_l is completely regular and second countable. Let X_l be the set X equipped with the topology \mathcal{T}_l ; the identity mapping of X determines a continuous bijection $i_l : X \rightarrow X_l$. Note that all sets $F_n, n \in \mathbf{N}$, are closed in X_l . Every open set in X is a union of a subfamily of \mathcal{F} , hence an F_σ -set in X_l , whence the mapping i_l^{-1} is measurable. \square

Theorem 1.2. *Let $f : X \rightarrow Y$ be a measurable mapping of a K -analytic space X onto a cosmic space Y . Then Y is analytic.*

PROOF: Define Y_u and $i_u : Y_u \rightarrow Y$ as in Lemma 1.1. The mapping $f \circ i_u^{-1} : X \rightarrow Y_u$ is measurable, and we are now in the conditions of Theorem 0.2, whence Y_u is analytic. Now Y is analytic because it is a continuous image of Y_u . \square

Theorem 1.3. *Let X be a cosmic space. If every metrizable continuous image of X is analytic then X is analytic.*

PROOF: Let X_l and i_l be as in Lemma 1.1. X_l is a metrizable continuous image of X , so X_l is analytic. The mapping $i_l^{-1} : X_l \rightarrow X$ is measurable and onto, so X is analytic by Theorem 1.2. \square

Theorem 1.3 is specific for cosmic spaces. Indeed, \mathbf{N}^{ω_1} is not Lindelöf, not to say K -analytic; nevertheless, all metrizable continuous images of \mathbf{N}^{ω_1} are analytic. To see that note that every metrizable continuous image of \mathbf{N}^{ω_1} is second countable because \mathbf{N}^{ω_1} satisfies c.c.c. Furthermore, every continuous mapping of \mathbf{N}^{ω_1} to a second countable space admits a continuous factorization through a countable face of \mathbf{N}^{ω_1} , homeomorphic to \mathbf{P} , hence has an analytic image. Furthermore, neither the Lindelöf Σ -property nor the hereditary Lindelöf property can replace the cosmicity in Theorem 1.3. To see the first, consider a Lindelöf Σ -space with a single nonisolated point which is not K -analytic [10]. Every disjoint covering of this space by G_δ -sets is countable, hence every metrizable image of this space is countable. The example for the second is the Sorgenfrey line (see [7]). Possibly, the answer to the following question is positive.

Question 1.4. Let X be a perfectly normal Lindelöf Σ -space all metrizable continuous images of which are analytic. Must X be K -analytic?

Theorem 1.5. *Let X be a cosmic space. If every metrizable continuous image of X is σ -compact, then X is σ -compact.*

PROOF: By Theorem 1.3, X is analytic. Now assume for contradiction that X is not σ -compact. Then by a theorem in [3] (a generalization of the Hurewicz theorem to cosmic spaces), X must contain a closed subset F homeomorphic to \mathbf{P} . The space \mathbf{P} is homeomorphic to a subspace of the real line \mathbf{R} (namely, to the space of irrationals), so we can extend the homeomorphism $F \rightarrow \mathbf{P}$ to a continuous function $f_1 : X \rightarrow \mathbf{R}$. Choose a continuous function $f_2 : X \rightarrow \mathbf{R}$ so that $F = f_2^{-1}(0)$ and define $f : X \rightarrow \mathbf{R}^2$ by $f(x) = (f_1(x), f_2(x))$. The set $\{(a, b) \in f(X) : b = 0\}$ is closed in $f(X)$ and is homeomorphic to \mathbf{P} whence $f(X)$ is not σ -compact, a contradiction. \square

2. Denote by $C_p(X)$ the space of all continuous real functions on the space X endowed with the topology of pointwise convergence, that is, the topology of subspace of \mathbf{R}^X endowed with the product topology. Recall that the sets of the form $\{f : f(x) \in U\}$ where $x \in X$ and U is an open subset of \mathbf{R} , constitute an open subbase for this topology. Christensen proved in [4] that if X is a separable metrizable space then $C_p(X)$ is analytic iff X is σ -compact. We extend this result to the following modification of $C_p(X)$.

Let A be a subset of X . Denote $C_p(X | A) = \{f \in C_p(A) : f \text{ admits a continuous extension over } X\}$. In different words, $C_p(X | A)$ is the subspace of $C_p(A)$ consisting of the functions on A which are restrictions to A of continuous functions on X . Note that if A is a dense subspace of X then the restriction mapping $r_A : C_p(X) \rightarrow C_p(X | A)$ is one-to-one.

In our proof below we will also use spaces $C_p(X | A, Y)$ defined as the spaces of all functions of A to Y that admit a continuous extension over X , with the topology of pointwise convergence on A .

Theorem 2.1. *Let X be a separable metrizable space and A a dense subspace of X . Then the following conditions are equivalent.*

- (i) X is σ -compact,

- (ii) $C_p(X | A)$ is a $K_{\sigma\delta}$ -space,
- (iii) $C_p(X | A)$ is analytic.

PROOF: (i) \Rightarrow (ii). Let $X = \bigcup\{X_l : l \in \mathbf{N}\}$ where each $X_l, l \in \mathbf{N}$, is compact. Fix a metric d on X compatible with the topology of X and put

$$M_{kln} = \{f \in [0, 1]^A : |f(x) - f(y)| \leq 1/k \text{ whenever } d(x, z) < 1/n \text{ and } d(y, z) < 1/n \text{ for some } z \in K_l\}, \quad k, l, n \in \mathbf{N}.$$

Claim 1. $C_p(X | A, [0, 1]) = \bigcap_{k \in \mathbf{N}} \bigcap_{l \in \mathbf{N}} \bigcup_{n \in \mathbf{N}} M_{kln}$.

Suppose $f \in \bigcap_{k \in \mathbf{N}} \bigcap_{l \in \mathbf{N}} \bigcup_{n \in \mathbf{N}} M_{kln}$. We need to check that f admits a continuous extension to (or is already continuous at) every point $z \in X$. Find an $l \in \mathbf{N}$ with $z \in X_l$. It suffices to verify that the oscillation of f on A near z is arbitrarily small, which is clear from $f \in \bigcap_{k \in \mathbf{N}} \bigcap_{l \in \mathbf{N}} \bigcup_{n \in \mathbf{N}} M_{kln}$. Thus, $\bigcap_{k \in \mathbf{N}} \bigcap_{l \in \mathbf{N}} \bigcup_{n \in \mathbf{N}} M_{kln} \subset C_p(X | A, [0, 1])$. To check the inverse inclusion, let $f \in C_p(X | A, [0, 1])$ and \hat{f} be the continuous extension of f over X . Fix k and l . Since \hat{f} is continuous and K_l is compact, there exists $n \in \mathbf{N}$ such that $|\hat{f}(x) - \hat{f}(y)| \leq 1/k$ as soon as $d(x, z) < 1/n$ and $d(y, z) < 1/n$ for some $z \in K_l$. Then $f \in M_{kln}$.

Claim 2. The sets M_{kln} are compact.

Indeed, the complement $[0, 1]^A \setminus M_{kln} = \{f \in [0, 1]^A : |f(x) - f(y)| > 1/k \text{ for some } x \in A \text{ and } y \in A \text{ such that } d(x, z) < 1/n \text{ and } d(y, z) < 1/n \text{ for some } z \in K_l\}$. is an open set because the evaluation mapping $f \mapsto f(x), x \in A$, are continuous on $[0, 1]^A$. Thus, M_{kln} is closed in the compact space $[0, 1]^A$, hence compact.

It follows from the claims 1 and 2 that $C_p(X | A, [0, 1])$ is a $K_{\sigma\delta}$ -space. Let $\overline{\mathbf{R}} = \mathbf{R} \cup \{-\infty, +\infty\}$ be the two-point compactification of \mathbf{R} homeomorphic to $[0, 1]$. Clearly, $C_p(X | A, \overline{\mathbf{R}})$ is homeomorphic to $C_p(X | A, [0, 1])$, hence is a $K_{\sigma\delta}$ -space.

Put $S = \bigcap_{l \in \mathbf{N}} \bigcup_{n \in \mathbf{N}} \{f \in \overline{\mathbf{R}}^A : |f(x)| \leq n \text{ for all } x \in A \cap K_l\}$. Clearly, S is an $F_{\sigma\delta}$ -set in $\overline{\mathbf{R}}^A$, hence a $K_{\sigma\delta}$ -space. Furthermore, $C_p(X | A) \subset S \subset \overline{\mathbf{R}}^A$, whence $C_p(X | A) = S \cap C_p(X | A, \overline{\mathbf{R}})$; $C_p(X | A)$ is an intersection of two $K_{\sigma\delta}$ -spaces, hence is a $K_{\sigma\delta}$ -space.

(ii) \Rightarrow (iii). The space A is second countable, hence $C_p(A)$, and consequently $C_p(X | A)$ are cosmic [1]. Every $K_{\sigma\delta}$ -space is K -analytic, and every K -analytic cosmic space is analytic [9], so we are done.

(iii) \Rightarrow (i). Let $e : C_p(X | A) \rightarrow C_p(X)$ be the “extension mapping” inverse to the restriction mapping $r_A : C_p(X) \rightarrow C_p(X | A)$. The mapping e is measurable. Indeed, $C_p(X)$ is cosmic [1], hence hereditary Lindelöf, so it suffices to check that $e^{-1}(V)$ are Baire sets for all elements of some subbase for $C_p(X)$. Let $V = \{f \in C_p(X) : f(x) \in U\}$ where $x \in X$ and U is open in \mathbf{R} ; as we already noted, the sets of this form constitute a subbase for $C_p(X)$. Choose a sequence $\{x_n : n \in \mathbf{N}\}$ of points of A converging to x ; we have

$$V = \bigcup_{n \in \mathbf{N}} \bigcap_{k > n} \{f \in C_p(X) : f(x_k) \in U\},$$

and

$$e^{-1}(V) = r_A(V) = \bigcup_{n \in \mathbf{N}} \bigcap_{k > n} \{f \in C_p(X | A) : f(x_k) \in U\},$$

a $G_{\delta\sigma}$ -set in $C_p(X | A)$. Thus, e is measurable, and $C_p(X)$ is analytic by Theorem 1.2. By theorem of Christensen [4], X is σ -compact. \square

Corollary 2.2. *If A is a dense subspace of \mathbf{P} then $C_p(\mathbf{P} | A)$ is a coanalytic nonanalytic subspace of \mathbf{R}^A .*

PROOF: Only coanalyticity now requires a proof. Put

$$\omega(f, U) = \sup\{|f(x) - f(y)| : x, y \in U\}$$

for $f \in \mathbf{R}^A$ and $U \subset A$. Clearly

$$C_p(\mathbf{P} | A) = \{f \in \mathbf{R}^A : \text{for any } x \in \mathbf{P} \text{ and } n \in \mathbf{N} \text{ there is a neighbourhood } U \text{ of } x \text{ with } \omega(f, U \cap A) \leq 1/n\}.$$

For $x \in \mathbf{P} = \mathbf{N}^{\mathbf{N}}$ denote by $x | n$ the sequence of the first n coordinates of x . Clearly, the sets of the form $V(x | k) = \{y \in \mathbf{P} : y | k = x | k\}$, $k \in \mathbf{N}$, form a base for \mathbf{P} at x . Hence $C_p(\mathbf{P} | A) = \{f \in \mathbf{R}^A : \text{for any } x \in \mathbf{P} \text{ and } n \in \mathbf{N} \text{ there exists } k \in \mathbf{N} \text{ with } \omega(f, V(x | k) \cap A) \leq 1/n\} = \bigcap_{x \in \mathbf{P}} \bigcap_{n \in \mathbf{N}} \bigcup_{k \in \mathbf{N}} F(x | k, n)$ where $F(x | k, n) = \{f \in \mathbf{R}^A : \omega(f, V(x | k) \cap A) \leq 1/n\}$. A straightforward check shows that the sets $F(x | k, n)$ are closed, hence $C_p(\mathbf{P} | A)$ is coanalytic. \square

Corollary 2.2 answers a question of T. Dobrowolski. W. Marciszewski communicated at the Seventh Prague Topological Symposium, August 1991, that this corollary had been also obtained by A. Krawczyk, who used quite a different approach. It is interesting to note that one can derive the nonanalyticity of $C_p(\mathbf{P} | A)$ also from Theorem 1.3 if he uses the facts that every mapping of $C_p(X)$ to a second countable space admits a factorization through $C_p(X | A)$ for some countable and, without loss of generality, dense A [2] and that all $C_p(\mathbf{P} | A)$ with dense countable A are homeomorphic (because the pairs (\mathbf{P}, A) are).

REFERENCES

- [1] Arhangel'skiĭ A.V., *Some topological spaces that arise in functional analysis*, Russ. Math. Surveys **31** (1976), 14–30.
- [2] ———, *Factorization theorems and function spaces: stability and monolithicity*, Soviet. Math. Doklady **26** (1982), 177–181.
- [3] ———, *Hurewicz spaces, analytic sets, and fan tightness of function spaces*, Soviet. Math. Doklady **33** (1986), 396–399.
- [4] Christensen J.P.R., *Topology and Borel Structure*, North Holland, Amsterdam, 1974.
- [5] Frolík Z., *A measurable map with analytic domain and metrizable range is quotient*, Bull. Amer. Math. Soc. **76** (1970), 1112–1117.
- [6] Kuratowski K., *Topology*, Vol. 1, Academic Press, N.Y.-London, 1966.
- [7] Motorov D.B., *Metrizable images of the arrow (Sorgenfrey line)*, Moscow Univ. Math. Bull. **39** (1984), 48–50.

- [8] Okunev O., *On analyticity in non-metrizable spaces*, Abstracts of the VII Prague Topol. Symp., p. 101.
- [9] Rogers C.A., Jayne J.E., and al., *Analytic Sets*, Academic Press, London, 1980.
- [10] Talagrand M., *A new countably determined Banach space*, Israel J. Math. **47** (1984), 75–80.

TVER STATE UNIVERSITY, PEDAGOGICAL DEPARTMENT, BOX 0236, 170002 TVER, RUSSIA

(Received May 26, 1992)