

Ghassan Alherk; Henryk Hudzik

Copies of l^1 and c_0 in Musielak-Orlicz sequence spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 35 (1994), No. 1, 9--19

Persistent URL: <http://dml.cz/dmlcz/118636>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Copies of l^1 and c_0 in Musielak-Orlicz sequence spaces

GHASSAN ALHERK, HENRYK HUDZIK

Abstract. Criteria in order that a Musielak-Orlicz sequence space l^Φ contains an isomorphic as well as an isomorphically isometric copy of l^1 are given. Moreover, it is proved that if $\Phi = (\Phi_i)$, where Φ_i are defined on a Banach space, X does not satisfy the δ_2^0 -condition, then the Musielak-Orlicz sequence space $l^\Phi(X)$ of X -valued sequences contains an almost isometric copy of c_0 . In the case of $X = \mathbb{R}$ it is proved also that if l^Φ contains an isomorphic copy of c_0 , then Φ does not satisfy the δ_2^0 -condition. These results extend some results of [A] and [H2] to Musielak-Orlicz sequence spaces.

Keywords: Musielak-Orlicz sequence space, copy of l^1 , copy of c_0

Classification: 46B20, 46B25, 46E30

0. Introduction

Two Banach spaces X, Y are said to be $(1 + \varepsilon)$ -isometric provided there exists a linear isomorphism $P : X \xrightarrow{\text{onto}} Y$ such that $\|P\| \|P^{-1}\| \leq 1 + \varepsilon$. It is easy to see that P is a $(1 + \varepsilon)$ -isometry if

$$\|x\|_X \leq \|Px\|_Y \leq (1 + \varepsilon)\|x\|_X$$

for any $x \in X$. We say a Banach space X contains an almost isometric (isomorphic) copy of Y if for any $\varepsilon > 0$ (for some $\varepsilon > 0$) there exists a subspace Z in X such that Z, Y are $(1 + \varepsilon)$ -isomorphic. We say a Banach space X contains an isomorphically isometric (shortly isometric) copy of Y if there exist a subspace Z of X and a linear isomorphism P from Z onto Y such $\|Px\|_Y = \|x\|_X$ for any $x \in Z$.

In the sequel X denotes a real Banach space and $\mathbb{N}, \mathbb{R}, \mathbb{R}_+$ and \mathbb{R}_+^c stand for the set of natural numbers, the set of reals, the set of nonnegative reals, and for $\mathbb{R}_+ \cup +\infty$, respectively. A map $\Phi : X \rightarrow \mathbb{R}_+^c$ is said to be an Orlicz function if it is convex, even, vanishing and continuous at 0, lower semicontinuous on the whole X and

$$(*) \quad \inf\{\Phi(x) : \|x\| = r\} \rightarrow \infty \quad \text{as} \quad r \rightarrow \infty.$$

We define a Musielak-Orlicz function Φ to be a sequence (Φ_i) of Orlicz functions (we write then $\Phi = (\Phi_i)$). Given a Banach space X , we denote by $l^\circ(X)$ the real space of all X -valued sequences $x = (x_n)$. We write shortly l° instead of $l^\circ(\mathbb{R})$.

Given an arbitrary Musielak-Orlicz function $\Phi = (\Phi_i)$ we define a functional $I_\Phi : l^o(X) \rightarrow \mathbb{R}_+^e$ by

$$I_\Phi(x) = \sum_{i=1}^{\infty} \Phi_i(x_i),$$

which is even and convex, $I_\Phi(0) = 0$ and for any $x \in l^o(X)$ the condition $I_\Phi(\lambda x) = 0$ for any $\lambda > 0$ yields $x = 0$.

Musielak-Orlicz space $l^\Phi(X)$ generated by a Musielak-Orlicz function Φ is defined as the set of all $x \in l^o(X)$ such that $I_\Phi(\lambda x) < \infty$ for some $\lambda > 0$ (cf. [T] and in the scalar case also [KR], [L], [M] and [RR]).

The subspace $h^\Phi(X)$ of $l^\Phi(X)$ is defined to be the closure in $l^\Phi(X)$ of the space $h(X)$ of all x in $l^o(X)$ which have only finite number of coordinates different from 0. The space $l^\Phi(X)$ can be equipped with the norm

$$\|x\|_\Phi = \inf\{\varepsilon > 0 : I_\Phi(x/\varepsilon) \leq 1\},$$

called the Luxemburg norm (cf. [M] and in the case of Orlicz spaces also [KR], [L] and [RR]). The space $h^\Phi(X)$ will be considered with the norm $\|\cdot\|_\Phi$ induced from $l^\Phi(X)$. $l^\Phi(X)$ and $h^\Phi(X)$ equipped with the norm $\|\cdot\|_\Phi$ are Banach spaces (cf. [T]).

We say that a Musielak-Orlicz function $\Phi = (\Phi_i)$ satisfies the δ_2^o -condition (we write $\Phi \in \delta_2^o$) if there are positive constants k and a , a sequence (c_i) with $c_i \in \mathbb{R}_+^e$ such that $\sum_{i=j}^{\infty} c_i < \infty$ for some $j \in \mathbb{N}$ and

$$\Phi_i(2x) \leq \Phi_i(x) + c_i$$

for any $i \in \mathbb{N}$ and $x \in X$ satisfying $\Phi_i(x) \leq a$.

If $\Phi = (\Phi_i)$ satisfies the δ_2^o -condition with $j = 1$ we say that Φ satisfies the δ_2 -condition. Of course, for any Musielak-Orlicz function $\Phi = (\Phi_i)$ with finite-valued Φ_i for any $i \in \mathbb{N}$ the δ_2 -condition is equivalent to the δ_2^o -condition (cf. [DH] and [K]).

Let us define for any Musielak-Orlicz function $\Phi = (\Phi_i)$ the sequence $\lambda = (\lambda_i)$ in \mathbb{R}_+ , where

$$\lambda_i = \sup\{u \in \mathbb{R}_+ : \Phi_i \text{ is linear on } [0, u] \text{ and } \Phi_i(u) \leq 1\}$$

for $i = 1, 2, \dots$.

1. Results

We start with the following theorem:

Theorem 1. *Let $\Phi = (\Phi_i)$ be a Musielak-Orlicz function with finite-valued Φ_i defined on \mathbb{R} for any $i \in \mathbb{N}$. Then $l^\Phi = (l^\Phi, \|\cdot\|_\Phi)$ contains an isometric copy of l^1 if and only if:*

- (i) Φ does not satisfy the δ_2 -condition.
- (ii) $\sum_{i=1}^{\infty} \Phi_i(\lambda_i) = \infty$.

PROOF: Sufficiency. Under our assumptions concerning Φ , the conditions δ_2 and δ_2^o are equivalent. Therefore, if Φ satisfies condition (i), then $\Phi \notin \delta_2^o$. This yields that l^Φ contains an isometric copy of l^∞ (cf. [K]) and so also an isometric copy of l^1 .

Assume now that Φ satisfies the condition (ii). Define i_1 to be the largest natural number satisfying

$$\sum_{i=1}^{i_1} \Phi_i(\lambda_i) \leq 1.$$

Then

$$\sum_{i=1}^{i_1+1} \Phi_i(\lambda_i) > 1.$$

There is a number $\alpha_i \in [0, \lambda_i)$ such that

$$\sum_{i=1}^{i_1} \Phi_i(\lambda_i) + \Phi_{i_1+1}(\alpha_1) = 1.$$

We have

$$\sum_{i=i_1+2}^{\infty} \Phi_i(\lambda_i) = \infty.$$

Define $i_2 \geq i_1 + 2$ to be the largest natural number such that

$$\sum_{i=i_1+2}^{i_2} \Phi_i(\lambda_i) \leq 1.$$

Then

$$\sum_{i=i_1+2}^{i_2+1} \Phi_i(\lambda_i) > 1.$$

There is a number $\alpha_2 \in [0, \lambda_{i_2+1})$ such that

$$\sum_{i=i_1+2}^{i_2} \Phi_i(\lambda_i) + \Phi_{i_2+1}(\alpha_2) = 1.$$

Proceeding in such a way by induction we find sequences (i_k) of natural numbers and (α_k) of numbers from the intervals $[0, \lambda_{i_k+1})$ such that

$$\sum_{i=i_{k-1}+2}^{i_k} \Phi_i(\lambda_i) + \Phi_{i_k+1}(\alpha_k) = 1$$

for $k = 1, 2, \dots$, where $i_0 = -1$ by definition. Denote

$$A_k = \{i_{k-1} + 1, \dots, i_k, i_k + 1\}, \quad k = 1, 2, \dots$$

The sets A_k are pairwise disjoint and $\bigcup_{k=1}^{\infty} A_k = \mathbb{N}$. Define a new sequence $d = (d_i)$ by

$$d_i = \begin{cases} \lambda_i & \text{if } i \in A_k \setminus \{i_k + 1\} \\ \alpha_k & \text{if } i = i_k + 1 \end{cases}$$

for $k = 1, 2, \dots$. Define now

$$f_k = \sum_{i \in A_k} d_i e_i,$$

where e_i is the i -th unit vector, i.e. $e_i = (0, \dots, 0, 1, 0, \dots)$ with 1 on the i -th place. We have

$$I_{\Phi}(f_k) = 1 \quad \text{for } k = 1, 2, \dots$$

We have also $f_k \perp f_l$ (i.e. the sequences f_k and f_l have disjoint supports) if $k \neq l$. Moreover, the coordinates of $f_k (k = 1, 2, \dots)$ belong to the intervals on which the respective Orlicz functions Φ_i are linear. Define an operator P from l^1 into l^{Φ} by the formula

$$Px = \sum_{k=1}^{\infty} x_k f_k \quad (\forall x = (x_k) \in l^1).$$

It is obvious that P is linear. Moreover

$$\begin{aligned} I_{\Phi} \left(\frac{Px}{\|x\|_{l^1}} \right) &= \sum_{k=1}^{\infty} I_{\Phi} \left(\frac{x_k f_k}{\|x\|_{l^1}} \right) \\ &= \sum_{k=1}^{\infty} \frac{|x_k|}{\|x\|_{l^1}} I_{\Phi}(f_k) = \sum_{k=1}^{\infty} \frac{|x_k|}{\|x\|_{l^1}} = 1. \end{aligned}$$

Hence

$$\left\| \frac{Px}{\|x\|_{l^1}} \right\|_{\Phi} = 1, \text{ i.e. } \|Px\|_{\Phi} = \|x\|_{l^1}.$$

This means that P is an isometry between l^1 and a closed subspace $P(l^1)$ of l^{Φ} .

Necessity. Assume that Φ satisfies the δ_2 -condition and $\sum_{i=1}^{\infty} (\lambda_i) < +\infty$. Then there is $n \in \mathbb{N}$, $n \geq 2$, such that $\sum_{i=1}^{\infty} \Phi_i(\lambda_i) \leq n$. We will prove that l^{Φ} is non- l_n^1 , i.e. for any elements x^1, \dots, x^n from the unit sphere $\mathcal{S}(l^{\Phi})$ of l^{Φ} there holds

$$\left\| \frac{1}{n} (x^1 \pm x^2 \pm \dots \pm x^n) \right\| < 1$$

for some choice of signs, which yields that l^1 can not be isometrically embedded into l^Φ .

Take arbitrary $x^1, \dots, x^n \in \mathcal{S}(l^\Phi)$. Then $I_\Phi(x^1) = \dots = I_\Phi(x^k) = 1$ for $i = 1, 2, \dots, n$. Define the set

$$A = \{i \in \mathbb{N} : \sum_{k=1}^n \Phi_i(x_i^k) > \Phi_i(\lambda_i)\}.$$

We will prove that for any $i \in A$

$$(1) \quad \Phi_i \left(\frac{(x_i^1 \pm \dots \pm x_i^n)}{n} \right) < \frac{1}{n} \sum_{k=1}^n \Phi_i(x_i^k)$$

for some choice of signs ± 1 , dividing the proof into two cases.

I. $\max\{|x_i^k| : k = 1, \dots, n\} \leq \lambda_i$ and $i \in A$. Then at least two numbers among $\Phi_i(x_i^k), k = 1, \dots, n$, must be positive. Assume that this is not true, i.e. that there is only one positive number $\Phi_i(x_i^j)$ among these numbers. Then

$$\sum_{k=1}^n \Phi_i(x_i^k) = \Phi_i(x_i^j) \leq \Phi_i(\lambda_i),$$

which contradicts the fact that $i \in A$. Therefore,

$$\max\{\Phi_i(x_i^k) : k = 1, \dots, n\} < \sum_{k=1}^n \Phi_i(x_i^k).$$

It is evident that

$$(2) \quad |x_i^1 \pm \dots \pm x_i^n| \leq \max\{|x_i^k| : k = 1, \dots, n\}$$

for some choice of signs ± 1 , whence

$$\begin{aligned} \Phi_i \left(\frac{1}{n} (x_i^1 \pm \dots \pm x_i^n) \right) &\leq \Phi \left(\frac{1}{n} \max_k |x_i^k| \right) \\ &= \frac{1}{n} \max_k \Phi_i(x_i^k) < \frac{1}{n} \sum_{k=1}^n \Phi_i(x_i^k). \end{aligned}$$

This means that inequality (1) holds true in case I.

II. $i \in A$ and $\max\{|x_i^k| : k = 1, \dots, n\} > \lambda_i$. Applying (2), we get for a choice of signs ± 1

$$\begin{aligned} \Phi_i \left(\frac{1}{n} (x_i^1 \pm \dots \pm x_i^n) \right) &\leq \Phi_i \left(\frac{1}{n} \max_k |x_i^k| \right) < \frac{1}{n} \Phi_i(\max_k |x_i^k|) \\ &= \frac{1}{n} \max_k \Phi_i(x_i^k) \leq \frac{1}{n} \sum_{k=1}^n \Phi_i(x_i^k). \end{aligned}$$

Combining both cases I and II we get inequality (1) for some choice of signs ± 1 . For the remaining $2^{n-1} - 1$ choices of signs ± 1 , we have by the convexity of Φ ,

$$(3) \quad \Phi_i \left(\frac{1}{n}(x_i^1 \pm \dots \pm x_i^n) \right) \leq \frac{1}{n} \sum_{k=1}^n \Phi_i(x_i^k) \quad (\forall i \in A).$$

Combining (1) and (3), we get

$$\sum_{\pm 1} \Phi_i \left(\frac{1}{n}(x_i^1 \pm \dots \pm x_i^n) \right) < \frac{2^{n-1}}{n} \sum_{k=1}^n \Phi_i(x_i^k) \quad (\forall i \in A).$$

Summing up both-side of the last inequality over $i \in A$, we have

$$\sum_{\pm 1} I_{\Phi} \left(\frac{1}{n}(x^1 \pm \dots \pm x^n) \chi_A \right) < \frac{2^{n-1}}{n} \sum_{k=1}^n I_{\Phi}(x^k \chi_A),$$

where χ_A denotes the characteristic function of A .

Hence it follows that

$$\begin{aligned} & 2^{n-1} - \sum_{\pm 1} I_{\Phi} \left(\frac{1}{n}(x^1 \pm \dots \pm x^n) \right) \\ &= \frac{2^{n-1}}{n} \sum_{k=1}^n I_{\Phi}(x^k) - \sum_{\pm 1} I_{\Phi} \left(\frac{1}{n}(x^1 \pm \dots \pm x^n) \right) \\ &\geq \frac{2^{n-1}}{n} \sum_{k=1}^n I_{\Phi}(x^k \chi_A) - \sum_{\pm 1} I_{\Phi} \left(\frac{1}{n}(x^1 \pm \dots \pm x^n) \chi_A \right) > 0, \end{aligned}$$

i.e.

$$\sum_{\pm 1} I_{\Phi} \left(\frac{1}{n}(x^1 \pm \dots \pm x^n) \right) < 2^{n-1}.$$

Therefore

$$I_{\Phi} \left(\frac{1}{n}(x^1 \pm \dots \pm x^n) \right) < 1$$

for at least one choice of signs ± 1 . Since Φ_i are finite-valued by the assumption and $\Phi \in \delta_2$, we get

$$\left\| \frac{1}{n}(x^1 \pm \dots \pm x^n) \right\| < 1$$

for at least one choice of signs (cf. [DH] and [K]), i.e. l^{Φ} is non- l_n^1 . This means that l^1 cannot be embedded isometrically into l^{Φ} , and the proof is finished. \square

Theorem 2. *Let $\Phi = (\Phi_i)$ be an arbitrary Musielak-Orlicz function defined on \mathbb{R} . Then $l^\Phi = (l^\Phi, \|\cdot\|_\Phi)$ contains an isomorphic copy of l^1 if and only if Φ or Φ^* (= the complementary function to Φ in the sense of Young) does not satisfy the δ_2^o -condition, i.e. if and only if l^Φ is not reflexive.*

PROOF: Sufficiency. If $\Phi \notin \delta_2^o$, then l^Φ contains an isometric copy of l^∞ (cf. [K]) and so of l^1 as well. Assume now that $\Phi \in \delta_2^o$ but $\Phi^* \notin \delta_2^o$. Then the dual of l^Φ is isomorphically isometric to l^{Φ^*} equipped with the Orlicz norm $\|\cdot\|_{\Phi^*}$ (cf. [M] and in the case of Orlicz spaces also [KR], [L] and [RR]). Therefore l^{Φ^*} contains an isomorphic copy of l^∞ (cf. again [K]), whence it follows that l^Φ contains an isomorphic copy of l^1 .

Necessity. Assume that both functions Φ and Φ^* satisfy the δ_2^o -condition. Then l^Φ is reflexive and so l^1 can not be embedded isomorphically into l^Φ as a nonreflexive space. \square

Theorem 3. *Let X be an arbitrary Banach space and $\Phi = (\Phi_i)$ be a Musielak-Orlicz function defined on X . Then:*

- (i) *if Φ does not satisfy the δ_2^o -condition, then $h^\Phi(X) = (h^\Phi(X), \|\cdot\|_\Phi)$ contains an almost isometric copy of c_o ;*
- (ii) *if $X = \mathbb{R}$ and $(h^\Phi = h^\Phi, \|\cdot\|_\Phi)$ contains an isomorphic copy of c_o , then Φ does not satisfy the δ_2^o -condition.*

PROOF: (i). Let

$$c_i^{k,\varepsilon} = \sup\{\Phi_i((1+\varepsilon)x) - 2^{k+1}\Phi_i(x) : \Phi_i(x) \leq 2^{-k-1}\} (\forall i, k \in \mathbb{N}, \varepsilon > 0).$$

We have that $\Phi \in \delta_2^o$ if and only if there exists $\varepsilon > 0$ and $m, k \in \mathbb{N}$ such that

$$\sum_{i=m}^{\infty} c_i^{k,\varepsilon} < \infty \quad (\text{cf. [DH] and [H1]}).$$

Define

$$\alpha_i^{k,\varepsilon} = \sup\{\Phi_i((1+\varepsilon)x) : \Phi_i(x) \leq 2^{-k-1} \quad \text{and} \quad \Phi_i((1+\varepsilon)x) - 2^{k+1}\Phi_i(x) \geq 0\}.$$

Since $\Phi_i(0) = 0 < 2^{-k-1}$, so 0 belongs to the set of these x over which the supremum in the definition of $c_i^{k,\varepsilon}$ is taken. Moreover,

$$\Phi((1+\varepsilon)0) - 2^{k+1}\Phi(0) = 0.$$

Hence it follows that $c_i^{k,\varepsilon} \geq 0$. Therefore, we can restrict ourselves in the definition of $c_i^{k,\varepsilon}$ to these x for which $\Phi_i((1+\varepsilon)x) - 2^{k+1}\Phi_i(x) \geq 0$. Hence we get $c_i^{k,\varepsilon} \leq d_i^{k,\varepsilon}$ for every $i, k \in \mathbb{N}$ and $\varepsilon > 0$. We have by the assumption that $\Phi \notin \delta_2^o$. Hence it follows that

$$\sum_{i=m}^{\infty} d_i^{k,\varepsilon} = \infty \quad (\forall m, k \in \mathbb{N}, \varepsilon > 0),$$

so we have among others $\sum_{i=1}^{\infty} d_i^{k,\varepsilon} = \infty$. Define i_1 as the largest natural number such that

$$\sum_{i=1}^{i_1} d_i^{k,\varepsilon} \leq 1,$$

whenever $d_1^{k,\varepsilon} \leq 1$ and $i_1 = 0$ whenever $d_1^{k,\varepsilon} > 1$. Then

$$\sum_{i=1}^{i_1+1} d_i^{k,\varepsilon} > 1.$$

Define in the first step $N_1 = \{1, \dots, i_1 + 1\}$. Define i_2 as the largest natural number such that

$$\sum_{i=i_1+2}^{i_2} d_i^{k,\varepsilon} \leq 1$$

if $d_{i_1+2}^{k,\varepsilon} \leq 1$ and $i_2 = i_1 + 2$ if $d_{i_1+2}^{k,\varepsilon} > 1$.

Then

$$\sum_{i=i_1+2}^{i_2+1} d_i^{k,\varepsilon} > 1.$$

Put $N_2 = \{i_1 + 2, \dots, i_2 + 1\}$. Proceeding in such a way by induction we can find a sequence (i_k) of nonnegative integers such that the sequence of pairwise disjoint sets (N_k) in \mathbb{N} defined by

$$N_k = \{i_{k-1} + 2, \dots, i_k + 1\}, i_0 := -1 \quad (k = 1, 2, \dots)$$

satisfies

$$(4) \quad \sum_{i \in N_k \setminus \{i_k + 1\}} d_i^{k,\varepsilon} \leq 1,$$

$$(5) \quad \sum_{i \in N_k} d_i^{k,\varepsilon} > 1.$$

In view of the definition of $d_i^{k,\varepsilon}$ and inequality (5), it follows that for any $\varepsilon > 0$ and $k \in \mathbb{N}$ there are $x_i \in X (x \in N_k)$ such that

$$(6) \quad \sum_{i \in N_k} \Phi_i((1 + \varepsilon)x_i) > 1,$$

$$(7) \quad \Phi_i(x_i) \leq 2^{-k-1} \quad \text{and} \quad \Phi_i((1 + \varepsilon)x_i) \geq 2^{k+1}\Phi_i(x_i) \quad (\forall i \in N_k).$$

Applying (6) and (7) we get

$$(8) \quad \sum_{i \in N_k} \Phi_i(x_i) \leq 2^{-k-1} \sum_{i \in N_k \setminus \{i_k+1\}} d_i^{k,\varepsilon} + 2^{-k-1} \leq 2^{-k-1} + 2^{-k-1} = 2^{-k}.$$

Define $y_k = \sum_{i=N_k} x_i e_i$. Then y_k have pairwise disjoint supports. In virtue of (6) and (8) we have

$$(9) \quad I_\Phi(y_k) = \sum_{i \in N_k} \Phi_i(x_i) \leq 2^{-k} \quad (\forall k \in \mathbb{N}),$$

$$(10) \quad I_\Phi((1+\varepsilon)y_k) = \sum_{i \in N_k} \Phi_i((1+\varepsilon)x_i) > 1 \quad (\forall k \in \mathbb{N}).$$

Define now an operator $P_1 : c_o \rightarrow h^\Phi(X)$ by

$$P_1 u = \sum_{k=1}^{\infty} u_k y_k \quad (\forall u = (u_k) \in c_o).$$

It is obvious that P_1 is linear. We will prove now that $P_1 u \in h^\Phi(X)$ for any $u \in c_o$. We need to prove that there is a sequence (y^l) in $l^o(X)$ such that y^l has only finite number of coordinates different from zero and $\|P_1 u - y^l\|_\Phi \rightarrow 0$ as $l \rightarrow \infty$, i.e.

$$(11) \quad I_\Phi(\lambda(P_1 u - y^l)) \rightarrow 0 \quad \text{as } l \rightarrow \infty \quad (\forall \lambda > 0).$$

Take arbitrary $\lambda, \varepsilon > 0$ and choose $k_o \in \mathbb{N}$ such that $\sum_{k=k_o}^{\infty} 2^{-k} < \varepsilon$. Define

$$y^l = \sum_{k=1}^l u_k y_k \quad (l = 1, 2, \dots).$$

Obviously any y^l has only finite number of coordinates different from zero. Let $l_o \geq k_o$ be such that $|u_k| \lambda \leq 1$ for any $k \geq l_o$. Such a number l_o exists because $u = (u_k) \in c_o$. We have for any $l \geq l_o$

$$\begin{aligned} I_\Phi(\lambda(P_1 u - y^l)) &= I_\Phi\left(\sum_{k=l+1}^{\infty} \lambda u_k y_k\right) \leq I_\Phi\left(\sum_{k=l+1}^{\infty} y_k\right) \\ &= \sum_{k=l+1}^{\infty} I_\Phi(y_k) \leq \sum_{k=l+1}^{\infty} 2^{-k} < \varepsilon, \end{aligned}$$

i.e. condition (11) holds. This means that $P_1 u \in h^\Phi$ for any $u \in c_o$. Applying (9), we get for any $u \in c_o$,

$$\begin{aligned} I_\Phi(P_1 u / \|u\|_\infty) &= \sum_{k=1}^{\infty} I_\Phi(u_k y_k / \|u\|_\infty) \\ &\leq \sum_{k=1}^{\infty} I_\Phi(y_k) \leq \sum_{k=1}^{\infty} 2^{-k} = 1, \end{aligned}$$

whence it follows that

$$(12) \quad \|P_1 u\|_\Phi \leq \|u\|_\infty \quad (\forall u \in c_o).$$

Let $k_o \in N$ be such that $|u_{k_o}| = \|u\|_\infty$. Then, in view of (10), we get

$$\begin{aligned} I_\Phi((1 + \varepsilon)\|u\|_\infty^{-1} P_1 u) &\geq I_\Phi((1 + \varepsilon)\|u\|_\infty^{-1} u_{k_o} y_{k_o}) \\ &= I_\Phi((1 + \varepsilon)y_{k_o}) > 1, \end{aligned}$$

whence it follows that

$$\|P_1 u\|_\Phi \geq \frac{1}{1 + \varepsilon} \|u\|_\infty \quad (\forall u \in c_o).$$

Defining now

$$Pu = (1 + \varepsilon)P_1 u \quad (\forall u \in c_o),$$

we get a linear operator from c_o into $h^\Phi(X)$ satisfying

$$\|u\|_\infty \leq \|Pu\|_\Phi \leq (1 + \varepsilon)\|u\|_\infty \quad (\forall u \in c_o),$$

which means that P is a $(1 + \varepsilon)$ -isometry. Since $\varepsilon > 0$ is arbitrary this means that c_o is embedded almost isometrically into h^Φ and the proof of statement (i) is finished.

(ii). Assume that $\Phi \in \delta_2^o$ and $X = \mathbb{R}$. Then $h^\Phi = l^\Phi$ and l^Φ is the dual space of h^{Φ^*} , where Φ^* is the Orlicz function complementary in the sense of Young to Φ (cf. [HY]). Assume that h^Φ contains an isomorphic copy of c_o . Then it contains (as a dual space) a copy of l^∞ (cf. [BP]). But this contradicts the fact that the norm $\|\cdot\|_\Phi$ is order continuous in h^Φ . This contradiction finishes the proof of statement (ii) and so of Theorem 3 as well. \square

Recall that a Banach lattice E is said to be a KB-space whenever every increasing bounded in the norm sequence of nonnegative elements in E is norm convergent to an element of E (cf. [AB] and [KA]).

Remark. It is known (cf. [AB, p. 227]) that if E is a Banach lattice that is not KB-space, then c_o is embeddable in E and conversely. The space h^Φ is a Banach lattice that is KB-space if and only if $\Phi \in \delta_2^o$. Therefore, h^Φ contains an isomorphic copy of c_o if and only if $\Phi \notin \delta_2^o$. It is worth to notice that if X is an arbitrary Banach space, then $h^\Phi(X)$ need not be a Banach lattice. However, in one direction an analogous result (cf. Theorem 3) still holds true.

REFERENCES

- [A] Alherk G., *On the non- $l_n^{(1)}$ and locally uniformly non- $l_n^{(1)}$ properties and l^1 copies in Musielak-Orlicz space*, Comment. Math. Univ. Carolinae **31** (1990), 435–443.
- [AB] Aliprantis C.D., Burkinshaw O., *Positive operators*, Pure and Applied Math., Academic Press, Inc., 1985.
- [BP] Bessaga C., Pelczyński A., *On bases and unconditional convergence of series in Banach spaces*, Studia Math. **17** (1958), 151–164.
- [DH] Denker M., Hudzik H., *Uniformly non- $l_n^{(1)}$ Musielak-Orlicz sequence spaces*, Proc. Indian Acad. Sci. **101** (1991), 71–86.
- [H1] Hudzik H., *On some equivalent conditions in Musielak-Orlicz spaces*, Comment. Math. (Prace Matem.) **24** (1984), 57–64.
- [H2] ———, *Orlicz spaces containing a copy of l^1* , Math. Japonica **34** (1989), 747–759.
- [HY] Hudzik H., Ye Y., *Support functionals and smoothness in Musielak-Orlicz sequence spaces endowed with the Luxemburg norm*, Comment. Math. Univ. Carolinae **31** (1990), 661–684.
- [K] Kamińska A., *Flat Orlicz-Musielak sequence spaces*, Bull. Acad. Polon. Sci. Math. **30** (1982), 347–352.
- [KA] Kantorovich L.V., Akilov G.P., *Functional Analysis* (in Russian), Nauka, Moscow, 1977.
- [KR] Krasnoselskii M.A., Rutickii Ya.B., *Convex functions and Orlicz spaces*, Groningen, 1961 (translation).
- [L] Luxemburg W.A.J., *Banach function spaces*, Thesis, Delft, 1955.
- [M] Musielak J., *Orlicz spaces and modular spaces*, Lecture Notes in Math. 1034, Springer-Verlag, 1983.
- [RR] Rao M.M., Ren Z.D., *Theory of Orlicz spaces*, Pure and Applied Mathematics, Marcel Dekker, 1991.
- [T] Turett B., *Fenchel-Orlicz spaces*, Dissertationes Math. **181** (1980), 1–60.

UNIVERSITY OF ALEPPO, ALEPPO, SYRIA

ADAM MICKIEWICZ UNIVERSITY, POZNAŃ, POLAND

(Received February 16, 1993)