

Josef Daněček

On  $\mathcal{L}_{loc}^{2,n}$ -regularity for the gradient of a weak solution to nonlinear elliptic systems

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 37 (1996), No. 3, 523--536

Persistent URL: <http://dml.cz/dmlcz/118860>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## On $\mathcal{L}_{loc}^{2,n}$ -regularity for the gradient of a weak solution to nonlinear elliptic systems

JOSEF DANĚČEK

*Abstract.* Interior  $\mathcal{L}_{loc}^{2,n}$ -regularity for the gradient of a weak solution to nonlinear second order elliptic systems is investigated.

*Keywords:* nonlinear elliptic system, regularity, Campanato-Morrey spaces

*Classification:* 35J60

### 1. Introduction

In this paper we consider the problem of the regularity of the first derivatives of weak solutions to a nonlinear elliptic system

$$(1) \quad -D_\alpha (A_i^\alpha (Du)) = 0, \quad (i = 1, \dots, N)$$

in a bounded open set  $\Omega \subset R^n$ . Throughout the whole text we use the summation convention over repeated indexes.

If  $n \geq 3$ , it is known that  $Du$  may not be continuous. Examples are provided by nonregular solutions of elliptic systems presented by Nečas in [8]. Campanato in [2] proved that  $Du \in \mathcal{L}_{loc}^{2,\lambda}(\Omega, R^N)$  with  $\lambda(n) < n$ , and  $u \in C_{loc}^{0,\alpha}(\Omega, R^N)$  for some  $\alpha < 1$  if  $n = 3, 4$ . In this paper we give sufficient condition on  $\mathcal{L}_{loc}^{2,n}$ -regularity for the gradient of a weak solution to (1). Recall that if  $Du \in \mathcal{L}_{loc}^{2,n}$ , then  $u$  is locally Zygmund continuous.

### 2. Preliminaries

Let  $\Omega$  be a bounded open set in  $R^n$  with points  $x = (x_1, \dots, x_n)$ ,  $n \geq 3$ . The notation  $\Omega_0 \Subset \Omega$  means that the closure of  $\Omega_0$  is contained in  $\Omega$ , i.e.  $\overline{\Omega_0} \subset \Omega$ . For the sake of simplicity we denote by  $|\cdot|$  and  $(\cdot, \cdot)$  the norm and scalar product in  $R^n$ ,  $R^N$  and  $R^{nN}$ . If  $x \in R^n$  and  $r$  is a positive real number, we set  $B(x, r) = \{y \in R^n : |y - x| < r\}$ , i.e. the open ball in  $R^n$ ,  $\Omega(x, r) = B(x, r) \cap \Omega$ . By  $\mu(\Omega(x, r))$  we denote the  $n$ -dimensional Lebesgue measure of  $\Omega(x, r)$ . A bounded domain  $\Omega \subset R^n$  is said to be of type  $\mathcal{A}$  if there exists a constant  $\mathcal{A} > 0$  such that for every  $x \in \overline{\Omega}$  and all  $0 < r < \text{diam } \Omega$  it holds  $\mu(\Omega(x, r)) \geq \mathcal{A}r^n$ .

Let  $u: \Omega \rightarrow R^N$ ,  $N \geq 1$ ,  $u(x) = (u^1(x), \dots, u^N(x))$  be a vector-valued function and  $Du = (D_1u, \dots, D_nu)$ ,  $D_\alpha = \partial/\partial x_\alpha$ .

By  $u_{x,r} = \mu^{-1}(\Omega(x,r)) \int_{\Omega(x,r)} u(y) dy = \int_{\Omega(x,r)} u(y) dy$  we denote mean value of  $u$  over the set  $\Omega(x,r)$  provided that  $u \in L^1(\Omega, R^N)$ . Besides usually used spaces as  $C_0^\infty(\Omega, R^N)$ , the Hölder spaces  $C^{0,\alpha}(\overline{\Omega}, R^N)$  and the Sobolev spaces  $H^{k,p}(\Omega, R^N)$ ,  $H_{loc}^{k,p}(\Omega, R^N)$ ,  $H_0^{k,p}(\Omega, R^N)$  (see e.g. [1], [6], [7] for definitions and basic properties) we use the following Campanato and Morrey spaces.

**Definition 1** (Campanato and Morrey spaces). Let  $\lambda \in [0, n]$ ,  $q \in [1, \infty)$ . The Morrey space  $L^{q,\lambda}(\Omega, R^N)$  is the subspace of such functions  $u \in L^q(\Omega, R^N)$  for which  $\|u\|_{L^{q,\lambda}(\Omega, R^N)}^q = \sup\{r^{-\lambda} \int_{\Omega(x,r)} |u(y)|^q dy : r > 0, x \in \Omega\}$  is finite.

Let  $\lambda \in [0, n + q]$ ,  $q \in [1, \infty)$ . The Campanato spaces  $\mathcal{L}^{q,\lambda}(\Omega, R^N)$  and  $\mathcal{L}_1^{q,\lambda}(\Omega, R^N)$  are subspaces of such functions  $u \in L^q(\Omega, R^N)$  for which  $[u]_{\mathcal{L}^{q,\lambda}(\Omega, R^N)}^q = \sup\{r^{-\lambda} \int_{\Omega(x,r)} |u(y) - u_{x,r}|^q dy : r > 0, x \in \Omega\}$  is finite and  $[u]_{\mathcal{L}_1^{q,\lambda}(\Omega, R^N)}^q = \sup\{\inf\{r^{-\lambda} \int_{\Omega(x,r)} |u(y) - P(y)|^q dy : P \in \mathcal{P}_1\} : r > 0, x \in \Omega\}$  is finite. Here  $\mathcal{P}_1$  is the set of all polynomials in  $n$  variables and of degree  $\leq 1$ . Let us denote  $\|u\|_{L^{q,\lambda}}$ ,  $\|u\|_{\mathcal{L}^{q,\lambda}} = \|u\|_{L^q} + [u]_{\mathcal{L}^{q,\lambda}}$  and  $\|u\|_{\mathcal{L}_1^{q,\lambda}} = \|u\|_{L^q(\Omega, R^N)} + [u]_{\mathcal{L}_1^{q,\lambda}}$ .

*Remark 1.* It is worth to recall a trivial however important property saying that  $\int_{\Omega} |u - u_{\Omega}|^2 dx = \min\{\int_{\Omega} |u - c|^2 dx : c \in R^N\}$  for every  $u \in L^2(\Omega, R^N)$ .

**Definition 2.** The Zygmund class  $\Lambda^1(\overline{\Omega}, R^N)$  is the subspace of such functions  $u \in C^0(\overline{\Omega}, R^N)$  for which  $[u]_{\Lambda^1(\overline{\Omega}, R^N)} = \sup\{|u(x) + u(y) - 2u((x+y)/2)| : |x - y| : x, y, (x+y)/2 \in \overline{\Omega}\}$  is finite.

For more details see [1], [4], [6], [7]. In particular, we will use the following result.

**Proposition 1.** *Let  $\Omega$  be of type  $\mathcal{A}$  and  $1 \leq q < \infty$ . Then it holds*

- (a)  $L^{q,\lambda}(\Omega, R^N)$ ,  $\mathcal{L}^{q,\lambda}(\Omega, R^N)$  and  $\mathcal{L}_1^{q,\lambda}(\Omega, R^N)$  equipped with norms  $\|u\|_{L^{q,\lambda}}$ ,  $\|u\|_{\mathcal{L}^{q,\lambda}}$  and  $\|u\|_{\mathcal{L}_1^{q,\lambda}}$  are Banach spaces.
- (b)  $\mathcal{L}^{q,\lambda}(\Omega, R^N)$  is isomorphic to the  $C^{0,(\lambda-n)/q}(\overline{\Omega}, R^N)$  if  $n < \lambda \leq n + q$ ,
- (c)  $L^{q,n}(\Omega, R^N)$  is isomorphic to the  $L^\infty(\Omega, R^N) \subsetneq \mathcal{L}^{q,n}(\Omega, R^N)$ ,
- (d)  $\mathcal{L}_1^{2,n+2}(\Omega, R^N)$  is isomorphic to the  $\Lambda^1(\overline{\Omega}, R^N)$ ,
- (e)  $C^{0,1}(\overline{\Omega}, R^N) \subsetneq \Lambda^1(\overline{\Omega}, R^N) \subsetneq \bigcap_{0 < \alpha < 1} C^{0,\alpha}(\overline{\Omega}, R^N)$ .

Further, we suppose

- (i) there is an  $M > 0$  such that for every  $p \in R^{nN}$

$$(2) \quad |A_i^\alpha(p)| \leq M(1 + |p|),$$

(ii)  $A_i^\alpha(p)$  are differentiable functions on  $R^{nN}$  with the bounded and continuous derivatives, i.e.

$$(3) \quad \left| \frac{\partial A_i^\alpha}{\partial p_\beta^j}(p) \right| \leq M \quad \text{for every } p \in R^{nN},$$

(iii) the strong ellipticity condition, i.e. there exists  $\nu > 0$  such that for every  $p, \xi \in R^{nN}$

$$(4) \quad \frac{\partial A_i^\alpha}{\partial p_\beta^j}(p) \xi_\alpha^i \xi_\beta^j \geq \nu |\xi|^2.$$

From (ii) it follows (see [3, p.169]) the existence of a real function  $\omega(s)$  defined on  $[0, \infty)$ , which is nonnegative, bounded, nondecreasing, concave,  $\omega(0) = 0$  (moreover,  $\omega$  is right continuous at 0 for uniformly continuous  $\partial A_i^\alpha / \partial p_\beta^j$ ) and such that for all  $p, q \in R^{nN}$

$$(5) \quad \left| \frac{\partial A_i^\alpha}{\partial p_\beta^j}(p) - \frac{\partial A_i^\alpha}{\partial p_\beta^j}(q) \right| \leq \omega(|p - q|^2).$$

By a weak solution of (1) we mean a function  $u \in H^{1,2}(\Omega, R^N)$  satisfying

$$(6) \quad \int_{\Omega} A_i^\alpha(Du) D_\alpha \varphi^i dx = 0$$

for every  $\varphi \in H_0^{1,2}(\Omega, R^N)$ .

We will also consider the pair of complementary Young functions

$$(7) \quad \Phi(t) = t \ln_+ at \quad \text{for } t \geq 0, \quad \Psi(t) = \begin{cases} t/a & \text{for } 0 \leq t < 1, \\ e^{t-1}/a & \text{for } t \geq 1, \end{cases}$$

where  $a > 0$  is a constant,  $\ln_+ at = 0$  for  $0 \leq t < 1/a$  and  $\ln_+ at = \ln at$  for  $t \geq 1/a$ . Recall Young's inequality

$$ts \leq \Phi(t) + \Psi(s), \quad t, s \geq 0.$$

For our consideration we also need to introduce quasiconvex functions.

**Definition 3** ([5, p.4]). A function  $G: [0, \infty) \rightarrow R$  is said to be quasiconvex (quasiconcave) on  $[0, \infty)$  if there exist a convex (concave) function  $g(\tilde{g})$  and a constant  $c > 0$  ( $\tilde{c} > 0$ ) such that

$$g(t) \leq G(t) \leq cg(ct), \quad (\tilde{g}(t) \leq G(t) \leq \tilde{c}\tilde{g}(\tilde{c}t)) \quad \text{for } t \geq 0.$$

Next, we will need the following properties of quasiconvex functions:

**Lemma 1** ([5, p. 4]). *Let us consider three statements:*

- (a)  $G(t)$  is quasiconvex (quasiconcave) on  $[0, \infty)$ ;
- (b) for all  $t_1, t_2 \in [0, \infty)$  and all  $\lambda \in (0, 1)$

$$G(\lambda t_1 + (1 - \lambda)t_2) \leq k_1(\lambda G(k_1 t_1) + (1 - \lambda)G(k_1 t_2))$$

$$\left( \lambda G(t_1) + (1 - \lambda)G(t_2) \leq l_1 G(l_1(\lambda t_1 + (1 - \lambda)t_2)) \right);$$

- (c) there exists a constant  $k_2$  ( $l_2$ ) such that for all  $u \in L^2_{loc}(\Omega, R^N)$  and all balls  $B(x, r) \subset \Omega$

$$\begin{aligned} G\left(\int_{B(x,r)} |u|^2 dy\right) &\leq k_2 \int_{B(x,r)} G(k_2 |u|^2) dy, \\ \left(\int_{B(x,r)} G(|u|^2) dy \leq l_2 G\left(l_2 \int_{B(x,r)} |u|^2 dy\right)\right). \end{aligned}$$

Then (a)  $\Rightarrow$  (b)  $\Rightarrow$  (c).

**Proposition 2.** *For all  $u, v \in L^2_{loc}(\Omega, R^N)$ , all balls  $B(x, r) \subset \Omega$  and an arbitrary nondecreasing quasiconvex function  $G$  on  $[0, \infty)$  we have*

(a)

$$\int_{B(x,r)} G(|u + v|^2) dy \leq \frac{k_1}{2} \left( \int_{B(x,r)} G(4k_1 |u|^2) dy + \int_{B(x,r)} G(4k_1 |v|^2) dy \right),$$

(b)

$$\int_{B(x,r)} G(|u - u_{x,r}|^2) dy \leq c_1 \int_{B(x,r)} G(c_2 |u - c|^2) dy,$$

where  $c_1 = \max\{k_1/2, k_2\}$ ,  $c_2 = \max\{4k_1, 4k_1 k_2\}$  and  $c \in R$  is arbitrary.

PROOF: (a) It follows from Lemma 1 (b).

(b) From (a) we get

$$\int_{B_r} G(|u - u_{x,r}|^2) dy \leq \frac{k_1}{2} \left( \int_{B_r} G(4k_1 |u - c|^2) dy + \int_{B_r} G(4k_1 |c - u_{x,r}|^2) dy \right).$$

Now, by means of Hölder's inequality and Lemma 1 (c)

$$\begin{aligned} \int_{B_r} G(4k_1 |c - u_{x,r}|^2) dy &= \mu(B_r) G(4k_1 |c - u_{x,r}|^2) \\ &= \mu(B_r) G\left(4k_1 \left|c - \int_{B_r} u(y) dy\right|^2\right) = \mu(B_r) G\left(\frac{4k_1}{\mu^2(B_r)} \left|\int_{B_r} (u(y) - c) dy\right|^2\right) \\ &\leq \mu(B_r) G\left(\int_{B_r} 4k_1 |u(y) - c|^2 dy\right) \leq k_2 \int_{B_r} G(4k_1 k_2 |u(y) - c|^2) dy \end{aligned}$$

and the result follows easily. □

**Lemma 2** ([9, p. 37]). *Let  $\varphi: [0, \infty] \rightarrow [0, \infty]$  be a monotone function which is absolutely continuous on every closed interval of finite length. If  $v \geq 0$  is measurable and  $E(t) = \{x \in R^n : v(x) > t\}$ , then*

$$\int_{R^n} \varphi \circ v dx = \int_0^\infty \mu(E(t)) \varphi'(t) dt.$$

**Proposition 3.** *Let  $v \in L_{loc}^2(\Omega, R^m)$ ,  $B(x, \sigma) \subset \Omega$ ,  $a > 0$  and  $s \in [1, \infty)$  be arbitrary. If the inequality*

$$\int_{B(x, \tau\sigma)} |v - v_{x, \tau\sigma}|^2 dy \leq \int_{B(x, \sigma)} |v - v_{x, \sigma}|^2 dy$$

holds for some  $\tau \in (0, 1)$ , then there exists a constant  $b$  such that

$$\int_{B(x, \tau\sigma)} \ln_+^s (a|v - v_{x, \tau\sigma}|^2) dy \leq b \int_{B(x, \sigma)} \ln_+^s (a|v - v_{x, \sigma}|^2) dy.$$

For the constant  $b$  we have the following estimate

$$b \leq h \left( \int_{B(x, \sigma)} |v - v_{x, \sigma}|^2 dy \right) \left( \int_{B(x, \sigma)} \ln_+^s (a|v - v_{x, \sigma}|^2) dy \right)^{-1},$$

where  $h(t) = (s/e(s-1))^{s/(s-1)} at$ ,  $t \in [0, e^{s/(s-1)}/a]$  and  $\ln^{s/(s-1)}(at)$ ,  $t \in (e^{s/(s-1)}/a, \infty)$ .

PROOF: We set  $E_{\tau\sigma}(t) = \{y \in B(x, \tau\sigma) : |v - v_{x,\tau\sigma}|^2 > t\}$  for  $t \geq 0$  and  $0 < \tau \leq 1$ . From Lemma 2 and by means of integration by parts we get

$$\begin{aligned} \int_{B_{\tau\sigma}} \ln_+^s(a|v - v_{\tau\sigma}|^2) dy &= \frac{s}{\mu(B_{\tau\sigma})} \int_{1/a}^\infty \mu(E_{\tau\sigma}(t)) \frac{\ln^{s-1}(at)}{t} dt \\ &= \frac{s}{\mu(B_{\tau\sigma})} \left[ \frac{\ln^{s-1}(at)}{t} \int_0^t \mu(E_{\tau\sigma}(\lambda)) d\lambda \right]_{1/a}^\infty \\ &\quad + \frac{s}{\mu(B_{\tau\sigma})} \int_{1/a}^\infty \left( \int_0^t \mu(E_{\tau\sigma}(\lambda)) d\lambda \right) \frac{\ln^{s-1}(at) - (s-1)\ln^{s-2}(at)}{t^2} dt. \end{aligned}$$

For the sake of simplicity we put  $V_r = \int_{B(x,r)} |v - v_{x,r}|^2 dy$ . The first integral is zero and on the second integral we can use the mean value theorem for the integrals and we have for some  $1/a < \xi_{\tau\sigma}$ ,  $\xi_\sigma < \infty$ ,

$$\begin{aligned} \int_{B_{\tau\sigma}} \ln_+^s(a|v - v_{\tau\sigma}|^2) dy &= sV_{\tau\sigma} \int_{\xi_{\tau\sigma}}^\infty \frac{\ln^{s-1}(at) - (s-1)\ln^{s-2}(at)}{t^2} dt \\ &= \frac{s \ln^{s-1}(a\xi_{\tau\sigma})}{\xi_{\tau\sigma}} V_{\tau\sigma} = \frac{\xi_\sigma \ln^{s-1}(a\xi_{\tau\sigma})}{\xi_{\tau\sigma} \ln^{s-1}(a\xi_\sigma)} \frac{V_{\tau\sigma}}{V_\sigma} \int_{B_\sigma} \ln_+^s(a|v - v_{x,\sigma}|^2) dy \\ &= b(\tau) \int_{B_\sigma} \ln_+^s(a|v - v_{x,\sigma}|^2) dy. \end{aligned}$$

Now the result follows from Lemma 1 (c). □

**3. The result**

For  $x \in \Omega$ ,  $r > 0$  we set  $U_r = U(x, r) = \int_{\Omega(x,r)} |Du - (Du)_{x,r}|^2 dy$ ,  $d_x = \text{dist}(x, \partial\Omega)$  and  $\alpha_n = \mu(B(0, 1))$ . We define  $S_0 = \{x \in \Omega : \overline{\lim}_{r \rightarrow 0+} U(x, r) > 0\}$ .

*Remark 2.* Let  $u$  be a solution of (1). It is well known (see [9, pp. 75, 122]) that  $\lim_{r \rightarrow 0+} U(x, r) = 0$  for all  $x \in \Omega \setminus E$  where  $n - 2 + \beta$  dimensional Hausdorff measure  $H^{n-2+\beta}(E) = 0$  for every  $\beta > 0$ .

Now we can formulate the main theorem.

**Theorem.** Let  $u \in H^{1,2}(\Omega, R^N)$  be a weak solution to the nonlinear system (1) under the hypotheses (i), (ii), (iii). Let  $x \in \mathcal{S}_0$  be arbitrary and suppose that there exists  $d \in (0, d_x/2)$  such that

$$(8) \quad \frac{Kl_2\omega^2}{\nu^2} \left( b \int_{B(x,2d)} \ln_+^{q/(q-1)} \left( \frac{4l_2\omega^2 |Du - (Du)_{x,2d}|^2}{CU_{2d}} \right) dy \right)^{1-1/q} < \frac{1}{4} \tau^n,$$

where  $K = c(n, N, q) (M/\nu)^8$ ,  $\tau = (2^{n+5}A)^{-1/2}$ ,  $l_2, A$  are the constants from Lemma 1 (c), Lemma 3,  $\omega = \omega(2^n l_2 U_{2d})$ ,  $\omega$  is from (5),  $C = 2^{n-8} \nu^2 \tau^n / \alpha_n A$  and  $b$  is the constant from Proposition 3 for the case  $a = 1/CU_{2d}$ ,  $\sigma = 2d$ ,  $v = 2\sqrt{l_2}\omega Du$ ,  $s = q/(q - 1)$  where  $q \in (1, n/(n - 2)]$ . Then there exists a ball  $B(x, r_x) \subset \Omega$  such that  $Du \in \mathcal{L}^{2,n}(B(x, r_x), R^{nN})$  and

$$(9) \quad [Du]_{\mathcal{L}^{2,n}(B(x,r_x),R^{nN})}^2 \leq \max\{2^n(4A\tau^{-n} + 1)U_{2d}, \mu^{-1}(B_{2d}) \int_{\Omega} |Du - (Du)_{\Omega}|^2 dx\}.$$

**Proposition 4.** Set  $\omega_{\infty} = \lim_{t \rightarrow \infty} \omega(t)$ ,  $V_1 = c_1 (M/\nu)^{3n+8} (\omega_{\infty}/\nu)^2$  and  $V_2 = c_2 (M/\nu)^{3n+6} (\omega_{\infty}/\nu)^2$ . If

$$(10) \quad V_2 \leq e^q \ \& \ q^{q-1} V_1 V_2^{1-1/q} < 1 \quad \text{or} \quad V_2 > e^q \ \& \ V_1 \ln^{q-1} V_2 < 1,$$

then condition (8) holds for every  $x \in \mathcal{S}_0$ . Here  $q \in (1, n/(n - 2)]$ ,  $c_1 = c_1(n, N, q)$  and  $c_2 = c_2(n, N)$ .

PROOF: Let  $x \in \mathcal{S}_0$  and  $d \in (0, d_x/2)$  be arbitrary such that  $U(x, 2d) > 0$ . From Proposition 3 it follows that the left hand side of (8) is equal or less than  $Kl_2\omega_{\infty}^2 h^{1-1/q}(4\omega_{\infty}^2 U_{2d})/\nu^2$ . From the definition of the function  $h(t)$  and assumption (10) it follows that (8) is satisfied. □

**Example.** We can consider the system (1) for  $n = 3, N = 2$  where  $A_i^{\alpha}(p) = (a \delta_{ij} \delta_{\alpha\beta} + b \delta_{i\alpha} \delta_{j\beta} \arctan |p|^2 / 2\pi) p_{\beta}^j$ ,  $a, b$  are constants,  $0 < b/6 < a$ . We have

$$\frac{\partial A_i^{\alpha}}{\partial p_{\beta}^j}(p) \xi_{\alpha}^i \xi_{\beta}^j \geq (a - b/6) |\xi|^2, \quad \forall \xi, p \in R^6,$$

$\omega_{\infty} \leq b$  and  $|\partial A_i^{\alpha} / \partial p_{\beta}^j(p)| \leq M = a + b$ . Setting  $P = b/a$  we get that  $V_1 < 4c_1 P^2 (1 + P)^{3n+8} / (1 - P/6)^{3n+10}$ ,  $V_2 < 4c_2 P^2 (1 + P)^{3n+6} / (1 - P/6)^{3n+8}$  and it is not difficult to see that (10) is satisfied for  $P$  sufficiently small.



**Corollary 1.** *Let  $\Omega_0 \Subset \Omega$  be arbitrary and of type  $\mathcal{A}$  and the assumptions of Theorem be satisfied for every  $x \in \overline{\Omega}_0 \cap \mathcal{S}_0$ . Then there are constants  $\mathcal{U}, d_0, r_0 > 0$  such that  $Du \in \mathcal{L}^{2,n}(\Omega_0, R^{nN})$  and the following estimate*

$$(11) \quad \begin{aligned} [Du]_{\mathcal{L}^{2,n}(\Omega_0, R^{nN})}^2 &\leq \max\{2^n(4A\tau^{-n} + 1)\mathcal{U}, \\ &\mu^{-1}(B_{2d_0}) \int_{\Omega} |Du - (Du)_{\Omega}|^2 dx, \\ &(Ar_0^n)^{-1} \int_{\Omega_0} |Du - (Du)_{\Omega_0}|^2 dx \} \end{aligned}$$

holds.

PROOF: From Remark 2, Theorem and the definition of the set  $\mathcal{S}_0$  it follows that for every  $x \in \overline{\Omega}_0$  there exists  $B(x, r_x) \subset \Omega$  such that  $Du \in \mathcal{L}^{2,n}(B(x, r_x), R^{nN})$ . As  $\overline{\Omega}_0$  is the compact set and the system balls  $\{B(x, r_x)\}$  covers of  $\overline{\Omega}_0$  we can choose a finite subcover  $\{B(x_j, r_{x_j})\}_{j=1}^m$ . If we set  $\mathcal{U} = \max\{U(x_j, 2d_{x_j}) : 1 \leq j \leq m\}$ ,  $r_0 = \min\{r_{x_j} : 1 \leq j \leq m\}$  and  $d_0 = \min\{d_{x_j} : 1 \leq j \leq m\}$ , then the estimate follows from Remark 1.  $\square$

**Corollary 2.** *Let the assumptions of Corollary 1 be satisfied. Then  $u \in A^1(\overline{\Omega}_0, R^N)$ .*

PROOF: It follows from Proposition 1 (d), Poincaré’s inequality and Corollary 1.  $\square$

#### 4. Lemmas

The statement of the following lemma is well known (see e.g. [1], [3], [7], [8]).

**Lemma 3.** *Let  $v \in H^{1,2}(\Omega, R^N)$  be a weak solution to the system (1) satisfying (i), (ii) and (iii), where  $\partial A_i^\alpha / \partial p_j^\beta$  are the constants. Then there exists a constant  $A = c(n, N)(M/\nu)^6$  such that for every  $x \in \Omega$  and  $0 < \sigma \leq R \leq \text{dist}(x, \partial\Omega)$  the following estimate holds*

$$\int_{B(x,\sigma)} |Dv(y) - (Dv)_{x,\sigma}|^2 dy \leq A \left(\frac{\sigma}{R}\right)^{n+2} \int_{B(x,R)} |Dv(y) - (Dv)_{x,R}|^2 dy.$$

The following lemma is possible to derive by the difference quotient method (see e.g. [1], [3], [7], [8]).

**Lemma 4.** *Let  $u \in H^{1,2}(\Omega, R^N)$  be a weak solution to the system (1) satisfying (i), (ii) and (iii). Then  $u \in H_{loc}^{2,2}(\Omega, R^N)$  and for all  $x \in \Omega$ ,  $0 < \sigma < \varrho \leq \text{dist}(x, \partial\Omega)$  we have*

$$\int_{B(x,\sigma)} |D^2u|^2 dy \leq \frac{6n(M/\nu)^2}{(\varrho - \sigma)^2} \int_{B(x,\varrho)} |Du - (Du)_{x,\varrho}|^2 dy.$$

**Lemma 5** ([6]). *Let  $1 \leq p, q < \infty, p^{-1} - q^{-1} \leq n^{-1}, R > 0, x \in \mathbb{R}^n$ . Then for  $u \in H^{1,p}(B(x, R), \mathbb{R}^N)$  we have*

$$\left( \int_{B(x,R)} |u(y)|^q dy \right)^{1/q} \leq cR^{1+n/q-n/p} \left( R^{-p} \int_{B(x,R)} |u(y)|^p dy + \int_{B(x,R)} |Du(y)|^p dy \right)^{1/p},$$

where  $c = c(n, N, p, q)$  is a constant independent of  $x, R$  and  $u$ .

**Lemma 6.** *Let  $u \in H^{1,2}(\Omega, \mathbb{R}^N)$  be a weak solution to (1) satisfying (i), (ii) and (iii). Then for every ball  $B(x, 2R) \subset \Omega$  and an arbitrary constant  $a > 0$  we have*

$$\begin{aligned} & \int_{B(x,R)} |Du - (Du)_{x,R}|^2 \ln_+(a |Du - (Du)_{x,R}|^2) dy \\ & \leq c \left( \frac{M}{\nu} \right)^2 \left( \int_{B(x,2R)} \ln_+^{q/(q-1)}(4a |Du - (Du)_{x,2R}|^2) dy \right)^{1-1/q} \\ & \qquad \qquad \qquad \int_{B(x,2R)} |Du - (Du)_{x,2R}|^2 dy, \end{aligned}$$

where  $1 < q \leq n/(n - 2)$  and  $c = c(n, N, q)$ .

PROOF: Let  $x \in \Omega$  and  $0 \leq R \leq \frac{1}{4} \text{dist}(x, \partial\Omega)$ . We denote  $B_R = B(x, R)$  for simplicity. From Lemma 4 it follows that  $Du \in H_{loc}^{1,2}(\Omega, \mathbb{R}^N)$ . By means of Sobolev's imbedding theorem  $H^{1,2}(B_R, \mathbb{R}^N) \hookrightarrow L^s(B_R, \mathbb{R}^N)$ , where  $B_R \subset \Omega$  be arbitrary and  $1 \leq s \leq 2n/(n - 2)$ . From this we obtain by Proposition 2 (b) and Lemma 5

$$\begin{aligned} & \int_{B_R} |Du - (Du)_R|^2 \ln_+(a |Du - (Du)_R|^2) dy \\ & \leq 4 \int_{B_R} |Du - (Du)_{2R}|^2 \ln_+(4a |Du - (Du)_{2R}|^2) dy \\ & \leq 4 \left( \int_{B_R} |Du - (Du)_{2R}|^{2q} dy \right)^{1/q} \left( \int_{B_R} \ln_+^{q/(q-1)}(4a |Du - (Du)_{2R}|^2) dy \right)^{1-1/q} \\ & \leq cR^{n(1/q-1)+2} \left( R^{-2} \int_{B_R} |Du - (Du)_{2R}|^2 + \int_{B_R} |D^2u|^2 dy \right) \times \end{aligned}$$

$$\begin{aligned} & \times \left( \int_{B_R} \ln_+^{q/(q-1)} (4a |Du - (Du)_{2R}|^2) dy \right)^{1-1/q} \\ & \leq c \left( \frac{M}{\nu} \right)^2 R^{-n(1-1/q)} \int_{B_{2R}} |Du - (Du)_{2R}|^2 dy \times \\ & \quad \times \left( \int_{B_R} \ln_+^{q/(q-1)} (4a |Du - (Du)_{2R}|^2) dy \right)^{1-1/q} \end{aligned}$$

and we finally obtain the result. □

**5. Proof of Theorem**

Set  $A_{ij}^{\alpha\beta}(\zeta) = \partial A_i^\alpha / \partial p_\beta^j(\zeta)$ ,  $A_{ij,0}^{\alpha\beta} = A_{ij}^{\alpha\beta}((Du)_R)$ ,

$$\tilde{A}_{ij}^{\alpha\beta} = \int_0^1 A_{ij}^{\alpha\beta}((Du)_R + t(Du - (Du)_R)) dt,$$

$B_R = B(x, R)$  and  $U_R = U(x, R)$  for simplicity. Then the system (1) can be rewritten as

$$-D_\alpha \left( A_{ij,0}^{\alpha\beta} D_\beta u^j \right) = -D_\alpha \left( \left( A_{ij,0}^{\alpha\beta} - \tilde{A}_{ij}^{\alpha\beta} \right) \left( D_\beta u^j - (D_\beta u^j)_R \right) \right).$$

Split  $u$  as  $v + w$  where  $v$  is the solution of the Dirichlet problem

$$\begin{cases} -D_\alpha \left( A_{ij,0}^{\alpha\beta} D_\beta v^j \right) = 0 & \text{in } B_R \\ v - u \in H_0^{1,2} (B_R, R^N). \end{cases}$$

For every  $0 < \sigma \leq R$  from Lemma 3 it follows

$$\int_{B_\sigma} |Dv - (Dv)_\sigma|^2 dy \leq A \left( \frac{\sigma}{R} \right)^{n+2} \int_{B_R} |Dv - (Dv)_R|^2 dy,$$

hence

$$(12) \quad \int_{B_\sigma} |Du - (Du)_\sigma|^2 dy \leq 2A \left( \frac{\sigma}{R} \right)^{n+2} \int_{B_R} |Dv - (Dv)_R|^2 dy + 2 \int_{B_R} |Dw|^2 dy.$$

Now  $w \in H_0^{1,2} (B_R, R^N)$  satisfies

$$\begin{aligned} & \int_{B_R} A_{ij,0}^{\alpha\beta} D_\beta w^j D_\alpha \varphi^i dy \leq \int_{B_R} \left| A_{ij,0}^{\alpha\beta} - \tilde{A}_{ij}^{\alpha\beta} \right| \left| D_\beta w^j - (D_\beta w^j)_R \right| \left| D_\alpha \varphi^i \right| dy \\ & \leq \left( \int_{B_R} \omega^2 \left( |Du - (Du)_R|^2 \right) |Du - (Du)_R|^2 dy \right)^{1/2} \left( \int_{B_R} |D\varphi|^2 dy \right)^{1/2} \end{aligned}$$

for any  $\varphi \in H_0^{1,2}(B_R, R^N)$ , where  $\omega$  is from (5). Hence, choosing  $\varphi = w$ , we get

$$\nu^2 \int_{B_R} |Dw|^2 dy \leq \int_{B_R} \omega^2 (|Du - (Du)_R|^2) |Du - (Du)_R|^2 dy.$$

Now applying the Young inequality (with the complementary functions (7)) on the right-hand side, we obtain for every  $\varepsilon > 0$

$$(13) \quad \nu^2 \int_{B_R} |Dw|^2 dy \leq \varepsilon \int_{B_R} |Du - (Du)_R|^2 \ln_+ (a\varepsilon |Du - (Du)_R|^2) dy + \frac{2}{a} \int_{B_R} e^{\omega_R^2/\varepsilon-1} dy,$$

where  $\omega_R^2 = \omega^2(|Du - (Du)_R|^2)$ .

From (12) and (13) it follows

$$(14) \quad \int_{B_\sigma} |Du - (Du)_\sigma|^2 dy \leq 4A \left(\frac{\sigma}{R}\right)^{n+2} \int_{B_R} |Du - (Du)_R|^2 dy + \frac{2(2A+1)}{\nu^2} \left( \varepsilon \int_{B_R} |Du - (Du)_R|^2 \ln_+ (a\varepsilon |Du - (Du)_R|^2) dy + \frac{2}{a} \int_{B_R} e^{\omega_R^2/\varepsilon-1} dy \right),$$

We can estimate the right-hand side by means of Lemma 1 (c) (for the quasiconcave case), Lemma 6 and we get

$$\int_{B_\sigma} |Du - (Du)_\sigma|^2 dy \leq 4A \left(\frac{\sigma}{R}\right)^{n+2} \int_{B_R} |Du - (Du)_R|^2 dy + \frac{2(2A+1)}{\nu^2} \left[ \varepsilon c \left(\frac{M}{\nu}\right)^2 \left( \int_{B_{2R}} \ln_+^{q/(q-1)} (4a\varepsilon |Du - (Du)_{2R}|^2) dy \right)^{1-1/q} \times \int_{B_{2R}} |Du - (Du)_{2R}|^2 dy + \frac{2\alpha_n R^n}{a} e^{l_2\omega^2(l_2U_R)/\varepsilon-1} \right].$$

Setting

$$\phi(t) = \int_{B_t} |Du - (Du)_t|^2 dy,$$

$$F_\varepsilon(t) = \left( \int_{B_t} \ln_+^{q/(q-1)} (4a\varepsilon |Du - (Du)_t|^2) dy \right)^{1-1/q},$$

we can rewrite the previous inequality as follows:

$$(15) \quad \phi(\sigma) \leq 4A \left(\frac{\sigma}{R}\right)^{n+2} \phi(R) + \frac{K\varepsilon}{\nu^2} F_\varepsilon(2R)\phi(2R) \\ + \frac{2^4 \alpha_n A}{a\nu^2} e^{l_2 \omega^2 (2^n l_2 U_{2R}) / \varepsilon - 1} R^n,$$

where  $K = c(n, N, q) (M/\nu)^8$ . From the assumptions of Theorem it follows that there exists  $d \in (0, d_x/2)$  such that (8) holds. Now we are going to prove that

$$(16) \quad \phi\left(2\tau^k d\right) \leq \tau^{kn} \phi(2d)$$

for every natural number  $k$  and  $\tau = (2^{n+5}A)^{-1/2}$ . Let  $k = 1$ . If we put in (15)  $a = 1/CU_{2d}$ ,  $\varepsilon = l_2 \omega^2 (2^n l_2 U_{2d})$ ,  $\sigma = 2\tau d$  and  $R = d$  we get

$$\phi(2\tau d) \leq 2^{n+4} A \tau^{n+2} \phi(d) + \frac{K l_2 \omega^2}{\nu^2} F_\varepsilon(2d) \phi(2d) + \frac{2^4 \alpha_n A}{\nu^2} C U_{2d} d^n \\ \leq 2^{n+4} A \tau^{n+2} \phi(2d) + \frac{K l_2 \omega^2}{\nu^2} b^{1-1/q} F_\varepsilon(2d) \phi(2d) + \frac{1}{4} \tau^n \phi(2d) \\ \leq \left(2^{n+4} A \tau^2 + \frac{1}{4} + \frac{1}{4}\right) \tau^n \phi(2d) = \tau^n \phi(2d).$$

Thus (16) holds for  $k = 1$ . Consequently  $U_{2\tau d} \leq U_{2d}$  and by means of Proposition 3 we have  $F_\varepsilon(2\tau d) \leq b^{1-1/q} F_\varepsilon(2d)$ .

Let us suppose that (16) holds for  $k \geq 1$ . Similarly to consideration above we have  $U_{2\tau^k d} \leq U_{2d}$  and  $F_\varepsilon(2\tau^k d) \leq b^{1-1/q} F_\varepsilon(2d)$ . We will show that (16) holds for  $k + 1$ . Setting  $a = 1/CU_{2d}$ ,  $\varepsilon = l_2 \omega^2 (2^n l_2 U_{2d})$ ,  $\sigma = 2\tau^{k+1} d$  and  $R = \tau^k d$  in (15) we obtain

$$\phi(2\tau^{k+1} d) \leq 2^{n+4} A \tau^{n+2} \phi\left(\tau^k d\right) + \frac{K l_2 \omega^2}{\nu^2} F_\varepsilon(2\tau^k d) \phi(2\tau^k d) \\ + \frac{2^4 \alpha_n A}{\nu^2} e^{\omega^2 (2^n l_2 U_{2\tau^k d}) / \omega^2 (2^n l_2 U_{2d}) - 1} \tau^{kn} C U_{2d} d^n \\ \leq 2^{n+4} A \tau^{n+2} \phi\left(2\tau^k d\right) + \frac{K l_2 \omega^2}{\nu^2} F_\varepsilon(2\tau^k d) \phi(2\tau^k d) + \frac{1}{4} \tau^{(k+1)n} \phi(2d) \\ \leq 2^{n+4} A \tau^{n+2} \tau^{kn} \phi(2d) + \frac{K l_2 \omega^2}{\nu^2} b^{1-1/q} F_\varepsilon(2d) \tau^{kn} \phi(2d) + \frac{1}{4} \tau^{(k+1)n} \phi(2d) \\ \leq \left(2^{n+4} A \tau^2 + \frac{1}{4} + \frac{1}{4}\right) \tau^{(k+1)n} \phi(2d) = \tau^{(k+1)n} \phi(2d).$$

Let us consider the two possibilities:

(a) if  $\tau \leq t < 1$ , then  $t^{-n}\phi(td) \leq \tau^{-n}\phi(td) \leq \tau^{-n} \sup_{t \in [\tau,1]} \phi(td)$  and also

$$(17) \quad \phi(td) \leq \left( \tau^{-n} \sup_{t \in [\tau,1]} \phi(td) \right) t^n,$$

(b) if  $0 < t < \tau$ , then there exists natural  $k \geq 1$  such that  $\tau^{k+1} \leq t < \tau^k$ . From Proposition 3, (8), (16) and (15) with  $a = 1/CU_{2d}$ ,  $\varepsilon = l_2\omega^2(2^n l_2 U_{2d})$ ,  $\sigma = td$  and  $R = \tau^k d$  we have

$$(18) \quad \begin{aligned} \phi(td) &= \phi\left(\frac{t}{\tau^k}(\tau^k d)\right) \\ &\leq 4A \left(\frac{t}{\tau^k}\right)^{n+2} \phi(\tau^k d) + \frac{K\varepsilon}{\nu^2} F_\varepsilon(2\tau^k d) \phi(2\tau^k d) \\ &\quad + \frac{2^4 \alpha_n A}{a\nu^2} e^{l_2\omega^2(2^n l_2 U_{2\tau^k d})/\varepsilon - 1} \tau^{kn} d^n \\ &\leq 4A \left(\frac{t}{\tau^k}\right)^{n+2} \tau^{kn} \phi(2d) + \frac{Kl_2\omega^2}{\nu^2} b^{1-1/q} F_\varepsilon(2d) \tau^{kn} \phi(2d) \\ &\quad + \frac{2^4 \alpha_n A}{\nu^2} CU_{2d} \tau^{kn} d^n \\ &\leq \left( 4A \left(\frac{t}{\tau^k}\right)^{n+2} \tau^{kn} + \tau^{(k+1)n} \right) \phi(2d) \\ &\leq \left( 4A\tau^{-n} \left(\frac{t}{\tau^k}\right)^{n+2} + 1 \right) \tau^{(k+1)n} \phi(2d) < (4A\tau^{-n} + 1) t^n \phi(2d). \end{aligned}$$

In both cases (17) and (18) we obtain

$$t^{-n}\phi(td) \leq c, \quad t \in (0, 1],$$

where  $c = \max\{\tau^{-n} \sup_{t \in [\tau,1]} \phi(td), (4A\tau^{-n} + 1)\phi(2d)\} = (4A\tau^{-n} + 1)\phi(2d)$ . Let  $0 < r < \text{dist}(B(x, r_x), \partial\Omega)$ . Hence  $U(y, r)$  is uniformly continuous for fixed  $r$  in  $\overline{B(x, r_x)} \subset \Omega$ . According to Proposition 3, the expression

$$\frac{Kl_2\omega^2}{\nu^2} \left( b \int_{B(y,r)} \ln_+^{q/(q-1)} \left( \frac{4l_2\omega^2 |Du - (Du)_{y,r}|^2}{CU(y,r)} \right) dz \right)^{1-1/q}$$

is also uniformly continuous with respect to  $y$  in  $\overline{B(x, r_x)}$  and we arrive at the conclusion. □

**Acknowledgement.** I thank Oldřich John, Jana Stará and Eugen Viszus for helpful discussions.

## REFERENCES

- [1] Campanato S., *Sistemi ellittici in forma divergenza. Regolarità all'interno*, Quaderni, Pisa, 1980.
- [2] Campanato S., *Hölder continuity of the solutions of some non-linear elliptic systems*, Adv. Math. **48** (1983), 16–43.
- [3] Giaquinta M., *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Annals of Mathematics Studies N. 105, Princeton University Press, Princeton, 1983.
- [4] Grevholm B., *On the structure of the spaces  $\mathcal{L}_k^{p,\lambda}$* , Math. Scand. **26** (1970), 189–196.
- [5] Kokilashvili V., Krbeč M., *Weighted Norm Inequalities in Lorentz and Orlicz Spaces*, World Scientific, London, 1991.
- [6] Kufner A., John O., Fučík S., *Function Spaces*, Academia, Prague, 1977.
- [7] Morrey C.B., Jr., *Multiple Integrals in the Calculus of Variations*, Springer-Verlag, Heidelberg, 1966.
- [8] Nečas J., *Introduction to the Theory of Nonlinear Elliptic Equations*, Band 52, Teubner-Texte zur Mathematik, Leipzig, 1983.
- [9] Ziemer W.P., *Weakly Differentiable Functions*, Springer-Verlag, Heidelberg, 1989.

ÚSTAV MATEMATIKY, FAST VUT, ŽIŽKOVA 17, 60200 BRNO, CZECH REPUBLIC  
E-mail: mddan@fce.vutbr.cz

(Received August 14, 1995, revised February 14, 1996)