

Yanzheng Duan; Shutao Chen

A note on Schroeder-Bernstein Property and Primary Property of Orlicz function spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 39 (1998), No. 1, 55--57

Persistent URL: <http://dml.cz/dmlcz/118984>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1998

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A note on Schroeder-Bernstein Property and Primary Property of Orlicz function spaces

YANZHENG DUAN, SHUTAO CHEN

Abstract. It is shown in the note that every reflexive Orlicz function space has the Schroeder-Bernstein Property and the Primary Property.

Keywords: Orlicz function spaces, Schroeder-Bernstein Property, Primary Property

Classification: 46B20

Let $G = [0, 1]$ and μ be the Lebesgue measure on G . We denote by $M : (-\infty, +\infty) \rightarrow [0, +\infty)$ a continuous, convex and even function satisfying $M(u) = 0$ iff $u = 0$ and $M(u)/u \rightarrow 0(+\infty)$ as $u \rightarrow 0(+\infty)$; by $N(v)$ the complementary function of $M(u)$, i.e., $N(v) := \max_u \{uv - M(u)\}$. We say $M \in \Delta_2$ if for any $u_0 > 0$ there exists $K > 2$ such that $M(2u) \leq KM(u)$, $u \geq u_0$. For every μ -measurable function $f : G \rightarrow (-\infty, +\infty)$, let $\varrho_M(f) = \int_G M(f(t)) d\mu$; then the Orlicz space

$$L_M = \{f : \varrho_M(af) < +\infty \text{ for some } a > 0\}$$

endowed with the Luxemburg norm

$$\|f\| = \inf\{r : \varrho_M(f/r) \leq 1\}$$

or Orlicz norm

$$\|f\|_M = \min_{k>0} [1 + \varrho_M(kf)]/k$$

is a Banach space and $\|f\|_M \leq \|f\| \leq 2\|f\|_M$ for every $f \in L_M$. More details about Orlicz spaces can be found in [2] and [4].

Let Y be a closed subspace of a Banach space X . Y is called a complemented subspace of X if there exists a linear, continuous and surjective projection from X to Y . A Banach space X is said to have the *Schroeder-Bernstein Property (SBP)* if for any Banach space Y , X is isomorphic to Y whenever X is isomorphic to a complemented subspace of X . A Banach space X is said to have the *Primary Property* if, for every linear, bounded projection P of X , X is isomorphic to PX or $(I - P)X$. Many spaces, for example, L^p ($1 < p < +\infty$) and James space J , have *SBP* and *Primary Property* (see [1]).

Without loss of generality, let $M(1) = 1$. Then $(L_M, \|\cdot\|)$ is an r.i. (i.e. rearrangement invariant) function space. More details about this space can be found in [3]. By Proposition 2.b.5 in [3], the Boyd indices for L_M are

$$p_{L_M} = \sup \left\{ p : \inf_{\lambda, t \geq 1} \frac{M(t\lambda)}{M(\lambda)t^p} > 0 \right\}$$

and

$$q_{L_M} = \inf \left\{ q : \sup_{\lambda, t \geq 1} \frac{M(t\lambda)}{M(\lambda)t^q} < +\infty \right\}.$$

In general, $1 \leq p_{L_M} \leq q_{L_M} \leq +\infty$.

Theorem 1. *For the Orlicz space $(L_M, \|\cdot\|)$, we have*

- (1) $q_{L_M} < +\infty$ if and only if $M \in \Delta_2$;
- (2) $p_{L_M} > 1$ if and only if $N \in \Delta_2$.

PROOF: (1) Necessity. If $q_{L_M} < +\infty$, then there exist constants $K > 1$ and $q_0 \geq 1$ such that $M(t\lambda)/(M(\lambda)t^{q_0}) \leq K$ for all $\lambda, t \geq 1$. Let $t = 2$, then $M(2\lambda) \leq 2^{q_0} K M(\lambda)$ for all $\lambda \geq 1$, i.e., $M \in \Delta_2$.

Sufficiency. Since $M \in \Delta_2$, by [4], there exists a constant $K > 2$ such that $M(2t) \leq K M(t)$ for all $t \geq 1$. Choose an integer $n \geq 0$ such that $2^n \leq t < 2^{n+1}$; then for all $\lambda \geq 1$ and $q > 1$ satisfying $K/2^q \leq 1$, we have

$$\frac{M(t\lambda)}{M(\lambda)t^q} \leq \frac{M(2^{n+1}\lambda)}{2^{nq}M(\lambda)} \leq \frac{K^{n+1}M(\lambda)}{2^{nq}M(\lambda)} = K \left(\frac{K}{2^q} \right)^n \leq K.$$

Thus, by the definition of q_{L_M} , we have $q_{L_M} < +\infty$.

(2) Necessity. If $p_{L_M} > 1$, then there exist $\varepsilon > 0$ and $\delta > 0$ such that $M(t\lambda)/(M(\lambda)t^{1+2\varepsilon}) \geq \delta$ for all $\lambda, t \geq 1$. Choose t_0 satisfying $t_0^\varepsilon \delta \geq 1$, then for all $\lambda \geq 1$, we have

$$\frac{M(t_0\lambda)}{M(\lambda)t_0^{1+\varepsilon}} \geq t_0^\varepsilon \delta \geq 1.$$

Therefore, $M(t_0\lambda) \geq t_0^{1+\varepsilon} M(\lambda)$ for all $\lambda \geq 1$. So $N \in \Delta_2$ by [4].

Sufficiency. If $N \in \Delta_2$, then, by [4], there exists $\varepsilon > 0$ such that $M(2\lambda) \geq 2^{1+\varepsilon} M(\lambda)$ for all $\lambda \geq 1$. Choose a positive integer k_0 such that $p = (1+\varepsilon)k_0/(1+k_0) > 1$. For all $\lambda \geq 1$ and $t \geq 2^{k_0}$ choose integer k such that $2^{k_0} \leq 2^k \leq t < 2^{k+1}$, then

$$\frac{M(t\lambda)}{M(\lambda)t^p} \geq \frac{M(2^k\lambda)}{M(\lambda)2^{(k+1)p}} \geq \frac{2^{(1+\varepsilon)k} M(\lambda)}{2^{(k+1)p} M(\lambda)} = 2^{(1+\varepsilon)k - (k+1)p} \geq 2^0 = 1.$$

Note the last inequality of the above formula is assured by the monotone increasing of the function $f(x) = (1+\varepsilon)x/(1+x)$ ($x > 0$).

On the other hand, for all $\lambda \geq 1$ and $t \in [1, 2^{k_0}]$, we have

$$\frac{M(t\lambda)}{M(\lambda)t^p} \geq \frac{M(\lambda)}{2^{k_0 p} M(\lambda)} = \frac{1}{2^{k_0 p}}.$$

Therefore, for all $\lambda, t \geq 1$, we have

$$\frac{M(t\lambda)}{M(\lambda)t^p} \geq \min \left\{ 1, \frac{1}{2^{k_0 p}} \right\} > 0.$$

Thus, $p_{L_M} \geq p > 1$. □

By [4] and Theorem 1, we immediately get the following corollary.

Corollary 2. L_M is reflexive if and only if $p_{L_M} > 1$ and $q_{L_M} < +\infty$.

Theorem 3. If L_M is reflexive, then L_M has SBP.

PROOF: If L_M is reflexive, then by Theorem 1, $p_{L_M} > 1$ and $q_{L_M} < +\infty$. Therefore, if a Banach space X is isomorphic to a complemented subspace of L_M and L_M is also isomorphic to a complemented subspace of X , then Proposition 2.d.5 in [3] implies that L_M is isomorphic to X . So L_M has SBP. □

Theorem 4. If L_M is reflexive, then L_M has the Primary Property.

PROOF: If L_M is reflexive, then by [4] and Theorem 1, L_M is separable, $p_{L_M} > 1$ and $q_{L_M} < +\infty$. Since L_M is an r.i. function space, Theorem 2.d.11 in [3] implies that L_M has the Primary Property. □

REFERENCES

- [1] Casazza P.G., *The Schroeder-Bernstein Property for Banach spaces*, Contemporary Mathematics **85** (1989), 61–77.
- [2] Shutao Chen, *Geometry of Orlicz spaces*, Dissert. Math. **356** (1996), 1–204.
- [3] Lindenstrauss J., Tzafriri L., *Classical Banach Spaces II. Function Spaces*, Springer-Verlag, Berlin, 1979.
- [4] Congxin Wu, Tingfu Wang, *Orlicz Spaces and Their Applications* (in Chinese), Heilongjiang Science and Technology Publishing House, Harbin, 1983.

DEPARTMENT OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN, 150001,
P.R. CHINA

DEPARTMENT OF MATHEMATICS, HARBIN NORMAL UNIVERSITY, HARBIN, 150080, P.R. CHINA

(Received April 3, 1997)