

Tiberiu Trif

Singularities and equicontinuity of certain families of set-valued mappings

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 39 (1998), No. 2, 353--365

Persistent URL: <http://dml.cz/dmlcz/119012>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1998

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Singularities and equicontinuity of certain families of set-valued mappings

TIBERIU TRIF

*Abstract.* In the present paper we establish an abstract principle of condensation of singularities for families consisting of set-valued mappings. By using it as a basic tool, the condensation of the singularities and the equicontinuity of certain families of generalized convex set-valued mappings are studied. In particular, a principle of condensation of the singularities of families of closed convex processes is derived. This principle immediately yields the uniform boundedness theorem stated in [1, Theorem 2.3.1].

*Keywords:* condensation of the singularities, equicontinuity, generalized convex set-valued mappings, closed convex processes

*Classification:* 46N10, 54C60, 26B25

### 1. Introduction

Many textbooks consider the following *principle of uniform boundedness* as one of the most important results in functional analysis.

**Theorem 1.1.** *Let  $X$  be a Banach space, let  $Y$  be a normed linear space, and let  $\mathcal{F}$  be a family of continuous linear mappings from  $X$  into  $Y$  such that*

$$\sup \{ \|f(x)\| \mid f \in \mathcal{F} \} < \infty \quad \text{for all } x \in X.$$

*Then*  $\sup \{ \|f\| \mid f \in \mathcal{F} \} < \infty$ .

This theorem reveals that if  $\sup \{ \|f\| \mid f \in \mathcal{F} \} = \infty$ , then there exists at least one singularity of  $\mathcal{F}$ , i.e. a point  $x \in X$  such that  $\sup \{ \|f(x)\| \mid f \in \mathcal{F} \} = \infty$ . More informations about the set of singularities of  $\mathcal{F}$  can be obtained if the following *principle of condensation of the singularities* is applied instead of the uniform boundedness principle.

**Theorem 1.2.** *Let  $X$  be a Banach space, let  $Y$  be a normed linear space, and let  $\mathcal{F}$  be a family of continuous linear mappings from  $X$  into  $Y$  such that*

$$\sup \{ \|f\| \mid f \in \mathcal{F} \} = \infty.$$

*Then the set  $S_{\mathcal{F}}$  of all  $x \in X$  for which*

$$\sup \{ \|f(x)\| \mid f \in \mathcal{F} \} = \infty$$

is residual.

Theorem 1.2 has been generalized by numerous authors and in several directions. Here we merely mention the investigations by W.W. Breckner [3], who established a principle of condensation of singularities for lower semicontinuous mappings defined on a topological space and taking values in the power set of a topological space. In the present paper we continue these investigations, but unlike W.W. Breckner we deal with families of set-valued mappings defined on topological spaces. In Section 2 we introduce the concept of a singularity for such families and after that we prove a general principle of condensation of the singularities of a family consisting of arbitrary upper semicontinuous set-valued mappings. In the following three sections we give several applications of this principle under the assumption that the involved set-valued mappings have some additional algebraic properties. They relate to families that consist either of  $(A, s)$ -convex set-valued mappings or of  $(A, s)$ -convex real-valued mappings that are not equicontinuous at the origin as well as to families of closed convex processes. An important corollary is the uniform boundedness theorem involving closed convex processes which has been stated by J.P. Aubin and H. Frankowska [1, Theorem 2.3.1].

Throughout the paper the set of all positive integers is denoted by  $\mathbb{N}$ . Given any subset  $M$  of a topological space, we denote by  $\text{cl } M$  its closure. Given any set  $Y$ , we denote by  $\mathcal{P}_0(Y)$  the set consisting of all nonempty subsets of  $Y$ . Given a topological linear space  $X$ , we denote by  $\mathcal{O}_{ac}(X)$  the class consisting of all nonempty, open and absolutely convex subsets of  $X$ .

## 2. An abstract principle of condensation of the singularities of families of set-valued mappings

Let  $X$  and  $Y$  be topological spaces, let  $F$  be a mapping from  $X$  to  $2^Y$ , and let  $x_0$  be any point of  $X$ . Recall that  $F$  is said to be *upper semicontinuous* at  $x_0$  if for every open subset  $Y_0$  of  $Y$  with  $F(x_0) \subseteq Y_0$ , there exists a neighbourhood  $V$  of  $x_0$  such that  $F(x) \subseteq Y_0$  for all  $x \in V$ . If  $F$  is upper semicontinuous at each point of  $X$ , then  $F$  is called *upper semicontinuous* (on  $X$ ).

Let  $I$  be a nonempty set, let  $B : I \times \mathbb{N} \rightarrow 2^Y$  be a mapping whose values are closed subsets of  $Y$ , and let  $\mathcal{F}$  be a family of mappings from  $X$  to  $\mathcal{P}_0(Y)$ .

We say that  $\mathcal{F}$  is *B-bounded at a point*  $x_0 \in X$  if there exists a family  $\{y_F \mid F \in \mathcal{F}\}$  with  $y_F \in F(x_0)$  ( $F \in \mathcal{F}$ ), satisfying the following condition: for every  $i \in I$  one can select a positive integer  $n$  such that  $\{y_F \mid F \in \mathcal{F}\} \subseteq B(i, n)$ . If  $\mathcal{F}$  is *B-bounded at each point of*  $X$ , then we say that  $\mathcal{F}$  is *pointwise B-bounded* (on  $X$ ). A point in  $X$  at which  $\mathcal{F}$  is not *B-bounded* is called a *B-singularity* of  $\mathcal{F}$ . The set of all *B-singularities* of  $\mathcal{F}$  will be denoted by  $S_{\mathcal{F}}(B)$ . Obviously,  $\mathcal{F}$  is pointwise *B-bounded* if and only if the set  $S_{\mathcal{F}}(B)$  is empty.

We say that  $\mathcal{F}$  is *uniformly B-bounded* if for every  $i \in I$  there exist a positive integer  $n$  and a nonempty open subset  $X_0$  of  $X$  satisfying the following condition:

for each point  $x \in X_0$  one can select a family  $\{y_F \mid F \in \mathcal{F}\}$  with  $y_F \in F(x)$  ( $F \in \mathcal{F}$ ) such that  $\{y_F \mid F \in \mathcal{F}\} \subseteq B(i, n)$ .

The above definitions are inspired by [1] and [3]. After these preliminaries we are in position to state the main result of the paper.

**Theorem 2.1.** *Let  $X$  and  $Y$  be topological spaces, and let  $\mathcal{F}$  be a family of upper semicontinuous mappings from  $X$  to  $\mathcal{P}_0(Y)$  which is not uniformly  $B$ -bounded. Then the following assertions are true:*

1°  $S_{\mathcal{F}}(B)$  is a residual set;

2° if in addition  $X$  is of second category, then  $S_{\mathcal{F}}(B)$  is of second category and hence nonempty;

3° if in addition  $X$  is a Baire space satisfying the separation axiom  $T_1$  and without isolated points, then  $S_{\mathcal{F}}(B)$  is of second category and uncountable.

PROOF: 1° Since  $\mathcal{F}$  is not uniformly  $B$ -bounded, choose  $i \in I$  such that for every positive integer  $n$  and every nonempty open subset  $X_0$  of  $X$  there exist a point  $x \in X_0$  and a mapping  $F \in \mathcal{F}$  satisfying  $F(x) \subseteq Y \setminus B(i, n)$ . Put

$$(2.1) \quad G_n := \{x \in X \mid \exists F \in \mathcal{F} : F(x) \subseteq Y \setminus B(i, n)\}$$

for all  $n \in \mathbb{N}$ . We claim that all the sets  $G_n$  ( $n \in \mathbb{N}$ ) are open and dense in  $X$ .

Indeed, let  $n$  be any positive integer, and let  $x_0$  be any point in  $G_n$ . According to (2.1) there exists a mapping  $F \in \mathcal{F}$  such that  $F(x_0) \subseteq Y \setminus B(i, n)$ . Taking into consideration that  $F$  is upper semicontinuous at  $x_0$  and that  $Y \setminus B(i, n)$  is open, it follows that there exists a neighbourhood  $V$  of  $x_0$  such that  $F(x) \subseteq Y \setminus B(i, n)$  for all  $x \in V$ . Hence  $V \subseteq G_n$ . Therefore  $x_0$  is an interior point of  $G_n$ . Since  $x_0$  was arbitrary in  $G_n$ , we can conclude that the set  $G_n$  is open.

Suppose now that there is a positive integer  $n$  for which  $G_n$  is not dense in  $X$ . Then  $X \setminus \text{cl } G_n$  is open and nonempty. The choice of  $i$  ensures that there exist a point  $x \in X \setminus \text{cl } G_n$  and a mapping  $F \in \mathcal{F}$  such that  $F(x) \subseteq Y \setminus B(i, n)$ . But, in view of (2.1), we have  $x \in G_n \subseteq \text{cl } G_n$ , which is a contradiction.

Consequently, all the sets  $G_n$  ( $n \in \mathbb{N}$ ) are open and dense, as claimed. Since

$$\bigcap_{n \in \mathbb{N}} G_n \subseteq S_{\mathcal{F}}(B),$$

it follows that  $S_{\mathcal{F}}(B)$  is a residual set.

The assertions 2° and 3° are immediate consequences of Proposition 2.4 and Proposition 2.6 in [3].  $\square$

### 3. Singularities and equicontinuity of families of generalized convex set-valued mappings

Assume that  $A$  is a subset of the open interval  $]0, 1[$  having 0 as a cluster point, and that  $s$  is a positive real number. Let  $X$  and  $Y$  be topological linear spaces,

and let  $M$  be a nonempty convex subset of  $X$ . According to W. W. Breckner [4] a mapping  $F : M \rightarrow \mathcal{P}_0(Y)$  is said to be  $(A, s)$ -convex if

$$(1 - a)^s F(x) + a^s F(y) \subseteq F((1 - a)x + ay)$$

whenever  $a \in A$  and  $x, y \in M$ .

Let  $X$  and  $Y$  be topological linear spaces, let  $o_X$  and  $o_Y$  denote the zero-elements of  $X$  and  $Y$ , respectively, let  $M$  be a nonempty subset of  $X$ , and let  $\mathcal{F}$  be a family of mappings from  $M$  to  $\mathcal{P}_0(Y)$ .

We say that  $\mathcal{F}$  is *bounded at a point*  $x_0 \in M$  if there exists a family  $\{y_F \mid F \in \mathcal{F}\}$  with  $y_F \in F(x_0)$  ( $F \in \mathcal{F}$ ) which is bounded, i.e. for every neighbourhood  $V$  of  $o_Y$  one can find a positive integer  $n$  such that  $\{y_F \mid F \in \mathcal{F}\} \subseteq nV$ . If  $\mathcal{F}$  is bounded at each point of  $M$ , then we say that  $\mathcal{F}$  is *pointwise bounded* (on  $M$ ). Any point in  $M$  at which  $\mathcal{F}$  is not bounded is called a *singularity* of  $\mathcal{F}$ . The set of all singularities of  $\mathcal{F}$  will be denoted by  $S_{\mathcal{F}}$ . If the set  $M$  is balanced, then we call a point  $x_0 \in M$  a *weak singularity* of  $\mathcal{F}$  if  $\mathcal{F}$  is not bounded either at  $x_0$  or at  $-x_0$ . The set of all weak singularities of  $\mathcal{F}$  will be denoted by  $WS_{\mathcal{F}}$ . Obviously  $WS_{\mathcal{F}} = S_{\mathcal{F}} \cup (-S_{\mathcal{F}})$  holds. The concept of weak singularity has already been introduced in [5].

We say that  $\mathcal{F}$  is *equi-lower semicontinuous* (respectively *equi-upper semicontinuous*) at a point  $x_0 \in M$  if for every neighbourhood  $V$  of  $o_Y$  there exists a neighbourhood  $U$  of  $x_0$  such that

$$F(x_0) \subseteq F(x) + V \quad (\text{respectively } F(x) \subseteq F(x_0) + V)$$

for all  $F \in \mathcal{F}$  and all  $x \in U \cap M$ . We say that  $\mathcal{F}$  is *equicontinuous at*  $x_0$  if it is both equi-lower semicontinuous and equi-upper semicontinuous at this point.

**Proposition 3.1.** *Let  $X$  and  $Y$  be topological linear spaces, let  $\mathcal{B}$  be a neighbourhood base at  $o_Y$  composed of closed sets, and let  $B : \mathcal{B} \times \mathbb{N} \rightarrow 2^Y$  be the mapping defined by*

$$B(V, n) := nV \quad \text{for all } (V, n) \in \mathcal{B} \times \mathbb{N}.$$

Further let  $\mathcal{F}$  be a family of  $(A, s)$ -convex mappings from a set  $M \in \mathcal{O}_{ac}(X)$  to  $\mathcal{P}_0(Y)$ , and let  $\mathcal{G} := \mathcal{F} \cup \{G_F \mid F \in \mathcal{F}\}$ , where  $G_F : M \rightarrow \mathcal{P}_0(Y)$  is the mapping defined by  $G_F(x) := F(-x)$  ( $F \in \mathcal{F}$ ). Then the following assertions are true:

- 1° if all mappings in  $\mathcal{F}$  are upper semicontinuous on  $M$ , then all mappings in  $\mathcal{G}$  are upper semicontinuous on  $M$ , too;
- 2°  $\mathcal{F}$  is bounded at a point  $x_0 \in M$  if and only if  $\mathcal{F}$  is  $B$ -bounded at  $x_0$ ;
- 3°  $WS_{\mathcal{F}} = S_{\mathcal{G}}(B)$ ;
- 4° if  $\mathcal{F}$  is bounded and equi-lower semicontinuous at  $o_X$ , then  $\mathcal{G}$  is uniformly  $B$ -bounded on  $M$ ;

5° if  $\mathcal{G}$  is uniformly  $B$ -bounded on  $M$ , and

$$\frac{1}{\lambda}F(o_X) \subseteq F(o_X) \subseteq \lambda F(o_X) \text{ for all } F \in \mathcal{F} \text{ and all } \lambda \in ]0, 1],$$

then  $\mathcal{F}$  is equicontinuous at  $o_X$ .

PROOF: The assertions 1°, 2° and 3° are obvious.

4° Let  $V$  be any closed neighbourhood of  $o_Y$ . Choose a balanced neighbourhood  $V_0$  of  $o_Y$  such that  $V_0 + V_0 \subseteq V$ . Since  $\mathcal{F}$  is bounded at  $o_X$ , there exists a bounded family  $\{x_F \mid F \in \mathcal{F}\}$  with  $x_F \in F(o_X)$  ( $F \in \mathcal{F}$ ). Therefore one can find a positive integer  $n$  so that

$$\{x_F \mid F \in \mathcal{F}\} \subseteq nV_0.$$

On the other hand, taking into consideration that  $\mathcal{F}$  is equi-lower semicontinuous at  $o_X$ , we can select a balanced neighbourhood  $U$  of  $o_X$  such that

$$F(o_X) \subseteq F(x) + nV_0 \text{ for all } F \in \mathcal{F} \text{ and all } x \in U \cap M.$$

Put  $M_0 := \text{int}(U \cap M)$ . Then  $M_0$  is a nonempty open subset of  $M$ . Let  $x$  be any point in  $M_0$ . For every  $F \in \mathcal{F}$  we have

$$x_F \in F(o_X) \subseteq F(x) + nV_0 \quad \text{and} \quad x_F \in F(o_X) \subseteq F(-x) + nV_0.$$

Therefore, for every  $F \in \mathcal{F}$  we can select elements  $y_F \in F(x)$  and  $z_F \in F(-x) = G_F(x)$  such that  $x_F \in y_F + nV_0$  and  $x_F \in z_F + nV_0$ . Consequently

$$y_F \in x_F - nV_0 \subseteq nV_0 + nV_0 \subseteq nV$$

and

$$z_F \in x_F - nV_0 \subseteq nV_0 + nV_0 \subseteq nV.$$

Hence

$$\{y_F \mid F \in \mathcal{F}\} \cup \{z_F \mid F \in \mathcal{F}\} \subseteq nV = B(V, n).$$

Since  $x$  was arbitrary in  $M_0$ , we can conclude that  $\mathcal{G}$  is uniformly  $B$ -bounded on  $M$ .

5° Let  $V$  be any neighbourhood of  $o_Y$ . Select a balanced neighbourhood  $V_0$  of  $o_Y$  such that  $V_0 + V_0 \subseteq V$  and a closed neighbourhood  $W$  of  $o_Y$  such that  $W \subseteq V_0$ . Since the family  $\mathcal{G}$  is uniformly  $B$ -bounded on  $M$ , there exist a positive integer  $n$  and a nonempty open subset  $M_0$  of  $M$  satisfying the following condition: for each point  $x \in M_0$  one can select the families  $\{y_F \mid F \in \mathcal{F}\}$  and  $\{z_F \mid F \in \mathcal{F}\}$  with  $y_F \in F(x)$ ,  $z_F \in F(-x)$  ( $F \in \mathcal{F}$ ) such that

$$(3.1) \quad \{y_F \mid F \in \mathcal{F}\} \cup \{z_F \mid F \in \mathcal{F}\} \subseteq nW.$$

Let  $x_0$  be any point in  $M_0$ . The choice of  $M_0$  ensures that  $-x_0 \in M_0$ . Since  $M_0$  is open, we can find a balanced neighbourhood  $U$  of  $o_X$  such that  $x_0 + U \subseteq M_0$  and  $-x_0 + U \subseteq M_0$ . Finally, choose  $a \in A$  so that  $a^s n / 2^s < 1$ , and put  $U_0 := aU$ .

Let  $x$  be an arbitrary point in  $U$ . From the equality

$$ax = \frac{a}{2}(x_0 + x) + \frac{a}{2}(-x_0 + x) + (1 - a)o_X,$$

we get

$$\left(\frac{a}{2}\right)^s F(x_0 + x) + \left(\frac{a}{2}\right)^s F(-x_0 + x) + (1 - a)^s F(o_X) \subseteq F(ax)$$

for all  $F \in \mathcal{F}$ . Taking into account that

$$F(o_X) \subseteq (1 - a)^s F(o_X) \quad (F \in \mathcal{F}),$$

we obtain

$$(3.2) \quad \left(\frac{a}{2}\right)^s F(x_0 + x) + \left(\frac{a}{2}\right)^s F(-x_0 + x) + F(o_X) \subseteq F(ax)$$

for all  $F \in \mathcal{F}$ . But  $x_0 + x$  and  $-x_0 + x$  are points in  $M_0$ . Therefore there exist families  $\{y_F \mid F \in \mathcal{F}\}$  and  $\{z_F \mid F \in \mathcal{F}\}$  such that  $y_F \in F(x_0 + x)$ ,  $z_F \in F(-x_0 + x)$  ( $F \in \mathcal{F}$ ) and which satisfy (3.1).

Let  $F$  be any mapping in  $\mathcal{F}$ . Taking into account (3.2) we get

$$\left(\frac{a}{2}\right)^s y_F + \left(\frac{a}{2}\right)^s z_F + F(o_X) \subseteq F(ax).$$

Consequently

$$\begin{aligned} F(o_X) &\subseteq F(ax) - \left(\frac{a}{2}\right)^s y_F - \left(\frac{a}{2}\right)^s z_F \subseteq \\ &\subseteq F(ax) - \left(\frac{a}{2}\right)^s nW - \left(\frac{a}{2}\right)^s nW \subseteq \\ &\subseteq F(ax) - \left(\frac{a}{2}\right)^s nV_0 - \left(\frac{a}{2}\right)^s nV_0 \subseteq F(ax) + V_0 + V_0 \subseteq \\ &\subseteq F(ax) + V. \end{aligned}$$

Hence we have proved that

$$F(o_X) \subseteq F(x) + V \quad \text{for all } F \in \mathcal{F} \text{ and all } x \in U_0.$$

Since  $V$  was an arbitrary neighbourhood of  $o_Y$ , we can conclude that  $\mathcal{F}$  is equi-lower semicontinuous at  $o_X$ .

Let again  $x$  be any point in  $U$ . From the equality

$$o_X = \frac{a}{2(1+a)}(x_0 - x) + \frac{a}{2(1+a)}(-x_0 - x) + \frac{1}{1+a}(ax),$$

we get

$$\left(\frac{a}{2}\right)^s F(x_0 - x) + \left(\frac{a}{2}\right)^s F(-x_0 - x) + F(ax) \subseteq (1 + a)^s F(o_X)$$

for all  $F \in \mathcal{F}$ . Taking into account that

$$(1 + a)^s F(o_X) \subseteq F(o_X) \quad (F \in \mathcal{F}),$$

we obtain

$$(3.3) \quad \left(\frac{a}{2}\right)^s F(x_0 - x) + \left(\frac{a}{2}\right)^s F(-x_0 - x) + F(ax) \subseteq F(o_X)$$

for all  $F \in \mathcal{F}$ . But  $x_0 - x$  and  $-x_0 - x$  are points in  $M_0$ . Therefore there exist families  $\{y'_F \mid F \in \mathcal{F}\}$  and  $\{z'_F \mid F \in \mathcal{F}\}$  such that  $y'_F \in F(x_0 - x)$ ,  $z'_F \in F(-x_0 - x)$  ( $F \in \mathcal{F}$ ) and

$$(3.4) \quad \{y'_F \mid F \in \mathcal{F}\} \cup \{z'_F \mid F \in \mathcal{F}\} \subseteq nW.$$

Let  $F$  be any mapping in  $\mathcal{F}$ . From (3.3) we obtain

$$\left(\frac{a}{2}\right)^s y'_F + \left(\frac{a}{2}\right)^s z'_F + F(ax) \subseteq F(o_X).$$

Taking now into account relation (3.4) we get

$$\begin{aligned} F(ax) &\subseteq F(o_X) - \left(\frac{a}{2}\right)^s y'_F - \left(\frac{a}{2}\right)^s z'_F \subseteq \\ &\subseteq F(o_X) - \left(\frac{a}{2}\right)^s nW - \left(\frac{a}{2}\right)^s nW \subseteq \\ &\subseteq F(o_X) - \left(\frac{a}{2}\right)^s nV_0 - \left(\frac{a}{2}\right)^s nV_0 \subseteq F(o_X) + V_0 + V_0 \subseteq \\ &\subseteq F(o_X) + V. \end{aligned}$$

Hence we have proved that

$$F(x) \subseteq F(o_X) + V \quad \text{for all } F \in \mathcal{F} \text{ and all } x \in U_0.$$

Since  $V$  was an arbitrary neighbourhood of  $o_Y$ , we can conclude that  $\mathcal{F}$  is equi-upper semicontinuous at  $o_X$ . Consequently,  $\mathcal{F}$  is equicontinuous at  $o_X$ .  $\square$

**Theorem 3.2.** *Let  $X$  and  $Y$  be topological linear spaces, and let  $\mathcal{F}$  be a family of upper semicontinuous  $(A, s)$ -convex mappings from a set  $M \in \mathcal{O}_{ac}(X)$  to  $\mathcal{P}_0(Y)$  which is not equicontinuous at  $o_X$  and satisfies the condition*

$$\frac{1}{\lambda} F(o_X) \subseteq F(o_X) \subseteq \lambda F(o_X) \quad \text{for all } F \in \mathcal{F} \text{ and all } \lambda \in ]0, 1].$$

Then the following assertions are true:

- 1°  $WS_{\mathcal{F}}$  is residual in  $M$ ;
- 2° if in addition  $X$  is of second category, then  $WS_{\mathcal{F}}$  is of second category in  $M$  and hence nonempty;
- 3° if in addition  $X$  is of second category and satisfies the separation axiom  $T_1$ , then  $WS_{\mathcal{F}}$  is of second category in  $M$  and uncountable.

PROOF: Follows immediately from Theorem 2.1 and Proposition 3.1.  $\square$

**Remark.** The assertion 1° of Theorem 3.2 does not remain true if we replace the set  $WS_{\mathcal{F}}$  by  $S_{\mathcal{F}}$ . This is shown by the following example. Let  $A := ]0, 1[$ , and let  $s := 1$ . For each positive integer  $n$ , define the function  $f_n : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f_n(x) := \begin{cases} 0 & \text{if } x < 0 \\ nx & \text{if } x \geq 0, \end{cases}$$

and then define the mapping  $F_n : \mathbb{R} \rightarrow \mathcal{P}_0(\mathbb{R})$  by  $F_n(x) := [f_n(x), \infty[$ . Obviously, the family  $\mathcal{F} := \{F_n \mid n \in \mathbb{N}\}$  satisfies the conditions of Theorem 3.2. Nevertheless  $S_{\mathcal{F}}$  is not residual because  $S_{\mathcal{F}} = ]0, \infty[$ .

**Theorem 3.3.** *Let  $X$  be a topological linear space of second category, let  $Y$  be a topological linear space, and let  $\mathcal{F}$  be a pointwise bounded family of upper semi-continuous  $(A, s)$ -convex mappings from a set  $M \in \mathcal{O}_{ac}(X)$  to  $\mathcal{P}_0(Y)$  satisfying the condition*

$$\frac{1}{\lambda}F(o_X) \subseteq F(o_X) \subseteq \lambda F(o_X) \text{ for all } F \in \mathcal{F} \text{ and all } \lambda \in ]0, 1].$$

Then  $\mathcal{F}$  is equicontinuous at  $o_X$ .

PROOF: Follows immediately from Theorem 3.2. □

#### 4. Singularities and equicontinuity of families of generalized convex real-valued functions

Let  $X$  be a linear space, and let  $M$  be a nonempty convex subset of  $X$ . Assume that  $A$  is a subset of  $]0, 1[$  having 0 as a cluster point, and that  $s$  is a positive real number. Having in mind the definitions of the  $s$ -convex and rationally  $s$ -convex functions given by W.W. Breckner [2], we say that a function  $f : M \rightarrow \mathbb{R}$  is  $(A, s)$ -convex if

$$f((1 - a)x + ay) \leq (1 - a)^s f(x) + a^s f(y)$$

whenever  $a \in A$  and  $x, y \in M$ .

It is easily seen that if  $f : M \rightarrow \mathbb{R}$  is an  $(A, s)$ -convex function, then the mapping  $F : M \rightarrow \mathcal{P}_0(\mathbb{R})$  defined by  $F(x) := [f(x), \infty[$  is  $(A, s)$ -convex, too.

Let  $X$  and  $Y$  be topological linear spaces, let  $o_X$  and  $o_Y$  be the zero-elements of  $X$  and  $Y$ , respectively, let  $M$  be a nonempty subset of  $X$ , and let  $\mathcal{F}$  be a family of mappings from  $M$  into  $Y$ .

If  $x_0$  is a point in  $M$ , then  $\mathcal{F}$  is said to be:

(i) *equicontinuous at  $x_0$*  if for every neighbourhood  $V$  of  $o_Y$  there exists a neighbourhood  $U$  of  $x_0$  such that

$$\{f(x) - f(x_0) \mid f \in \mathcal{F}\} \subseteq V \text{ for every } x \in U \cap M;$$

(ii) *bounded at  $x_0$*  if the set  $\{f(x_0) \mid f \in \mathcal{F}\}$  is bounded, i.e. for every neighbourhood  $V$  of  $o_Y$  there exists a positive integer  $n$  such that

$$\{f(x_0) \mid f \in \mathcal{F}\} \subseteq nV.$$

If  $\mathcal{F}$  is equicontinuous (respectively bounded) at each point of  $M$ , then  $\mathcal{F}$  is called *equicontinuous* (respectively *pointwise bounded*) on  $M$ .

Any point in  $M$  at which  $\mathcal{F}$  is not bounded is said to be a *singularity* of  $\mathcal{F}$ . The set of all singularities of  $\mathcal{F}$  is denoted by  $S_{\mathcal{F}}$ .

If  $M$  is balanced, then we say that a point  $x_0 \in M$  is a *weak singularity* of  $\mathcal{F}$  if the set

$$\{f(x_0) \mid f \in \mathcal{F}\} \cup \{f(-x_0) \mid f \in \mathcal{F}\}$$

is unbounded.

The set of all weak singularities of  $\mathcal{F}$  is denoted by  $WS_{\mathcal{F}}$ . Obviously,  $WS_{\mathcal{F}} = S_{\mathcal{F}} \cup (-S_{\mathcal{F}})$  holds.

The next theorem is an improvement of Theorem 3.1 in [5].

**Theorem 4.1.** *Let  $X$  be a topological linear space, and let  $\mathcal{F}$  be a family of lower semicontinuous  $(A, s)$ -convex functions from a set  $M \in \mathcal{O}_{ac}(X)$  to  $\mathbb{R}$  which is not equicontinuous at  $o_X$ . If either  $s \in ]0, 1[$ , or  $s = 1$  and  $\mathcal{F}$  is bounded at  $o_X$ , then the following assertions are true:*

1°  $WS_{\mathcal{F}}$  is residual in  $M$ ;

2° if in addition  $X$  is of second category, then  $WS_{\mathcal{F}}$  is of second category in  $M$  and hence nonempty;

3° if in addition  $X$  is of second category and satisfies the separation axiom  $T_1$ , then  $WS_{\mathcal{F}}$  is of second category in  $M$  and uncountable.

PROOF: Construct a new family  $\mathcal{G} := \{G_f \mid f \in \mathcal{F}\}$  of set-valued mappings, where  $G_f : M \rightarrow \mathcal{P}_0(\mathbb{R})$  is defined by

$$G_f(x) := \begin{cases} [f(x), \infty[ & \text{if } s \in ]0, 1[ \\ [f(x) - f(o_X), \infty[ & \text{if } s = 1. \end{cases}$$

It is immediately seen that  $\mathcal{G}$  is a family of upper semicontinuous  $(A, s)$ -convex mappings from  $M$  to  $\mathcal{P}_0(\mathbb{R})$  which is not equicontinuous at  $o_X$  and satisfies the condition

$$\frac{1}{\lambda}G(o_X) \subseteq G(o_X) \subseteq \lambda G(o_X) \text{ for all } G \in \mathcal{G} \text{ and all } \lambda \in ]0, 1].$$

On the other hand, it is clear that  $\mathcal{F}$  is bounded at a point  $x_0 \in M$  if and only if  $\mathcal{G}$  is bounded at  $x_0$ . Hence  $WS_{\mathcal{F}} = WS_{\mathcal{G}}$ . Therefore, the assertions of the theorem are consequences of Theorem 3.2.  $\square$

**Theorem 4.2.** *Let  $X$  be a topological linear space of second category, let  $s \in ]0, 1]$ , and let  $\mathcal{F}$  be a pointwise bounded family of lower semicontinuous  $(A, s)$ -convex functions from a set  $M \in \mathcal{O}_{ac}(X)$  to  $\mathbb{R}$ . Then  $\mathcal{F}$  is equicontinuous on  $M$ .*

PROOF: Let  $x_0$  be any point in  $M$ . We choose a balanced neighbourhood  $U$  of  $x_0$  such that  $x_0 + U \subseteq M$ . Then  $M_0 := \text{int}(\text{conv} U)$  lies in  $\mathcal{O}_{ac}(X)$  and satisfies  $x_0 + M_0 \subseteq M$ . To each  $f \in \mathcal{F}$  we assign the function  $g_f : M_0 \rightarrow \mathbb{R}$  defined by  $g_f(x) := f(x_0 + x)$ . Obviously, all the functions  $g_f$  ( $f \in \mathcal{F}$ ) are lower semicontinuous and  $(A, s)$ -convex. Moreover, the family  $\mathcal{G} := \{g_f \mid f \in \mathcal{F}\}$  is bounded on  $M_0$ . Therefore, in view of Theorem 4.1, the family  $\mathcal{G}$  must be equicontinuous at  $o_X$ . In other words,  $\mathcal{F}$  must be equicontinuous at  $x_0$ .  $\square$

**5. Singularities and uniform boundedness of families of closed convex processes**

Let  $X$  and  $Y$  be normed linear spaces. In the sequel we shall denote by  $B_X(r)$  (respectively by  $B_Y(r)$ ) the closed ball centered at  $o_X$  (respectively at  $o_Y$ ) and of radius  $r$ . The balls  $B_X(1)$  and  $B_Y(1)$  will be simply denoted by  $B_X$  and  $B_Y$ , respectively.

A mapping  $F : X \rightarrow 2^Y$  is called a *convex process* if the following conditions are satisfied:

- (i)  $o_Y \in F(o_X)$ ;
- (ii)  $F(\lambda x) = \lambda F(x)$  for all  $x \in X$  and all  $\lambda \in ]0, \infty[$ ;
- (iii)  $F(x_1) + F(x_2) \subseteq F(x_1 + x_2)$  for all  $x_1, x_2 \in X$ .

The norm  $\|F\|$  of a convex process  $F$  is defined by

$$\|F\| := \sup_{x \in \text{Dom}(F) \cap B_X} \inf_{y \in F(x)} \|y\|.$$

The mapping  $F$  is called a *closed convex process* if it is a convex process whose graph is closed.

**Proposition 5.1.** *Let  $X$  and  $Y$  be normed linear spaces, and let  $\mathcal{F}$  be a family of convex processes from  $X$  to  $\mathcal{P}_0(Y)$ . Then the following assertions are equivalent:*

- 1°  $\mathcal{F}$  is equicontinuous at  $o_X$ ;
- 2° there exists a real number  $k$  such that  $\|F\| \leq k$  for all  $F \in \mathcal{F}$ .

PROOF: 1°  $\Rightarrow$  2° Since  $\mathcal{F}$  is equicontinuous at  $o_X$ , there exists a positive real number  $r$  such that

$$F(o_X) \subseteq F(x) + B_Y \quad \text{for all } F \in \mathcal{F} \text{ and all } x \in B_X(r).$$

We shall prove that

$$(5.1) \quad \|F\| \leq \frac{1}{r} \quad \text{for all } F \in \mathcal{F}.$$

Let  $F$  be any mapping in  $\mathcal{F}$ , and let  $x$  be any point in  $B_X$ . Because of  $\|rx\| \leq r$ , we have

$$F(o_X) \subseteq F(rx) + B_Y = rF(x) + B_Y.$$

Since  $o_Y \in F(o_X)$ , we can choose the points  $y_0 \in F(x)$  and  $z_0 \in B_Y$  such that  $ry_0 + z_0 = o_Y$ . Hence  $\|y_0\| = 1/r\|z_0\| \leq 1/r$ . Therefore we have

$$\inf_{y \in F(x)} \|y\| \leq \frac{1}{r}.$$

Since  $x$  was arbitrarily chosen in  $B_X$ , we can conclude that  $\|F\| \leq 1/r$ . Consequently (5.1) holds.

$2^\circ \Rightarrow 1^\circ$  Let  $V$  be any neighbourhood of  $o_Y$ . We choose a positive real number  $r$  such that  $B_Y(r) \subseteq V$ . Put  $\alpha := \frac{k+1}{r}$  and  $U := \frac{1}{\alpha}B_X$ . We shall prove that

$$(5.2) \quad F(o_X) \subseteq F(x) + V \quad \text{and} \quad F(x) \subseteq F(o_X) + V$$

whenever  $F \in \mathcal{F}$  and  $x \in U$ .

Let  $F$  be any mapping in  $\mathcal{F}$ , and let  $x$  be any point in  $U$ . Then  $\alpha x \in B_X$ . Since  $\|F\| \leq k$ , we can find a point  $y_0 \in F(\alpha x)$  such that  $\|y_0\| \leq k+1$ . Hence we can find a point  $z_0 \in F(x) \cap B_Y(r)$  such that  $y_0 = \alpha z_0$ . Then

$$z_0 + F(o_X) \subseteq F(x) + F(o_X) \subseteq F(x).$$

Consequently

$$F(o_X) \subseteq F(x) - z_0 \subseteq F(x) + B_Y(r) \subseteq F(x) + V.$$

On the other hand, we have  $-\alpha x \in B_X$ . Hence we can find a point  $y'_0 \in F(-\alpha x)$  such that  $\|y'_0\| \leq k+1$ . Therefore we can find a point  $z'_0 \in F(-x) \cap B_Y(r)$  such that  $y'_0 = \alpha z'_0$ . We have

$$z'_0 + F(x) \subseteq F(-x) + F(x) \subseteq F(o_X).$$

Hence

$$F(x) \subseteq F(o_X) - z'_0 \subseteq F(o_X) + B_Y(r) \subseteq F(o_X) + V.$$

Consequently (5.2) holds as claimed. Therefore  $\mathcal{F}$  is equicontinuous at  $o_X$ .  $\square$

**Theorem 5.2.** *Let  $X$  be a Banach space, let  $Y$  be a normed linear space, and let  $\mathcal{F}$  be a family of closed convex processes from  $X$  to  $\mathcal{P}_0(Y)$  such that*

$$\sup \{\|F\| \mid F \in \mathcal{F}\} = \infty.$$

*Then  $WS_{\mathcal{F}}$  is an uncountable residual subset of  $X$ .*

PROOF: For every  $F \in \mathcal{F}$  define the function  $f_F : X \rightarrow \mathbb{R}$  by

$$f_F(x) := \inf_{y \in F(x)} \|y\|.$$

Since  $F$  is a closed convex process, it is easily seen that  $f_F$  is a sublinear lower semicontinuous function. Put  $\mathcal{G} := \{f_F \mid F \in \mathcal{F}\}$ . Obviously  $\mathcal{G}$  is bounded at  $o_X$  because  $f_F(o_X) = 0$  for all  $F \in \mathcal{F}$ .

Suppose that  $\mathcal{G}$  is equicontinuous at  $o_X$ . Then we can find a positive real number  $\delta$  such that  $f_F(x) < 1$  for all  $x \in B_X(\delta)$  and all  $F \in \mathcal{F}$ . Choose  $F_0 \in \mathcal{F}$  such that  $\|F_0\| > 2/\delta$ . Then we can find a point  $x_0^* \in B_X$  such that

$$\inf_{y \in F_0(x_0^*)} \|y\| > \frac{2}{\delta}.$$

Therefore we have  $\|y\| > 2/\delta$  for all  $y \in F_0(x_0^*)$ . Put  $x_0 := \delta x_0^*$ . Then  $x_0 \in B_X(\delta)$  and  $\|y\| > 2$  for all  $y \in F_0(x_0)$ . Hence  $f_{F_0}(x_0) \geq 2$ , which is a contradiction. Summing up, we conclude that the family  $\mathcal{G}$  is not equicontinuous at  $o_X$ .

Since  $WS_{\mathcal{F}} = WS_{\mathcal{G}}$ , the conclusion of the theorem follows immediately from Theorem 4.1.  $\square$

The next result is Theorem 2.3.1 in [1].

**Theorem 5.3.** *Let  $X$  be a Banach space, let  $Y$  be a normed linear space, and let  $\mathcal{F}$  be a pointwise bounded family of closed convex processes from  $X$  to  $\mathcal{P}_0(Y)$ . Then*

$$\sup \{\|F\| \mid F \in \mathcal{F}\} < \infty.$$

PROOF: Follows immediately from Theorem 5.2.  $\square$

The following theorem is similar to Theorem 5.3, but upper semicontinuous convex processes are considered instead of closed convex ones.

**Theorem 5.4.** *Let  $X$  be a Banach space, let  $Y$  be a normed linear space, and let  $\mathcal{F}$  be a pointwise bounded family of upper semicontinuous convex processes from  $X$  to  $\mathcal{P}_0(Y)$ . Then*

$$\sup \{\|F\| \mid F \in \mathcal{F}\} < \infty.$$

PROOF: Follows immediately from Theorem 3.3 and Proposition 5.1.  $\square$

**Acknowledgement.** The author expresses his gratitude to Professor Wolfgang W. Breckner for the suggestions that helped to improve the initial version of this paper.

#### REFERENCES

- [1] Aubin J.P., Frankowska H., *Set-Valued Analysis*, Birkhäuser, Boston, 1990.
- [2] Breckner W.W., *Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen*, Publ. Inst. Math. (Beograd) **23 (37)** (1978), 13–20.
- [3] Breckner W.W., *A principle of condensation of singularities for set-valued functions*, Mathematica-Rev. Anal. Numér. Théor. Approx., Sér. L'Anal. Numér. Théor. Approx. **12** (1983), 101–111.

- [4] Breckner W.W., *Continuity of generalized convex and generalized concave set-valued functions*, Rev. Anal. Numér. Théor. Approx. **22** (1993), 39–51.
- [5] Breckner W.W., Trif T., *On the singularities of certain families of nonlinear mappings*, Pure Math. Appl. **6** (1995), 121–137.

UNIVERSITATEA BABEȘ-BOLYAI, FAC. DE MATEMATICĂ ȘI INFORMATICĂ, STR. KOGĂLNICEANU NR. 1, RO-3400 CLUJ-NAPOCA, ROMANIA

*E-mail:* ttrif@math.ubbcluj.ro

(Received May 12, 1997)