

S. A. Peregudov

On n -in-countable bases

Commentationes Mathematicae Universitatis Carolinae, Vol. 41 (2000), No. 1, 175--178

Persistent URL: <http://dml.cz/dmlcz/119151>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2000

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On n -in-countable bases

S.A. PEREGUDOV

Abstract. Some results concerning spaces with countably weakly uniform bases are generalized for spaces with n -in-countable ones.

Keywords: weakly uniform base, n -in-countable base, countably compact space, separable space

Classification: Primary 54A25; Secondary 54D30

All spaces in the paper are assumed to be T_1 . If τ and λ are cardinals, then one says that a family \mathcal{B} of sets is τ -in- λ if for every set A of cardinality τ , $A \subset B$ holds for no more than λ members $B \in \mathcal{B}$. τ -in- $< \lambda$ is defined similarly. One says that a family is τ -in-countable in place of τ -in- ω and that a family is τ -in-finite in place of τ -in- $< \omega$. A 2-in-finite family is called *weakly uniform* and a 2-in-countable family is also called *countably weakly uniform*.

We are going to extend results of paper [6] on spaces with *countably weakly uniform bases* to spaces with *n -in-countable bases*, where n is a natural number. A few results concern *n -in-finite bases*.

Lemma 1. *Let $n \geq 1$ be a natural number. An n -in-countable open family in a separable space is countable.*

PROOF: It is evident. □

Theorem 1. *Let $n \geq 1$ be a natural number. A regular countably compact space with an n -in-countable T_1 -separating open cover is metrizable.*

PROOF: Let X be a regular countably compact space and let \mathcal{B} be its n -in-countable T_1 -separating open cover. Denote by $\mathcal{L}(X)$ the set of all nonisolated points of X . It is evident that if $x \in \mathcal{L}(X)$ then $ord(x, \mathcal{B}) \leq \min\{|A| : x \text{ is an accumulation point of } A\}$. Let us consider the set $Z = \{x \in \mathcal{L}(X) : ord(x, \mathcal{B}) \leq \omega\}$. Because X is regular countably compact, $\bar{Z} = \mathcal{L}(X)$. Now, by analogy with the proof of Miscenco’s theorem [3] with using Lemma 1 one can prove that there exists a countable set $Y \subset Z$ such that $\bar{Y} = \mathcal{L}(X)$. Hence $|\{B \in \mathcal{B} : B \cap \mathcal{L}(X) \neq \emptyset\}| \leq \omega$. We can assume that if $B \in \mathcal{B}$ and $B \cap \mathcal{L}(X) = \emptyset$ then B is an one-point set. Hence \mathcal{B} is countable. Therefore X is metrizable. □

Corollary 1 ([1]). *Let $n \geq 1$ be a natural number. A regular countably compact space with an n -in-countable base is metrizable.*

Later on we will denote by $\mathcal{I}(X)$ the set of all isolated points of a space X .

Lemma 2. *Let $n \geq 1$ be a natural number. Let X be a space with an n -in-countable base \mathcal{B} and let $|\mathcal{I}(X)| = \tau$, where τ is an uncountable cardinal. Then X has an n -in-countable base \mathcal{B}^* such that for each $x \in \mathcal{I}(X)$, $\text{ord}(x, \mathcal{B}^*) < \tau$.*

PROOF: Let $\mathcal{I}(X) = \{x_\xi : \xi < \tau\}$. Let us define $\mathcal{B}_0 = \{B \in \mathcal{B} : x_0 \in B\}$. Let $\eta < \tau$. Assume that \mathcal{B}_ξ are defined for each $\xi < \eta$. Then we define $\mathcal{B}_\eta = \{B \in \mathcal{B} : x_\eta \in B, B \notin \mathcal{B}_\xi \text{ under } \xi < \eta\}$. Now put $\mathcal{B}_\xi^1 = \{B \setminus \{x_\xi\} : B \in \mathcal{B}_\xi\}$ for each $\xi < \tau$ and $\mathcal{B}^1 = \bigcup \{\mathcal{B}_\xi^1 : \xi < \tau\}$. It is evident that if $A \subset \mathcal{I}(X)$ and $|A| = n - 1$ then $|\{B \in \mathcal{B}^1 : A \subset B\}| < \tau$. Repeating this step $n - 1$ times (of course we assume $n \geq 2$), we obtain a family $\mathcal{B}^{(n-1)}$ such that $\mathcal{B}^* = \mathcal{B}^{(n-1)} \cup \{B \in \mathcal{B} : B \cap \mathcal{I}(X) = \emptyset\} \cup \{\{x\} : x \in \mathcal{I}(X)\}$ is a base with all the required properties. \square

Lemma 3. *Let $n \geq 1$ be a natural number. Let X be a space with an n -in-countable base \mathcal{B} and let $|\mathcal{I}(X)| \leq \omega$. Then X has an n -in-countable base \mathcal{B}^* such that for each $x \in \mathcal{I}(X)$, $\text{ord}(x, \mathcal{B}^*) \leq \omega$.*

PROOF: The statement can be proved just as Lemma 2. \square

Theorem 2. *Let $n \geq 1$ be a natural number. A regular locally separable space with an n -in-countable base is metrizable.*

PROOF: The statement is proved in just the same way as Theorem from [7] with using above-mentioned Lemmas 1, 2, and 3. Here is a sketch of the proof. By Lemma 1 the space is locally metrizable. Hence every its n -in-countable base has a countable order at each nonisolated point. Let us maintain induction on the cardinality of the set of all isolated points. Let a space X satisfy the assumptions of Theorem 2, and let $|\mathcal{I}(X)| \leq \omega$. Then with respect to the above remark and by Lemma 3, X has a point-countable base. Therefore X is divided into a disjoint family of open metrizable subspaces. Consequently, X is metrizable. Now, let τ be an uncountable cardinal. Let us assume that the statement of the theorem is true for each cardinal $\lambda < \tau$. Suppose now that $|\mathcal{I}(X)| = \tau$. We will consider two cases.

Case 1. τ is regular. By Lemmas 1 and 2 there is a base \mathcal{B} of X of order $< \tau$ at each point of X . Without loss of generality we may assume that every element of the base is separable. For every $B^* \in \mathcal{B}$ define a family $\mathcal{E}(B^*)$ in the following way: $\mathcal{E}_0(B^*) = \{B^*\}$, $\mathcal{E}_1(B^*) = \{B \in \mathcal{B} : B \cap B^* \neq \emptyset\}$, \dots , $\mathcal{E}_n(B^*) = \{B \in \mathcal{B} : B \cap (\bigcup \mathcal{E}_{n-1}) \neq \emptyset\}$, etc., $\mathcal{E}(B^*) = \bigcup \{\mathcal{E}_n(B^*) : n \in \omega\}$. Then we have $|\mathcal{E}(B^*)| < \tau$, and $\bigcup \mathcal{E}(B^*)$ has less than τ isolated points. It follows that $\bigcup \mathcal{E}(B^*)$ is metrizable, hence X is metrizable too.

Case 2. τ is singular. Then there are a cardinal $\lambda < \tau$ and a partition $\{I_\xi : \xi < \lambda\}$ of the set $\mathcal{I}(X)$ such that $|I_\xi| = \tau_\xi < \tau$. Let \mathcal{B} be a base of X with the same properties as in Case 1. Fix an ordinal $\xi < \lambda$. For each point $a \in I_\xi$ put $\mathcal{S}_a = \{B \in \mathcal{B} : a \in B\}$. Put $A_\eta = \bigcup \{\mathcal{S}_a : |\mathcal{S}_a| < \tau_\eta\}$. Since A_η is an open subspace of X , which has the set of all isolated points of cardinality less than τ , it is metrizable. Hence, the space $G_\xi = \bigcup \{B \in \mathcal{B} : B \cap I_\xi \neq \emptyset\} = \bigcup \{A_\eta : \eta < \lambda\}$

has a base of order no more than λ at each its point. It follows that the space $G = \bigcup\{G_\xi : \xi < \lambda\} = \bigcup\{B \in \mathcal{B} : B \cap \mathcal{I}(X) \neq \emptyset\}$ has a base \mathcal{U} of order no more than λ at each its point. Without loss of generality we may assume that every element of the base is separable. In the same way as in Case 1 it can be proved that G is metrizable. Because $X \setminus G$ is contained in an open subspace of X without isolated points, it is metrizable. Thus, X is the union of two open locally separable metrizable subspaces. Hence X is metrizable. \square

Corollary 2. *Let $n \geq 1$ be a natural number. A regular space which admits an n -in-countable cover of open separable metrizable subspaces is metrizable.*

Corollary 3. *Let $n \geq 1$ be a natural number. A regular locally countably compact space with an n -in-countable base is metrizable.*

Corollary 4 ([2]). *Let $n \geq 1$ be a natural number. A Hausdorff locally compact space with an n -in-countable base is metrizable.*

Lemma 4 (MA + \neg CH). *Let $n \geq 1$ be a natural number. If X is a Čech-complete space with $c(X) = \omega$, then every open n -in-countable family of X is countable.*

PROOF: Under $n = 1$ this is a result of Shapirovskii [9]. Then the statement is proved by induction on n . \square

It follows from [8] that the statement of Lemma 4 is false under \neg SH even if $n = 1$.

Theorem 3 (MA + \neg CH). *Let $n \geq 1$ be a natural number. Suppose that X is a regular space which is locally Čech-complete and locally has the Souslin property. If X has an n -in-countable base, then X is metrizable.*

PROOF: The statement follows from Lemma 4 and Theorem 2. \square

Theorem 4. *Let $n \geq 1$ be a natural number. Every pseudocompact space with an n -in-finite base is Čech complete first countable.*

PROOF: Let X be a pseudocompact space and \mathcal{B} be an n -in-finite base for X . For each $B \in \mathcal{B}$ choose an open in βX set B' such that $B = X \cap B'$, and consider the family \mathcal{B}' consisting of such sets. We show that the family \mathcal{B}' is n -in-finite in βX . Let $A \subset \beta X$ and let $|A| = n$. Let \mathcal{S} be an infinite countable subfamily of the family \mathcal{B} . Denote by F the set $\bigcap \mathcal{S} \cap A$. By definition, the cardinality of F is less than n . Then $G = \bigcap\{B' \setminus F : B \in \mathcal{S}\} \subset \beta X \setminus X$. Because X is pseudocompact, the set G being a G_δ -set of βX that is contained in $\beta X \setminus X$ is empty by the well known result of Hewitt [5]. Thus \mathcal{B}' is an n -in-finite family in βX . The same arguments show that the family \mathcal{B}' has finite order at each point of $\beta X \setminus X$. Denote by \mathcal{I} the set of all isolated points of X and define $F_m = \{x \in \beta X : ord(x, \mathcal{B}') \leq m\} \setminus \mathcal{I}$. Because \mathcal{I} is open in βX , each F_m is a closed subset of βX . Moreover, $\beta X \setminus X = \bigcup\{F_m : m \in \omega\}$. So X is Čech

complete; therefore, X is a k -space. It is evident that a k -space with an n -infinite base is first countable. \square

Corollary 5. *Let $n \geq 1$ be a natural number. Every submetacompact pseudo-compact space with an n -infinite base is metrizable.*

Theorem 5. *Let $n \geq 2$ be a natural number. Every space X with an n -infinite base has cardinality at most $\exp_{n-1}(L(X))$, where $L(X)$ is the Lindelöf degree of X .*

PROOF: Let \mathcal{B} be an n -infinite base of X . Put $\tau = L(X)$. We will use the theorem of Erdős and Rado: $(\exp_{n-1}(\tau))^+ \rightarrow (\tau^+)_\tau^n$ ([4]). Let us assume that $|X| > \exp_{n-1}(\tau)$. Consider a mapping $P : [X]^n \rightarrow \omega$ defined by the rule: $A \mapsto |\{B \in \mathcal{B} : A \subset B\}|$. There exists a homogeneous with respect to P set H of cardinality τ^+ . It is easy to show that the set H is a closed discrete subset of X , a contradiction. \square

Let us note that there exists an example of a Hausdorff Lindelöf space with a weakly uniform base which is not first countable ([6]).

REFERENCES

- [1] Arhangel'skii A.V., Just W., Reznichenko E., Szeptyczki P.J., *Sharp bases and weakly uniform bases versus point countable bases*, to appear.
- [2] Balogh Z., Gruenhage G., *Base multiplicity in compact and generalized compact spaces*, to appear.
- [3] Engelking R., *General Topology*, Warsaw, Heldermann Verlag, Berlin, 1989.
- [4] Erdős P., Rado R., *A partition calculus in set theory*, Bull. Amer. Math. Soc. **62** (1956), 427–489.
- [5] Hewitt E., *Rings of real-valued continuous functions, 1*, Trans. Amer. Math. Soc. **64** (1948), 45–99.
- [6] Peregudov S.A., *Weakly uniform bases and the first axiom of countability* (in Russian), Math. Zametki **3** (1986), 331–340.
- [7] Peregudov S.A., *On metrizability in a class of topological spaces with a weakly uniform base* (in Russian), Bull. Pol. Acad. Nauk **28** (1980), 609–612.
- [8] Ponomarev V.I., *On metrizability of finally compact spaces with a point countable base* (in Russian), Dokl. Akad. Nauk SSSR **174** (1967), 1274–1277.
- [9] Shapirovskii B.E., *On separability and metrizability of spaces with Souslin condition* (in Russian), Dokl. Akad. Nauk SSSR **207** (1972), 800–803.

STATE UNIVERSITY OF MANAGEMENT, RJAZANSKIY PROSPEKT 99, MOSCOW, RUSSIA 109 542
E-mail: matem@acman.msk.su

(Received March 3, 1999, revised July 22, 1999)