

Michael K. Kinyon

Global left loop structures on spheres

Commentationes Mathematicae Universitatis Carolinae, Vol. 41 (2000), No. 2, 325--346

Persistent URL: <http://dml.cz/dmlcz/119167>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2000

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Global left loop structures on spheres

MICHAEL K. KINYON

Abstract. On the unit sphere \mathbb{S} in a real Hilbert space \mathbf{H} , we derive a binary operation \odot such that (\mathbb{S}, \odot) is a power-associative Kikkawa left loop with two-sided identity \mathbf{e}_0 , i.e., it has the left inverse, automorphic inverse, and A_l properties. The operation \odot is compatible with the symmetric space structure of \mathbb{S} . (\mathbb{S}, \odot) is not a loop, and the right translations which fail to be injective are easily characterized. (\mathbb{S}, \odot) satisfies the left power alternative and left Bol identities “almost everywhere” but not everywhere. Left translations are everywhere analytic; right translations are analytic except at $-\mathbf{e}_0$ where they have a nonremovable discontinuity. The orthogonal group $O(\mathbf{H})$ is a semidirect product of (\mathbb{S}, \odot) with its automorphism group. The left loop structure of (\mathbb{S}, \odot) gives some insight into spherical geometry.

Keywords: loop, quasigroup, sphere, Hilbert space, spherical geometry

Classification: 20N05

1. Introduction

Let \mathbf{H} be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$. Let

$$(1.1) \quad \mathbb{S} = \{\mathbf{x} \in \mathbf{H} : \langle \mathbf{x}, \mathbf{x} \rangle = 1\}$$

be the unit sphere in \mathbf{H} . For $\mathbf{x}, \mathbf{y} \in \mathbb{S}$, set

$$(1.2) \quad \mathbf{x} * \mathbf{y} = 2\langle \mathbf{x}, \mathbf{y} \rangle \mathbf{x} - \mathbf{y}.$$

([16, p. 66]). If \mathbf{x} and \mathbf{y} span a plane Π through the origin, then $\mathbf{x} * \mathbf{y}$ is a point lying in $\mathbb{S} \cap \Pi$ which is obtained by reflecting \mathbf{y} in Π about the line passing through \mathbf{x} and the origin. Equivalently, thinking of $\mathbb{S} \cap \Pi$ as the great circle passing through \mathbf{x} and \mathbf{y} , $\mathbf{x} * \mathbf{y}$ is the point on $\mathbb{S} \cap \Pi$ whose angle with \mathbf{x} is the same as the angle between \mathbf{x} and \mathbf{y} , but with the opposite orientation. The magma $(\mathbb{S}, *)$ has the structure of a symmetric space.

Definition 1.1. A *symmetric space* $(M, *)$ is a topological space M together with a continuous binary operation $* : M \times M \rightarrow M$ such that the following properties hold. For $x, y, z \in M$,

- (L1) (idempotence) $x * x = x$;
- (L2) (left keyes identity) $x * (x * y) = y$;
- (L3) (left distributivity) $x * (y * z) = (x * y) * (x * z)$;
- (L4) each $a \in M$ has a neighborhood $U \subseteq M$ such that $a * x = x$ implies $x = a$ for all $x \in U$.

We make the implicit assumption that the binary operation $*$: $M \times M \rightarrow M$ possesses as much smoothness as the space M allows. Thus if M is a smooth manifold, $*$ is assumed to be smooth, and not just continuous. In the smooth case, Definition 1.1 is due to Loos [16]. It is equivalent to the more standard ones (cf. [7]), but has the advantage of depending only on the topology. This motivates our trivial adaptation of his definition.

The magma $(\mathbb{S}, *)$ defined by (1.2) is easily seen to satisfy (L1), (L2) and (L3). Also, $\mathbf{x} * \mathbf{y} = \mathbf{y}$ if and only if $\mathbf{y} = \pm \mathbf{x}$, and thus (L4) holds.

Observe that the first three axioms of a symmetric space are purely algebraic. For $x \in M$, we define a mapping $s_x : M \rightarrow M$, called the *symmetry about x* , by

$$(1.3) \quad s_x(y) = x * y.$$

In terms of the symmetries, the key identity (L2) reads $s_x^2 = I$, where I is the identity mapping on M . Therefore $(M, *)$ is a left quasigroup, i.e., each symmetry s_x is a bijection. The topological axiom (L4) asserts that each $a \in M$ is an isolated fixed point of the symmetry s_a .

As another example of a symmetric space, let G be a topological group with the property that $1 \in G$ has a neighborhood $U \subseteq G$ such that no element of U has order 2. (For instance, if G is a Lie group, then the existence of such a neighborhood is implied by the existence of the exponential map.) Define

$$(1.4) \quad x * y = xy^{-1}x$$

for $x, y \in G$. Axioms (L1), (L2) and (L3) are easily checked. Now $1 * x = x^{-1}$ for all $x \in G$, and thus the assumed property of G implies that (L4) holds for $a = 1$. In addition, we have $x(y * z) = xy * xz$ for $x, y, z \in G$ ([16, p. 65]). This implies that (L4) holds for all $a \in G$.

The case where G is a (multiplicative) abelian group is of particular interest here; in this case the symmetric space operation is simply $x * y = x^2y^{-1}$ for $x, y \in G$. Now assume G is uniquely 2-divisible, i.e., that the squaring map is bijective. For $x \in G$, let $x^{1/2}$ denote the unique element of G whose square is x . Then one can recover the group product from the symmetric space operation by defining

$$(1.5) \quad x \cdot y = x^{1/2} * (1 * y)$$

for all $x, y \in G$. Thus if G is uniquely 2-divisible, (G, \cdot) and $(G, *)$ are isotopic. It is clear from (1.5) that the two essential ingredients for the isotopy are: a distinguished point $1 \in G$ and a well-defined notion of " $x^{1/2}$ " for all $x \in G$. It is useful to note that the latter ingredient can be described entirely in terms of the symmetric space operation. If G is a uniquely 2-divisible abelian group, then *for $x \in G$, there exists a unique $x^{1/2} \in G$ such that $x^{1/2} * 1 = x$.*

This idea can be generalized to a wider class of symmetric spaces.

Lemma 1.2. *Let $(M, *)$ be a left keyesian, left distributive, left quasigroup, and fix $e \in M$. The following properties are equivalent.*

1. *For each $x \in M$, there exists a unique $x^{1/2} \in M$ such that $x^{1/2} * e = x$.*
2. *$(M, *)$ is a right quasigroup.*

PROOF: (1) \implies (2): Let $a, b \in G$ be given, and assume $x \in M$ satisfies $x * a = b$. Then using (L3) and (L2),

$$a^{1/2} * b = a^{1/2} * (x * a) = (a^{1/2} * x) * (a^{1/2} * a) = (a^{1/2} * x) * e.$$

By the uniqueness assumption, $a^{1/2} * x = (a^{1/2} * b)^{1/2}$, and thus by (L2), $x = a^{1/2} * (a^{1/2} * b)^{1/2}$. Conversely, set $x = a^{1/2} * (a^{1/2} * b)^{1/2}$. Then using (L3) and (L2),

$$\begin{aligned} x * a &= \left(a^{1/2} * (a^{1/2} * b)^{1/2} \right) * (a^{1/2} * e) \\ &= a^{1/2} * \left((a^{1/2} * b)^{1/2} * e \right) \\ &= a^{1/2} * (a^{1/2} * b) \\ &= b. \end{aligned}$$

Therefore $(M, *)$ is a right quasigroup and hence a quasigroup.

(2) \implies (1): Obvious. □

Lemma 1.3. *Let $(M, *)$ be a left distributive quasigroup. Then $(M, *)$ is idempotent. Moreover, for each $a \in M$, $a * x = x$ implies $x = a$ for all $x \in M$.*

PROOF: For all $x, a \in M$, $(x * x) * a = x * (x * (x \setminus a)) = x * a$. Cancelling a , we have $x * x = x$, and thus $(M, *)$ is idempotent. If $a * x = x = x * x$, then clearly $x = a$. □

These considerations lead us to the following definition.

Definition 1.4. A reflection quasigroup $(M, *)$ is a left keyesian, left distributive quasigroup.

The term “reflection quasigroup” is due to Kikkawa [10]. Reflection quasigroups are also known as “left-sided quasigroups”, following a convention of Robinson [21]. Reflection quasigroups are also equivalent to the recently studied “point reflection structures” of Gabrieli and Karzel [3], [4], [5]. Given a nonempty set P and a mapping $\tilde{\cdot} : P \rightarrow \text{Sym}(P) : x \mapsto \tilde{x}$, Gabrieli and Karzel call the pair $(P, \tilde{\cdot})$ a “point-reflection structure” if the following hold: (i) $\forall x, y \in P$, $(\tilde{x} \circ \tilde{x})(y) = y$; (ii) $\forall x, y \in P$, $\tilde{x}(y) = y$ implies $y = x$; (iii) $\forall a, b \in P$, $\exists! x \in P$ such that $\tilde{x}(a) = b$; (iv) $\forall a, b \in P$, $\exists c \in P$ such that $\tilde{a} \circ \tilde{b} \circ \tilde{a} = \tilde{c}$. This is simply the notion of a reflection quasigroup in different notation. Indeed, for $x, y \in P$,

set $x * y = \tilde{x}(y)$. Then $(P, *)$ is left keyesian by (i), idempotent by (ii), and is a right quasigroup by (iii). Applying both sides of (iv) to $a * b$, and using the left keyesian identity and idempotence, we have $a * b = c * (a * b)$. By idempotence and the fact that $(P, *)$ is a right quasigroup, $c = a * b$. Thus applying (iv) to $a * x$, and using the left keyesian identity, we have $a * (b * x) = (a * b) * (a * x)$ for all $a, b, x \in P$. This is left distributivity. Conversely, if $(P, *)$ is a reflection quasigroup, then for $x, y \in P$, set $\tilde{x}(y) = x * y$. Properties (i), (ii), and (iii) are clear. For all $a, b, x \in P$, we have $(\tilde{a} \circ \tilde{b} \circ \tilde{a})(x) = a * (b * (a * x)) = (a * b) * x$, using the left distributive and left keyesian properties. Thus (iv) holds with $c = a * b$.

Lemma 1.3 implies that a topological reflection quasigroup is necessarily a symmetric space. However, the symmetric space $(\mathbb{S}, *)$ is not a reflection quasigroup. For instance, given $\mathbf{a} \in \mathbb{S}$, the equation $\mathbf{x} * \mathbf{a} = \mathbf{a}$ has two solutions $\mathbf{x} = \pm \mathbf{a}$. The obstructions to $(\mathbb{S}, *)$ being a reflection quasigroup can also be seen in geometric terms. If $\dim \mathbf{H} > 2$, then for a fixed $\mathbf{e} \in \mathbb{S}$, there is a nontrivial family of great circles on \mathbb{S} which connect \mathbf{e} to its antipode $-\mathbf{e}$. Thus consider the equation

$$(1.6) \quad \mathbf{x} * \mathbf{e} = -\mathbf{e}$$

in the symmetric space $(\mathbb{S}, *)$. We have $-\mathbf{e} = 2\langle \mathbf{e}, \mathbf{x} \rangle \mathbf{x} - \mathbf{e}$, or $\langle \mathbf{e}, \mathbf{x} \rangle \mathbf{x} = \mathbf{0}$. Hence (1.6) is equivalent to

$$(1.7) \quad \langle \mathbf{e}, \mathbf{x} \rangle = 0.$$

Therefore any point $\mathbf{x} \in \mathbb{S}$ which is orthogonal to \mathbf{e} will satisfy (1.6). But in general, there is no distinguished solution to (1.6). (This nonuniqueness problem holds even for the unit circle $S^1 \subset \mathbb{R}^2$. However, since there are only two solutions, a choice based on orientation leads to the usual group structure of S^1 .) Despite these concerns, reflection quasigroups form a model for our later discussion of $(\mathbb{S}, *)$.

The aforementioned obstructions extend to other compact symmetric spaces. Indeed, in a smooth symmetric space, each symmetry s_x is locally a geodesic symmetry; i.e., for y near x , $s_x(y)$ is the image of y under the reflection about x along a (maximal) geodesic connecting x and y . For a distinguished point e and any other point x in a smooth reflection quasigroup (such as a Riemannian symmetric space of noncompact type), there is only one geodesic connecting e and x , and thus there is a unique point $x^{1/2}$ on that geodesic such that $x^{1/2} * e = s_{x^{1/2}}(e) = x$. However, in compact symmetric spaces, the distinguished point e (or any point for that matter) has a nonempty cut locus, i.e., set of points conjugate to e ([13]). For each such conjugate point x , there is a nontrivial family of distinct geodesics connecting e and x , and thus there is a nontrivial family of distinct points z such that $z * e = x$. Since there is generally no distinguished choice for “ $x^{1/2}$ ”, compact smooth symmetric spaces are generally not reflection quasigroups.

Let $(M, *)$ be a reflection quasigroup, fix a distinguished element $e \in M$, and as before, for each $x \in M$, let $x^{1/2} = x/e$, where $/$ denotes right division in

$(M, *)$. (Since a reflection quasigroup is left keyesian, left division \setminus agrees with the multiplication $*$.) Define

$$(1.8) \quad x \cdot y = x^{1/2} * (e * y)$$

for $x, y \in M$. Then (M, \cdot) is a loop with identity element e , and in fact, (M, \cdot) is the principal e, e -isotope of $(M, *)$ ([1], [19]): $x \cdot y = (x/e) * (e \setminus y)$. Following some definitions, we will identify the class of loops to which (M, \cdot) belongs.

In a left loop, denote the left translations by $L_x : y \mapsto x \cdot y$, and the left inner mappings by $L(x, y) = L_{x \cdot y}^{-1} L_x L_y$. A left loop is said to have the *left inverse property* if

$$(LIP) \quad L_x^{-1} = L_{x^{-1}}$$

for all x , where x^{-1} is the right (and necessarily, two-sided) inverse of x . A left loop is said to have the A_l *property* if every left inner mapping is an automorphism. A left loop with two-sided inverses is said to have the *automorphic inverse property* if

$$(AIP) \quad (x \cdot y)^{-1} = x^{-1} \cdot y^{-1}$$

for all x, y . An A_l , LIP, AIP left loop is called a *Kikkawa* left loop ([9]). A loop is called a (left) *Bol loop* if it satisfies the (left) *Bol identity*:

$$(Bol) \quad L_x L_y L_x = L_{x \cdot (y \cdot x)}$$

for all x, y, z . Bol loops necessarily satisfy LIP (see, e.g., [9, 6.4]). A Bol loop with AIP is called a *Bruck loop*. Bruck loops are necessarily Kikkawa loops (see, e.g., [9, 6.6(3)]). A Bruck loop is called a *B-loop* if the mapping $x \mapsto x \cdot x$ is a permutation.

Robinson's original definition of "Bruck loop" ([20], [21]) included the property that squaring is a permutation, and is thus equivalent to what we call a B-loop. Contemporary usage of the term "Bruck loop" in the literature (with some exceptions) tends to be as we have given it here, cf. [14], [15]. This usage seems to stem from a remark of Glauberman ([6, p. 376]).

Bruck loops are equivalent to the class of loops Ungar dubbed "gyrogroups" ([29]) and later "gyrocommutative gyrogroups" ([31]). Bruck loops are also equivalent to "K-loops", which are the additive loops of near-domains. Near-domains were introduced by Karzel [8], and the additive loop structure was later axiomatized and named in unpublished work of Kerby and Wefelscheid (the first appearance of the term "K-loop" in the literature was in a paper of Ungar [27]). The aforementioned equivalences have been established independently by various authors. Kreuzer showed the equivalence of Bruck loops with K-loops [15], and Sabinin *et al* showed the equivalence of Bruck loops with (gyrocommutative)

gyrogroups [25]. (The direct equivalence of gyrocommutative gyrogroups with K-loops is a well-known folk result.)

The term “B-loop” was introduced by Glauberman [6] to describe a finite, odd order Bol loop with the automorphic inverse property. Having odd order implies that the squaring map is a permutation, and Glauberman noted that for some results, it is this latter property which is essential, not the finiteness ([6, p. 374]). Since the contemporary usage of “Bruck loop” no longer implies that squaring is a bijection, it is quite natural to extend Glauberman’s terminology to the not necessarily finite case. (A less acronymic alternative would be “uniquely 2-divisible Bruck loop”.)

Proposition 1.5. 1. *Let $(M, *)$ be a reflection quasigroup with distinguished element $e \in M$, and define $x \cdot y = x^{1/2} * (e * y)$ for $x, y \in M$. Then (M, \cdot) is a B-loop.*

2. *Let (M, \cdot) be a B-loop and define $x * y = x^2 \cdot y^{-1}$ for $x, y \in M$. Then $(M, *)$ is a reflection quasigroup.*

In short, every reflection quasigroup is isotopic to a B-loop and conversely.

With different terminology and motivation than that employed here, Proposition 1.5 seems to have been first obtained by D. Robinson in his 1964 dissertation [20]; the result was not published, however, until 1979 [21]. Again with different jargon and with different axioms for what we are calling a B-loop, Kikkawa discovered the result independently in 1973 [10]. Kikkawa’s version emphasized the connection with symmetric spaces. The isotopic relationship between reflection quasigroups and B-loops is quite natural, and has been rediscovered other times in the literature, e.g., [3], [4], [5].

We have already noted that a smooth compact symmetric space $(M, *)$ is not a reflection quasigroup, and thus the symmetric space operation $*$ is not obtained from a B-loop operation by the previously described isotopy. However, for smooth symmetric spaces, whether compact or not, there is a well-developed *local* theory which guarantees that in a neighborhood of any distinguished point e , there exists a local B-loop structure with e as its identity element. This theory was worked out in detail primarily by Sabinin [23]; expositions can be found in [17] and [24]. For a smooth reflection quasigroup, the globally-defined B-loop operation agrees with the locally-defined operation guaranteed by the general theory wherever the latter operation is defined.

Every Bol loop, and hence every B-loop, is left alternative, and thus the relationship between a reflection quasigroup $(M, *)$ and the B-loop (M, \cdot) can be written

$$x * y = x \cdot (x \cdot y^{-1})$$

for $x, y \in M$. In the next section, we derive a *globally*-defined binary operation $\odot : \mathbb{S} \times \mathbb{S} \rightarrow \mathbb{S}$ which is compatible with the symmetric space structure $(\mathbb{S}, *)$ in the same sense:

$$(1.9) \quad \mathbf{x} * \mathbf{y} = \mathbf{x} \odot (\mathbf{x} \odot \mathbf{y}^{-1})$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{S}$. Our operation \odot will agree with the local B-loop operation guaranteed by general theory wherever the latter operation is defined. However, our approach will be elementary, in that we will not use any tools from differential geometry, nor even the intrinsic spherical distance on \mathbb{S} . Rather, we will simply use (1.2) and the Hilbert space structure.

2. Notation

We now introduce some notation which will be used throughout the paper. As in §1, let \mathbf{H} be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle : \mathbf{H} \times \mathbf{H} \rightarrow \mathbb{R}$ and corresponding norm $\|\mathbf{x}\| = \langle \mathbf{x}, \mathbf{x} \rangle^{1/2}$ for $\mathbf{x} \in \mathbf{H}$. For any subset $\mathbf{W} \subseteq \mathbf{H}$, we will denote by \mathbf{W}^\perp the orthogonal complement of \mathbf{W} in \mathbf{H} . Let \mathbf{e}_0 be a distinguished unit vector in \mathbf{H} , and let $\mathbf{V} = \mathbf{e}_0^\perp$ be the orthogonal complement in \mathbf{H} of \mathbf{e}_0 . Elements of the orthogonal direct sum $\mathbf{H} = \mathbb{R}\mathbf{e}_0 \oplus \mathbf{V}$ will be denoted by $\mathbf{x} = x_0\mathbf{e}_0 + \mathbf{x}_\perp$ where $x_0 = \langle \mathbf{x}, \mathbf{e}_0 \rangle$. In particular, for $\mathbf{V} = \mathbb{R}^n$, it is useful to identify $\mathbf{e}_0 \in \mathbb{R}^{n+1}$ with the standard basis element $(1, 0, \dots, 0)^T$.

For any closed subspace $\mathbf{E} \subseteq \mathbf{H}$, let $\mathcal{B}(\mathbf{E})$ denote the set of all bounded linear operators on \mathbf{E} . Given $A \in \mathcal{B}(\mathbf{E})$, the *transpose* of A is the operator $A^T \in \mathcal{B}(\mathbf{E})$ defined by

$$(2.1) \quad \langle A^T \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, A\mathbf{y} \rangle$$

for $\mathbf{x}, \mathbf{y} \in \mathbf{E}$. A bounded linear operator A is *symmetric* if $A^T = A$, and is *orthogonal* if $A^T A = I$, the identity transformation.

For a given $\mathbf{x} \in \mathbf{H}$, we define a linear functional $\mathbf{x}^T : \mathbf{H} \rightarrow \mathbb{R}$ by

$$(2.2) \quad \mathbf{x}^T \mathbf{y} = \langle \mathbf{x}, \mathbf{y} \rangle$$

for $\mathbf{y} \in \mathbf{H}$. For $\mathbf{a}, \mathbf{b} \in \mathbf{H}$, we define an operator $\mathbf{a}\mathbf{b}^T \in \mathcal{B}(\mathbf{H})$ by

$$(2.3) \quad \mathbf{a}\mathbf{b}^T \mathbf{x} = \langle \mathbf{b}, \mathbf{x} \rangle \mathbf{a}$$

for $\mathbf{x} \in \mathbf{H}$. This operator satisfies $(\mathbf{a}\mathbf{b}^T)^T = \mathbf{b}\mathbf{a}^T$. We denote the orthogonal projection onto the subspace $\mathbb{R} \cdot \mathbf{a}$ by

$$(2.4) \quad P_{\mathbf{a}} = \frac{\mathbf{a}\mathbf{a}^T}{\|\mathbf{a}\|^2}$$

The operator $P_{\mathbf{a}}$ is symmetric and $P_{\mathbf{a}}^2 = P_{\mathbf{a}}$.

For any closed subspace $\mathbf{E} \subseteq \mathbf{H}$, let

$$(2.5) \quad O(\mathbf{E}) = \{A \in \mathcal{B}(\mathbf{E}) : A^T A = I\}$$

be the orthogonal group of \mathbf{E} . We will identify $O(\mathbf{V})$ with a particular subgroup of $O(\mathbf{H})$, namely

$$(2.6) \quad O(\mathbf{V}) \cong \{A \in O(\mathbf{H}) : A\mathbf{e}_0 = \mathbf{e}_0\}.$$

For any $\mathbf{a} \in \mathbf{H}$, $A \in O(\mathbf{H})$, we have

$$(2.7) \quad AP_{\mathbf{a}} = P_{A\mathbf{a}}A^T.$$

Finally, let $J \in \mathcal{B}(\mathbf{H})$ be defined by

$$(2.8) \quad J\mathbf{x} = x_0\mathbf{e}_0 - \mathbf{x}_{\perp}$$

for $\mathbf{x} \in \mathbf{H}$. Then J is symmetric and orthogonal, and the following properties hold:

$$(2.9) \quad J^2 = I$$

$$(2.10) \quad J\mathbf{e}_0 = \mathbf{e}_0$$

$$(2.11) \quad JV = V$$

$$(2.12) \quad I + J = 2\mathbf{e}_0\mathbf{e}_0^T$$

3. Derivation of the operation

We now derive the operation \odot on \mathbb{S} , which will turn out to give (\mathbb{S}, \odot) a left loop structure. The distinguished point $\mathbf{e}_0 \in \mathbb{S}$ will be the identity element. We will use (1.8) as a guide. For $\mathbf{x} \in \mathbb{S}$, we wish to define $\mathbf{x}^{1/2}$ to be an element of \mathbb{S} satisfying $\mathbf{x}^{1/2} * \mathbf{e}_0 = \mathbf{x}$, provided such an element exists, and provided there is some canonical choice among such elements. For $\mathbf{u}, \mathbf{x} \in \mathbb{S}$, we have $\mathbf{u} * \mathbf{e}_0 = \mathbf{x}$ if and only if

$$(3.1) \quad 2\langle \mathbf{u}, \mathbf{e}_0 \rangle \mathbf{u} = \mathbf{e}_0 + \mathbf{x}.$$

If $\mathbf{x} = -\mathbf{e}_0$, then $\langle \mathbf{u}, \mathbf{e}_0 \rangle = 0$, and hence \mathbf{u} could be any element of $\mathbb{S} \cap \mathbf{V}$. Thus there is no unique choice of square root for the antipode $-\mathbf{e}_0 \in \mathbb{S}$. Therefore assume for now that $\mathbf{x} \neq -\mathbf{e}_0$. Taking norms of both sides of (3.1), we have $2|\mathbf{u}^T \mathbf{e}_0| = \|\mathbf{e}_0 + \mathbf{x}\|$. Therefore (3.1) implies

$$(3.2) \quad \mathbf{u} = \pm \frac{\mathbf{e}_0 + \mathbf{x}}{\|\mathbf{e}_0 + \mathbf{x}\|}.$$

Geometrically, it is clear from (3.2) that the preferred choice of square root is

$$(3.3) \quad \mathbf{x}^{1/2} = \frac{\mathbf{e}_0 + \mathbf{x}}{\|\mathbf{e}_0 + \mathbf{x}\|}$$

for $\mathbf{x} \neq -\mathbf{e}_0$. This can also be seen by a continuity argument: as $\mathbf{x} \rightarrow \mathbf{e}_0$, we have $\mathbf{u} \rightarrow \pm\mathbf{e}_0$. Taking $\mathbf{e}_0^{1/2} = \mathbf{e}_0$ and assuming $\mathbf{x} \mapsto \mathbf{x}^{1/2}$ to be continuous at \mathbf{e}_0 gives the plus sign convention in (3.2).

With (1.8) as motivation, for $\mathbf{x}, \mathbf{y} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{e}_0$, we set

$$(3.4) \quad \mathbf{x} \odot \mathbf{y} = \mathbf{x}^{1/2} * (\mathbf{e}_0 * \mathbf{y}).$$

Now for $\mathbf{y} \in \mathbb{S}$, we have

$$\mathbf{e}_0 * \mathbf{y} = 2\langle \mathbf{e}_0, \mathbf{y} \rangle \mathbf{e}_0 - \mathbf{y} = 2y_0 \mathbf{e}_0 - \mathbf{y} = y_0 \mathbf{e}_0 - \mathbf{y}_\perp = J\mathbf{y}.$$

Therefore for $\mathbf{x}, \mathbf{y} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{e}_0$, we have the following expression for the operation \odot :

$$(3.5) \quad \mathbf{x} \odot \mathbf{y} = (2P_{\mathbf{x}^{1/2}} - I) J\mathbf{y}.$$

What remains is to find a suitable definition of $-\mathbf{e}_0 \odot \mathbf{y}$. As noted, (3.4) is not appropriate here, because there is no canonical choice for “ $(-\mathbf{e}_0)^{1/2}$ ”. However, the correct definition of $-\mathbf{e}_0 \odot \mathbf{y}$ will be clear from the following remarks. Let $\Pi \subset \mathbf{H}$ be a plane passing through \mathbf{e}_0 and the origin, and fix $\mathbf{e}_1 \in \mathbb{S} \cap \Pi \cap \mathbf{V}$. For $\mathbf{x} \in \mathbb{S} \cap \Pi$, we have $\mathbf{x} = x_0 \mathbf{e}_0 + x_1 \mathbf{e}_1$, where $x_1 = \mathbf{x}_1^T \mathbf{e}_1$. In particular, for $\mathbf{x}, \mathbf{y} \in \Pi$, $\mathbf{e}_0 + \mathbf{x} = (1 + x_0)\mathbf{e}_0 + x_1 \mathbf{e}_1$, $\|\mathbf{e}_0 + \mathbf{x}\|^2 = (1 + x_0)^2 + x_1^2 = 2(1 + x_0)$, and $J\mathbf{y} = y_0 \mathbf{e}_0 - y_1 \mathbf{e}_1$. Thus for $\mathbf{x}, \mathbf{y} \in \mathbb{S} \cap \Pi$, it is easy to show with a few tedious calculations using (3.3) that (3.5) simplifies as follows:

$$(3.6) \quad \mathbf{x} \odot \mathbf{y} = (x_0 y_0 - x_1 y_1) \mathbf{e}_0 + (x_0 y_1 + x_1 y_0) \mathbf{e}_1.$$

Therefore the set $\mathbb{S} \cap \Pi$ is closed under the partial operation \odot . Now the limit of the expression in (3.6) as $\mathbf{x} \rightarrow -\mathbf{e}_0$ is $-y_0 \mathbf{e}_0 - y_1 \mathbf{e}_1 = -\mathbf{y}$; note that this is independent of our choice of \mathbf{e}_1 . Thus we define

$$(3.7) \quad -\mathbf{e}_0 \odot \mathbf{y} = -\mathbf{y}$$

for all $\mathbf{y} \in \mathbb{S}$.

The definition (3.7) has a geometric interpretation analogous to (3.4). While (3.4) defines $\mathbf{x} \odot \mathbf{y}$ as being the reflection of $\mathbf{e}_0 * \mathbf{y} = J\mathbf{y}$ about the point $\mathbf{x}^{1/2}$, (3.7) defines $-\mathbf{e}_0 \odot \mathbf{y}$ as being the reflection of $J\mathbf{y}$ across the subspace \mathbf{V} :

$$-\mathbf{e}_0 \odot \mathbf{y} = J\mathbf{y} - 2y_0 \mathbf{e}_0.$$

The following definition summarizes the preceding discussion.

Definition 3.1. Let \mathbb{S} denote the unit sphere in a real Hilbert space. Define a binary operation $\odot : \mathbb{S} \times \mathbb{S} \rightarrow \mathbb{S}$ by

$$(3.8) \quad \mathbf{x} \odot \mathbf{y} = \begin{cases} (2P_{\mathbf{x}^{1/2}} - I) J\mathbf{y}, & \text{if } \mathbf{x} \neq -\mathbf{e}_0 \\ -\mathbf{y}, & \text{if } \mathbf{x} = -\mathbf{e}_0 \end{cases}$$

for $\mathbf{x}, \mathbf{y} \in \mathbb{S}$.

Although most of the properties of the magma (\mathbb{S}, \odot) will follow from (3.8), there is another form for the operation \odot which is also occasionally useful. For $\mathbf{x}, \mathbf{y} \in \mathbb{S}$, $\mathbf{x} \neq -\mathbf{e}_0$, it easy to show by direct computation that

$$(3.9) \quad \mathbf{x} \odot \mathbf{y} = \langle \mathbf{x}, J\mathbf{y} \rangle \mathbf{e}_0 + \frac{y_0 + \langle \mathbf{x}, J\mathbf{y} \rangle}{1 + x_0} \mathbf{x}_\perp + \mathbf{y}_\perp.$$

Fix a plane $\Pi \subset \mathbf{H}$ through \mathbf{e}_0 and the origin, fix $\mathbf{e}_1 \in \mathbb{S} \cap \Pi \cap \mathbf{V}$ and consider the mapping $\mathbb{S} \cap \Pi \rightarrow \mathbb{C} : \mathbf{x} \mapsto x_0 + ix_1$, where $x_1 = \mathbf{x}_\perp^T \mathbf{e}_1$. This mapping is injective, and its image is the unit circle $S^1 \subset \mathbb{C}$. From (3.6) we see that this mapping is an isomorphism from the submagma $(\mathbb{S} \cap \Pi, \odot)$ to the circle group (S^1, \cdot) where the operation \cdot refers to multiplication of complex numbers. (This isomorphism depends on the choice of \mathbf{e}_1 , which fixes an orientation of the circle $\mathbb{S} \cap \Pi$.) We thus have the following result.

Theorem 3.2. *Let $\Pi \subset \mathbf{H}$ be a plane through \mathbf{e}_0 and the origin. Then $(\mathbb{S} \cap \Pi, \odot)$ is an abelian subgroup of (\mathbb{S}, \odot) .*

If we consider the case where $\mathbb{S} = S^1$, we have an immediately corollary.

Corollary 3.3. $(S^1, \odot) = (S^1, \cdot)$.

For $\mathbf{x} \in \mathbb{S}$, define

$$(3.10) \quad \mathbf{x}^{-1} = J\mathbf{x} = x_0 \mathbf{e}_0 - \mathbf{x}_\perp \in \mathbb{S}.$$

In particular, $\mathbf{e}_0^{-1} = \mathbf{e}_0$ and $(-\mathbf{e}_0)^{-1} = -\mathbf{e}_0$. The following is another corollary of Theorem 3.2.

- Corollary 3.4.** 1. *The element $\mathbf{e}_0 \in \mathbb{S}$ is a two-sided identity for (\mathbb{S}, \odot) .*
 2. *For each $\mathbf{x} \in \mathbb{S}$, \mathbf{x}^{-1} is a two-sided inverse of \mathbf{x} .*
 3. *(\mathbb{S}, \odot) is power-associative.*

Theorem 3.2 and the definitions (3.3) and (3.10) suggest another useful operation, namely an action of \mathbb{R} on \mathbb{S} .

Definition 3.5. Let \mathbb{S} denote the unit sphere in a real Hilbert space. For $\mathbf{x} \in \mathbb{S}$, $\mathbf{x} \neq -\mathbf{e}_0$, and for $t \in \mathbb{R}$, define

$$(3.11) \quad \mathbf{x}^t = \cos\left(t \cos^{-1} x_0\right) \mathbf{e}_0 + \sin\left(t \cos^{-1} x_0\right) \frac{\mathbf{x}_\perp}{\|\mathbf{x}_\perp\|}.$$

(Here $\cos^{-1} u = \arccos u$.) Also define $\mathbf{e}_0^t = \mathbf{e}_0$ for all $t \in \mathbb{R}$. Let $\omega : \mathbb{S} \setminus \{-\mathbf{e}_0\} \times \mathbb{R} \rightarrow \mathbb{S}$ denote the operation $(\mathbf{x}, t) \mapsto \mathbf{x}^t$.

Some properties easily follow from this definition. Once we have shown that the magma (\mathbb{S}, \odot) is a left loop, it will follow from the next result that $(\mathbb{S}, \odot, \omega)$ is a partial real odule (cf. [17, XII.1.16]).

Theorem 3.6. Fix $\mathbf{x} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{e}_0$. The following hold.

1. $\mathbf{x}^1 = \mathbf{x}$.
2. $(\mathbf{x}^s)^t = \mathbf{x}^{st}$ for $0 \leq s < \pi / \cos^{-1} x_0$ and for all $t \in \mathbb{R}$.
3. $\mathbf{x}^t \odot \mathbf{x}^s = \mathbf{x}^{t+s}$ for all $s, t \in \mathbb{R}$ such that $\mathbf{x}^t, \mathbf{x}^s, \mathbf{x}^{t+s} \neq -\mathbf{e}_0$.

PROOF: (1) is immediate from (3.11) since $\sin(\cos^{-1} x_0) = \|\mathbf{x}_\perp\|$. To simplify further calculations, set $a = \cos^{-1} x_0$. For (2), the restriction on s implies $\cos^{-1}(\cos(sa)) = sa$, and the result follows immediately. For (3), we compute

$$\langle \mathbf{x}^s, J\mathbf{x}^t \rangle = \cos(sa) \cos(ta) - \sin(ta) \sin(sa) = \cos((s + t)a)$$

and

$$\begin{aligned} \frac{\cos(sa) + \langle \mathbf{x}^s, J\mathbf{x}^t \rangle}{1 + \cos(ta)} \sin(ta) &= \cos(sa) \sin(ta) - \frac{\sin^2(ta) \sin(sa)}{1 + \cos(ta)} \\ &= \cos(sa) \sin(ta) - \sin(sa)(1 - \cos(ta)) \\ &= \sin((t + s)a) - \sin(sa). \end{aligned}$$

Using these calculations in (3.9) gives the desired result. □

The restriction on s in Theorem 3.6 (2) is generically unavoidable. For instance, if $\pi / \cos^{-1} x_0 < s < 2\pi / \cos^{-1} x_0$, then

$$(\mathbf{x}^s)^t = \cos(t(2\pi - s \cos^{-1} x_0))\mathbf{e}_0 + \sin\left(t(2\pi - s \cos^{-1} x_0)\right) \frac{\mathbf{x}_\perp}{\|\mathbf{x}_\perp\|} \neq \mathbf{x}^{st},$$

unless t is an integer.

Some consequences of Definition 3.5 and Theorem 3.6 are worth noting explicitly. First, $\mathbf{x}^0 = \mathbf{e}_0$, and in fact, for any integer $n \neq 0$, $\mathbf{x}^n = \mathbf{x} \odot \dots \odot \mathbf{x}$ (n factors). In particular, (3.11) agrees with (3.10) for $t = -1$. Also (3.11) agrees with (3.3) for $t = 1/2$.

We will consider additional properties of the magma (\mathbb{S}, \odot) in the next section. For now we make a couple of remarks which anticipate later results. We have seen that (S^1, \odot) is exactly the circle group. However, (S^3, \odot) is not (isomorphic to) the group of unit quaternions, and (S^7, \odot) is not (isomorphic to) the Moufang loop of unit octonions. This is because (\mathbb{S}, \odot) turns out to be a left loop, but not a loop for $\dim \mathbf{H} > 2$. Also (S^{15}, \odot) is not (isomorphic to) the left loop studied by Smith [26] using sedenian multiplication; the latter left loop is not power-associative.

Finally, recall that the 2-sphere S^2 may be identified with the *Riemann sphere* $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ by stereographic projection, where \mathbf{e}_0 , the “north pole”, is mapped to $0 \in \mathbb{C}$ and $-\mathbf{e}_0$, the “south pole”, is mapped to ∞ . Using this mapping, it is straightforward to transfer the operation \odot to $\hat{\mathbb{C}}$. For $x, y \in \hat{\mathbb{C}}$, we find that

$$(3.12) \quad x \odot y = \begin{cases} \frac{x + y}{1 - \bar{x}y} & \text{if } x, y \neq \infty \\ -1/\bar{y} & \text{if } x = \infty \\ -1/\bar{x} & \text{if } y = \infty \end{cases} .$$

Here \bar{x} denotes the complex conjugate of x , and we are using the usual conventions of complex arithmetic: $1/0 = \infty$ and $1/\infty = 0$. For $x, y \neq \infty$, (3.12) agrees with the local geodesic operation on $\hat{\mathbb{C}}$ found by Nesterov and Sabinin [18]. For the most part, results about $(\hat{\mathbb{C}}, \odot)$ are just special cases of results about (\mathbb{S}, \odot) .

4. Algebraic structure

It is clear from (3.8) that the left translation mappings $L_{\mathbf{x}} : \mathbb{S} \rightarrow \mathbb{S}$ defined by $L_{\mathbf{x}}\mathbf{y} = \mathbf{x} \odot \mathbf{y}$ are given by the restriction to \mathbb{S} of bounded linear operators defined on all of \mathbf{H} . Abusing notation slightly, we will denote these operators by $L_{\mathbf{x}}$ also. We have

$$(4.1) \quad L_{\mathbf{x}} = \begin{cases} (2P_{\mathbf{x}^{1/2}} - I)J, & \text{if } \mathbf{x} \neq -\mathbf{e}_0 \\ -I & \text{if } \mathbf{x} = -\mathbf{e}_0 \end{cases}.$$

Theorem 4.1. *Let $\mathbf{x} \in \mathbb{S}$ be given. The following hold.*

$$(4.2) \quad L_{\mathbf{x}} \in O(\mathbf{H}).$$

$$(4.3) \quad JL_{\mathbf{x}} = L_{J\mathbf{x}}J.$$

$$(4.4) \quad L_{\mathbf{x}}^T = L_{\mathbf{x}^{-1}}.$$

PROOF: All three assertions are trivial for $\mathbf{x} = -\mathbf{e}_0$, and thus we may assume $\mathbf{x} \neq -\mathbf{e}_0$. Equation (4.2) follows from the computation

$$L_{\mathbf{x}}L_{\mathbf{x}}^T = 4P_{\mathbf{x}^{1/2}}^2 - 4P_{\mathbf{x}^{1/2}} + I = I$$

using (2.9) and the fact that $P_{\mathbf{x}^{1/2}}$ is an orthogonal projection. Now

$$J\mathbf{x}^{1/2} = \mathbf{x}^{-1/2} = (J\mathbf{x})^{1/2},$$

and thus

$$JL_{\mathbf{x}} = (2JP_{\mathbf{x}^{1/2}} - J)J = (2P_{(J\mathbf{x})^{1/2}} - I)JJ = L_{J\mathbf{x}}J$$

using (2.7) (with $A = J$). This establishes (4.3). Finally, (4.4) follows from (4.3), (2.9), and the symmetry of $P_{\mathbf{x}^{1/2}}$:

$$L_{\mathbf{x}}^T = JL_{\mathbf{x}}J = L_{\mathbf{x}^{-1}}.$$

This completes the proof. □

Let $\text{Aut}(\mathbb{S}, \odot)$ denote the group of continuous automorphisms of (\mathbb{S}, \odot) .

Theorem 4.2. $\text{Aut}(\mathbb{S}, \odot) = O(\mathbf{V})$.

We omit the (long) proof, but note that the containment $O(\mathbf{V}) \subseteq \text{Aut}(\mathbb{S}, \odot)$ is clear from (3.9).

We will identify each left inner mapping $L(\mathbf{x}, \mathbf{y}) = L_{\mathbf{x} \odot \mathbf{y}}^{-1}L_{\mathbf{x}}L_{\mathbf{y}}$ on \mathbb{S} with its extension to a bounded linear operator on \mathbf{H} .

Theorem 4.3. For all $\mathbf{x}, \mathbf{y} \in \mathbb{S}$, $L(\mathbf{x}, \mathbf{y}) \in O(\mathbf{V})$.

PROOF: Since $O(\mathbf{H})$ is a group, $L(\mathbf{x}, \mathbf{y}) = L_{\mathbf{x} \odot \mathbf{y}}^{-1} L_{\mathbf{x}} L_{\mathbf{y}} \in O(\mathbf{H})$ by (4.2). We have $L(\mathbf{x}, \mathbf{y})\mathbf{e}_0 = L_{\mathbf{x} \odot \mathbf{y}}^{-1}(\mathbf{x} \odot \mathbf{y}) = \mathbf{e}_0$. In view of (2.6), this establishes the result. \square

Putting the preceding discussion together, we have identified the class of left loops to which (\mathbb{S}, \odot) belongs.

Corollary 4.4. (\mathbb{S}, \odot) is a Kikkawa left loop.

PROOF: The left inverse property follows from (4.2) and (4.4). The automorphic inverse property follows from (4.3). The A_l property follows from Theorems 4.3 and 4.2. \square

Not only do operators in $O(\mathbf{V})$ preserve the operation \odot , they also preserve the scalar multiplication ω given by (3.11). Indeed, for all $\mathbf{x} \in \mathbb{S}$, $\mathbf{x} \neq -\mathbf{e}_0$, for all $t \in \mathbb{R}$, and for all $A \in O(\mathbf{V})$, we have

$$(4.5) \quad (A\mathbf{x})^t = A\mathbf{x}^t$$

as an obvious consequence of the definitions. From (4.5) and Theorem 4.3, we have

$$(4.6) \quad (L(\mathbf{x}, \mathbf{y})\mathbf{z})^t = L(\mathbf{x}, \mathbf{y})\mathbf{z}^t$$

for all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{S}$, $\mathbf{z} \neq -\mathbf{e}_0$, $t \in \mathbb{R}$. Thus the \mathbb{R} -odule $(\mathbb{S}, \odot, \omega)$ satisfies Sabinin’s so-called “second A_l property” (cf. [17, XII.3.18]).

Kiechle has shown that there exist Kikkawa left loops which are not loops ([9, 11.3.2]). Here we will show that (\mathbb{S}, \odot) is not a loop. For $\mathbf{a} \in \mathbb{S}$, consider the equation

$$(4.7) \quad \mathbf{x} \odot \mathbf{a} = -\mathbf{a}^{-1}$$

in \mathbb{S} . From Theorem 3.2, $\mathbf{x} = -\mathbf{a}^{-2}$ is a solution. If $\mathbf{a} = \pm\mathbf{e}_0$, then clearly $\mathbf{x} = -\mathbf{a}^{-2} = -\mathbf{e}_0$ is the only solution. Conversely, if $\mathbf{x} = -\mathbf{e}_0$ is a solution, then $-\mathbf{a} = -\mathbf{a}^{-1}$, i.e., $\mathbf{a} = \pm\mathbf{e}_0$. In general, (4.7) has solutions other than $\mathbf{x} = -\mathbf{a}^{-2}$.

Theorem 4.5. For all $\mathbf{a} \in \mathbb{S}$, $\mathbf{a} \neq \pm\mathbf{e}_0$, the equation (4.7) has the solution set

$$(4.8) \quad \mathbf{X} = \left[-\mathbf{a}^{-2} + (\mathbf{a}^{-1})^\perp \right] \cap \mathbb{S} \setminus \{-\mathbf{e}_0\}.$$

If $\dim \mathbf{H} > 2$, then \mathbf{X} is not a singleton.

PROOF: By the preceding discussion, $\mathbf{x} \neq -\mathbf{e}_0$. Thus (4.7) becomes $(2P_{\mathbf{x}^{1/2}} - I)\mathbf{J}\mathbf{a} = -\mathbf{a}^{-1}$, i.e., $P_{\mathbf{x}^{1/2}}\mathbf{a}^{-1} = \mathbf{0}$. This is equivalent to $\langle \mathbf{x}^{1/2}, \mathbf{a}^{-1} \rangle = 0$, and thus to $\langle \mathbf{e}_0 + \mathbf{x}, \mathbf{a}^{-1} \rangle = 0$, or $\langle \mathbf{x}, \mathbf{a}^{-1} \rangle = -a_0$, with the constraint $\mathbf{x} \in \mathbb{S} \setminus \{-\mathbf{e}_0\}$. An obvious particular solution is $\mathbf{x} = -\mathbf{a}^{-2}$. Thus the solution set in \mathbf{H} is the affine

subspace $-\mathbf{a}^{-2} + (\mathbf{a}^{-1})^\perp$, and hence the solution set in \mathbb{S} is \mathbf{X} , as claimed. The nonuniqueness assertion follows from geometric considerations: \mathbf{X} is a punctured sphere of codimension 2 in \mathbf{H} with center $-a_0\mathbf{a}^{-1}$ and containing $-\mathbf{a}^{-2}$. The antipode of $-\mathbf{a}^{-2}$ is the removed point $-\mathbf{e}_0$. If \mathbf{X} is a singleton, then either $(\mathbf{a}^{-1})^\perp \cap \mathbb{S} = \{-\mathbf{a}^{-2}, -\mathbf{e}_0\}$, in which case $\dim \mathbf{H} \leq 2$, or $(\mathbf{a}^{-1})^\perp$ is tangent to \mathbb{S} at $-\mathbf{a}^{-2}$, in which case $-\mathbf{a}^{-2} = \pm\mathbf{a}^{-1}$ or $\mathbf{a} = \pm\mathbf{e}_0$, a contradiction. \square

We now turn to properties of (\mathbb{S}, \odot) which fail to hold everywhere, but which hold “almost everywhere”. First we consider the left power alternative property: $\mathbf{x}^s \odot (\mathbf{x}^t \odot \mathbf{y}) = \mathbf{x}^{s+t} \odot \mathbf{y}$. Special cases are the left inverse property ($s = \pm 1$, $t = \mp 1$) (which we already know to hold everywhere) and the left alternative property ($s = t = 1$).

Theorem 4.6. *For all $\mathbf{x} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{e}_0$ and for all $s, t \in \mathbb{R}$ such that $\mathbf{x}^s, \mathbf{x}^t \neq -\mathbf{e}_0$,*

$$(4.9) \quad L_{\mathbf{x}^s} L_{\mathbf{x}^t} = (2P_{\mathbf{x}^{(s+t)/2}} - I)J.$$

In addition, $\mathbf{x}^{s+t} \neq -\mathbf{e}_0$ if and only if

$$(4.10) \quad L_{\mathbf{x}^s} L_{\mathbf{x}^t} = L_{\mathbf{x}^{s+t}}.$$

PROOF: (Sketch) Fix $\mathbf{x} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{e}_0$. To simplify notation, set $C(t) = \cos((t/2) \cos^{-1} x_0)$ and $S(t) = \sin((t/2) \cos^{-1} x_0)$. For $s, t \in \mathbb{R}$ such that $\mathbf{x}^s, \mathbf{x}^t \neq -\mathbf{e}_0$, we have

$$(4.11) \quad L_{\mathbf{x}^s} L_{\mathbf{x}^t} J = 4P_{\mathbf{x}^{t/2}} P_{\mathbf{x}^{s/2}} - 2(JP_{\mathbf{x}^{s/2}} + P_{\mathbf{x}^{t/2}}J) + J$$

using (4.1). Now

$$\begin{aligned} P_{\mathbf{x}^{t/2}} P_{\mathbf{x}^{s/2}} &= \langle \mathbf{x}^{t/2}, J\mathbf{x}^{s/2} \rangle \mathbf{x}^{t/2} (\mathbf{x}^{s/2})^T \\ &= (C(t)C(s) - S(t)S(s)) \mathbf{x}^{t/2} (\mathbf{x}^{s/2})^T \\ &= C(t+s) \mathbf{x}^{t/2} (\mathbf{x}^{s/2})^T. \end{aligned}$$

Using (2.12) and (3.11), we may expand (4.11) into an expression of the form

$$L_{\mathbf{x}^s} L_{\mathbf{x}^t} J = \alpha \mathbf{e}_0 \mathbf{e}_0^T + \frac{\beta}{\|\mathbf{x}_\perp\|} \mathbf{e}_0 \mathbf{x}_\perp^T + \frac{\gamma}{\|\mathbf{x}_\perp\|} \mathbf{x}_\perp \mathbf{e}_0^T + \frac{\delta}{\|\mathbf{x}_\perp\|^2} \mathbf{x}_\perp \mathbf{x}_\perp^T - I,$$

where the coefficients $\alpha, \beta, \gamma, \delta$ are to be determined. We find, for instance,

$$\begin{aligned} \alpha &= 4C(t+s)C(t)C(s) - 2C(s)^2 - 2S(t)^2 + 2 \\ &= 2C(t+s)(C(t+s) + C(t-s)) - 2(C(s)^2 C(t)^2 - S(s)^2 S(t)^2) \\ &= 2C(t+s)^2. \end{aligned}$$

Similar calculations give

$$\begin{aligned} \beta &= \gamma = 2C(t+s)S(t+s) \\ \delta &= 2S(t+s)^2. \end{aligned}$$

Thus $L_{\mathbf{x}^s}L_{\mathbf{x}^t}J =$

$$2 \left(C(t+s)\mathbf{e}_0 + S(t+s)\frac{\mathbf{x}_\perp}{\|\mathbf{x}_\perp\|} \right) \left(C(t+s)\mathbf{e}_0 + S(t+s)\frac{\mathbf{x}_\perp}{\|\mathbf{x}_\perp\|} \right)^T - I.$$

This and (3.11) give the desired result. The remaining assertion is clear. □

Next we consider the cases ruled out in the hypotheses of Theorem 4.6.

Theorem 4.7. *If $\dim \mathbf{H} > 2$, then for all $\mathbf{x} \in \mathbb{S}$ with $\mathbf{x} \neq \pm\mathbf{e}_0$, $L_{-\mathbf{e}_0}L_{\mathbf{x}} = L_{\mathbf{x}}L_{-\mathbf{e}_0} \neq L_{-\mathbf{x}}$.*

PROOF: Fix $\mathbf{x} \in \mathbb{S}$ with $\mathbf{x} \neq \pm\mathbf{e}_0$, and assume $L_{\mathbf{x}}L_{-\mathbf{e}_0} = L_{-\mathbf{x}}$. Then $L_{\mathbf{x}^{-1}}L_{-\mathbf{x}} = -I$. Choose $\mathbf{y} \in \mathbb{S} \cap \mathbf{V} \cap \mathbf{x}^\perp$. Then $\mathbf{y} \in (J\mathbf{x})^\perp$. Suppressing details, we have $-\mathbf{y} = L_{\mathbf{x}^{-1}}L_{-\mathbf{x}}\mathbf{y} = -L_{\mathbf{x}^{-1}}J\mathbf{y} = \mathbf{y}$, a contradiction. □

The remaining case is clear:

$$(4.12) \quad L_{-\mathbf{e}_0}L_{-\mathbf{e}_0} = I = L_{\mathbf{e}_0} = L_{(-\mathbf{e}_0)^2}.$$

Putting together Theorems 4.6 and 4.7 with (4.12), we easily characterize those points in \mathbb{S} for which the left alternative property holds.

Corollary 4.8. *Assume $\dim \mathbf{H} > 2$. For all $\mathbf{x} \in \mathbb{S}$, $L_{\mathbf{x}}^2 = L_{\mathbf{x}^2}$ if and only if $\mathbf{x} \notin \mathbf{V}$.*

In particular, the left alternative property holds “almost everywhere”, since it fails only on the set $\mathbb{S} \cap \mathbf{V}$ of “measure zero”. In the finite dimensional case, we may remove the quotation marks.

A globally smooth Kikkawa left loop necessarily satisfies the left alternative property ([11, Lemma 6.2]. The result is stated for loops, but the proof clearly works for left loops.) It follows from Corollary 4.8 that the multiplication \odot on \mathbb{S} is not globally smooth. We will examine the continuity of \odot more closely in the next section.

Next we consider the Bol identity $L_{\mathbf{x}}L_{\mathbf{y}}L_{\mathbf{x}} = L_{\mathbf{x} \odot (\mathbf{y} \odot \mathbf{x})}$. Since the left alternative law does not hold everywhere, the Bol identity cannot hold everywhere either. Observe that the identity is trivial for $\mathbf{x} = \pm\mathbf{e}_0$.

Theorem 4.9. *Assume $\dim \mathbf{H} > 2$. Let $\mathbf{x} \in \mathbb{S}$, $\mathbf{x} \neq \pm\mathbf{e}_0$, be given. If $\mathbf{y} = -\mathbf{e}_0$ or if $\mathbf{y} \neq -\mathbf{e}_0$, $-\mathbf{x}^{-2}$ and $\mathbf{y} \odot \mathbf{x} = -\mathbf{x}^{-1}$, then $L_{\mathbf{x}}L_{\mathbf{y}}L_{\mathbf{x}} \neq L_{\mathbf{x} \odot (\mathbf{y} \odot \mathbf{x})}$.*

PROOF: We have

$$L_{\mathbf{x}}L_{-\mathbf{e}_0}L_{\mathbf{x}} = -L_{\mathbf{x}}^2 = -(2P_{\mathbf{x}} - I)J$$

by (4.9), and

$$L_{\mathbf{x} \odot (-\mathbf{e}_0 \odot \mathbf{x})} = L_{\mathbf{x} \odot (-\mathbf{x})} = L_{-\mathbf{x}^2} = (2P_{(-\mathbf{x}^2)^{1/2}} - I)J.$$

Now choose $\mathbf{z} \in \mathbf{V} \cap \mathbf{x}^\perp$. We have $L_{\mathbf{x}}L_{-\mathbf{e}_0}L_{\mathbf{x}}\mathbf{z} = J\mathbf{z} = -\mathbf{z}$ and $L_{\mathbf{x} \odot (-\mathbf{e}_0 \odot \mathbf{x})}\mathbf{z} = -J\mathbf{z} = \mathbf{z}$. Thus $L_{\mathbf{x}}L_{-\mathbf{e}_0}L_{\mathbf{x}} = L_{\mathbf{x} \odot (-\mathbf{e}_0 \odot \mathbf{x})}$ if and only if $\mathbf{V} \cap \mathbf{x}^\perp = \{\mathbf{0}\}$, i.e., if and only if $\dim \mathbf{H} \leq 2$.

Now assume $\mathbf{y} \neq -\mathbf{e}_0$ and $\mathbf{y} \odot \mathbf{x} = -\mathbf{x}^{-1}$ (i.e., $\mathbf{y} \in \mathbf{X}$ is given by (4.8)). Then $\mathbf{x} \odot (\mathbf{y} \odot \mathbf{x}) = -\mathbf{e}_0$, and thus $L_{\mathbf{x}}^{-1}L_{\mathbf{x} \odot (\mathbf{y} \odot \mathbf{x})}L_{\mathbf{x}}^{-1} = -L_{\mathbf{x}}^{-2}$. If the Bol identity holds, then $L_{\mathbf{y}} = -L_{\mathbf{x}}^{-2}$. Applying both sides to \mathbf{e}_0 , we have $\mathbf{y} = -\mathbf{x}^{-2}$. \square

The conditions described in the previous theorem turn out to be the only obstructions to the Bol identity.

Theorem 4.10. *Let $\mathbf{x}, \mathbf{y} \in \mathbb{S}$, $\mathbf{x} \neq \pm \mathbf{e}_0$, $\mathbf{y} \neq -\mathbf{e}_0$ and $\mathbf{y} \odot \mathbf{x} \neq -\mathbf{x}^{-1}$. Then $L_{\mathbf{x}}L_{\mathbf{y}}L_{\mathbf{x}} = L_{\mathbf{x} \odot (\mathbf{y} \odot \mathbf{x})}$.*

We omit the proof, but outline the idea. Applying both sides of the Bol identity to \mathbf{e}_0 gives an obvious equality, so it is only necessary to check that both sides agree when applied to an arbitrary element of \mathbf{V} . The hypotheses guarantee that all the left translations in the identity have the form $(2P_{\mathbf{u}^{1/2}} - I)J$, so all that remains is a calculation. This is tedious, but straightforward.

5. Continuity and smoothness

We have already noted in the previous section that the operation \odot cannot be globally smooth on $\mathbb{S} \times \mathbb{S}$ because (\mathbb{S}, \odot) does not everywhere satisfy the left alternative property.

For $\mathbf{x} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{e}_0$, we have

$$(5.1) \quad \lim_{t \rightarrow \pi / \cos^{-1} x_0} \mathbf{x}^t = -\mathbf{e}_0.$$

On the other hand, $\lim_{t \rightarrow \pi / \cos^{-1} x_0} \mathbf{x}^{t/2} = \mathbf{x}_\perp / \|\mathbf{x}_\perp\|$, and thus

$$(5.2) \quad \lim_{t \rightarrow \pi / \cos^{-1} x_0} L_{\mathbf{x}^t} = \left(2 \frac{\mathbf{x}_\perp \mathbf{x}_\perp^T}{\|\mathbf{x}_\perp\|^2} - I \right) J.$$

Now for $\mathbf{y} \in \mathbb{S}$, $((2/\|\mathbf{x}_\perp\|^2)\mathbf{x}_\perp \mathbf{x}_\perp^T - I)J\mathbf{y} = -\mathbf{y}$ if and only if $-\|\mathbf{x}_\perp\|^2 \mathbf{y}_\perp = (\mathbf{x}_\perp^T \mathbf{y}_\perp)\mathbf{x}_\perp$ i.e., if and only if \mathbf{x} , \mathbf{y} , and \mathbf{e}_0 are coplanar with $\mathbf{0}$. Therefore, for $\mathbf{x}, \mathbf{y} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{e}_0$, and $\mathbf{x}, \mathbf{y}, \mathbf{e}_0$ not coplanar with $\mathbf{0}$,

$$(5.3) \quad \lim_{t \rightarrow \pi / \cos^{-1} x_0} \mathbf{x}^t \odot \mathbf{y} \neq -\mathbf{e}_0 \odot \mathbf{y}.$$

For $\mathbf{y} \in \mathbb{S}$, define the right translation operator $R_{\mathbf{y}} : \mathbb{S} \rightarrow \mathbb{S} : \mathbf{x} \mapsto \mathbf{x} \odot \mathbf{y}$. We have established the following.

Theorem 5.1. *For each $\mathbf{y} \in \mathbb{S}$, $R_{\mathbf{y}} : \mathbb{S} \rightarrow \mathbb{S}$ has a nonremovable discontinuity at $-\mathbf{e}_0$.*

This turns out to be the only topological, or even analytic, deficiency.

Proposition 5.2. *The following hold.*

1. *For each $\mathbf{x} \in \mathbb{S}$, $L_{\mathbf{x}} : \mathbb{S} \rightarrow \mathbb{S}$ is analytic.*
2. *For each $\mathbf{y} \in \mathbb{S}$, $R_{\mathbf{y}} : \mathbb{S} \setminus \{-\mathbf{e}_0\} \rightarrow \mathbb{S}$ is analytic.*
2. *$\odot : \mathbb{S} \setminus \{-\mathbf{e}_0\} \times \mathbb{S} \rightarrow \mathbb{S}$ is analytic.*

PROOF: (Sketch) Note that (1) is immediate since $L_{\mathbf{x}}$ is the restriction of a bounded linear operator to \mathbb{S} . For (2), it is clear from (3.9) that $R_{\mathbf{y}}$ ($\mathbf{y} \in \mathbb{S}$) has an analytic extension to $\mathbf{H} \setminus \{-\mathbf{e}_0\}$. Similarly, (3.9) shows that \odot has an analytic extension to $\mathbf{H} \setminus \{-\mathbf{e}_0\} \times \mathbf{H}$, and this implies (3). □

6. Semidirect product structure

In this section we examine the relationship between the left loop structure of (\mathbb{S}, \odot) and the structure of the orthogonal group $O(\mathbf{H})$ relative to the subgroup $O(\mathbf{V})$. This discussion is a particular case of the general theory of semidirect products of left loops with groups. This theory was worked out in general by Sabinin [22] (see also [17], [24]), and was later rediscovered in the particular case of A_l left loops with LIP by Kikkawa [11] and Ungar [28]. A survey with recent extensions can be found in [12].

Since (\mathbb{S}, \odot) is an A_l left loop, we may form its *standard semidirect product* with $\text{Aut}(\mathbb{S}, \odot)$. This is the group denoted by $\mathbb{S} \rtimes \text{Aut}(\mathbb{S}, \odot)$, consisting of the set $\mathbb{S} \times \text{Aut}(\mathbb{S}, \odot)$ with multiplication defined by

$$(6.1) \quad (\mathbf{x}, A)(\mathbf{y}, B) = (\mathbf{x} \odot A\mathbf{y}, L(\mathbf{x}, A\mathbf{y})AB).$$

Define

$$(6.2) \quad L(\mathbb{S}) = \{L_{\mathbf{x}} : \mathbf{x} \in \mathbb{S}\}$$

to be the set of all left translations. Then $L(\mathbb{S}) \subset O(\mathbf{H})$ by (4.2), and $L(\mathbb{S})$ is a left transversal of the subgroup $O(\mathbf{V})$. Indeed, for $A \in O(\mathbf{H})$, let $\mathbf{u} = A\mathbf{e}_0$. Then $U = L_{\mathbf{u}}^{-1}A \in O(\mathbf{V})$. If $L_{\mathbf{u}}U = L_{\mathbf{v}}V$ with $\mathbf{v} \in \mathbb{S}$, $V \in O(\mathbf{H})$, then applying both sides to \mathbf{e}_0 , we obtain $\mathbf{u} = \mathbf{v}$, and thus $U = V$. Thus $A = L_{\mathbf{u}}U$ is a unique factorization of A into a left translation in $L(\mathbb{S})$ and an operator in $O(\mathbf{V})$.

The transversal decomposition $O(\mathbf{H}) = L(\mathbb{S})O(\mathbf{V})$ defines a left loop structure on $L(\mathbb{S})$ itself by projection: $L_{\mathbf{x}} \odot L_{\mathbf{y}} = L_{\mathbf{u}}$ where $L_{\mathbf{u}}$ is the unique representative of $L_{\mathbf{x}}L_{\mathbf{y}}$ in $L(\mathbb{S})$. Thus $O(\mathbf{H})$ is an *internal semidirect product* of the left loop $(L(\mathbb{S}), \odot)$ by the subgroup $O(\mathbf{V})$.

Now for $\mathbf{x}, \mathbf{y} \in \mathbb{S}$, $A, B \in O(\mathbf{V})$, $L_{\mathbf{x}}AL_{\mathbf{y}}B = L_{\mathbf{x} \odot A\mathbf{y}}L(\mathbf{x}, A\mathbf{y})AB$. Thus the left loops (\mathbb{S}, \odot) and $(L(\mathbb{S}), \odot)$ are just isomorphic copies (take $A = I$), and since $O(\mathbf{V}) \cong \text{Aut}(\mathbb{S}, \odot)$ (Theorem 4.1), we also have the following result.

Theorem 6.1. $O(\mathbf{H}) \cong \mathbb{S} \times O(\mathbf{V})$.

Interestingly, the transversal $L(\mathbb{S})$ in $O(\mathbf{H})$ is *not* connected. The element $L_{-\mathbf{e}_0} = -I$ is an isolated point. An intuitive argument can be seen as follows (cf. (5.3)). For any $\mathbf{u} \in \mathbb{S}$ with $\mathbf{u} \neq \pm \mathbf{e}_0$,

$$\lim_{\mathbf{x} \rightarrow \mathbf{u}} L_{\mathbf{x}} = \lim_{\mathbf{x} \rightarrow \mathbf{u}} (2P_{\mathbf{x}} - I)J = (2P_{\mathbf{u}} - I)J.$$

If $(2P_{\mathbf{u}} - I)J = -I$, then $2P_{\mathbf{u}} = I - J$. Apply this to any $\mathbf{v} \in \mathbb{S} \cap \mathbf{V} \cap \mathbf{u}^\perp$. Then $\mathbf{0} = (I - J)\mathbf{v} = 2\mathbf{v}$, a contradiction. Thus $\lim_{\mathbf{x} \rightarrow \mathbf{u}} L_{\mathbf{x}} \neq L_{-\mathbf{e}_0}$. This argument can easily be made more precise; we omit the details here. The nonconnectedness of $L(\mathbb{S})$ shows once again that the mapping $L : \mathbb{S} \rightarrow O(\mathbf{H}) : \mathbf{x} \mapsto L_{\mathbf{x}}$ is not everywhere continuous.

7. Spherical geometry

We now make a few remarks about the relationship between the Kikkawa left loop structure of (\mathbb{S}, \odot) and spherical geometry on \mathbb{S} . This relationship is analogous to that between the B -loop structure of the unit ball in a Hilbert space and hyperbolic geometry; see [31] and references therein.

On \mathbb{S} we define a norm by

$$(7.1) \quad \|\mathbf{x}\|_s = \cos^{-1} x_0 = \cos^{-1} \langle \mathbf{x}, \mathbf{e}_0 \rangle$$

for $\mathbf{x} \in \mathbb{S}$. Note the formal similarities between the following properties of $\|\cdot\|_s : \mathbb{S} \rightarrow \mathbb{R}$, and properties of norms in vector spaces.

Theorem 7.1. *The following hold.*

1. For all $\mathbf{x} \in \mathbb{S}$, $0 \leq \|\mathbf{x}\|_s \leq \pi$. In addition, $\|\mathbf{x}\|_s = 0$ if and only if $\mathbf{x} = \mathbf{e}_0$, and $\|\mathbf{x}\|_s = \pi$ if and only if $\mathbf{x} = -\mathbf{e}_0$.
2. For all $\mathbf{x} \in \mathbb{S}$, $\mathbf{x} \neq -\mathbf{e}_0$, $t \in \mathbb{R}$, $\|\mathbf{x}^t\|_s = |t| \|\mathbf{x}\|_s \pmod{\pi}$.
3. (Triangle inequality) For all $\mathbf{x}, \mathbf{y} \in \mathbb{S}$,

$$(7.2) \quad \|\mathbf{x} \odot \mathbf{y}\|_s \leq \pi - \|\mathbf{x}\|_s + \|\mathbf{y}\|_s - \pi \leq \|\mathbf{x}\|_s + \|\mathbf{y}\|_s.$$

The first inequality is an equality if and only if \mathbf{x} , \mathbf{y} , and \mathbf{e}_0 are coplanar with $\mathbf{0}$.

4. (Invariance under $O(\mathbf{V})$) For all $\mathbf{x} \in \mathbb{S}$, $A \in O(\mathbf{V})$, $\|A\mathbf{x}\|_s = \|\mathbf{x}\|_s$.

PROOF: (1), (2), and (4) are easy consequences of the definitions. For (3), we use (3.9) and the Cauchy-Schwarz inequality to compute

$$\begin{aligned} \cos \|\mathbf{x} \odot \mathbf{y}\|_s &= \langle \mathbf{x} \odot \mathbf{y}, \mathbf{e}_0 \rangle = \langle \mathbf{x}, J\mathbf{y} \rangle = x_0 y_0 - \langle \mathbf{x}_\perp, \mathbf{y}_\perp \rangle \\ &\geq x_0 y_0 - \|\mathbf{x}_\perp\| \|\mathbf{y}_\perp\| \\ &= \cos(\cos^{-1} x_0) \cos(\cos^{-1} y_0) - \sin(\cos^{-1} x_0) \sin(\cos^{-1} y_0) \\ &= \cos(\|\mathbf{x}\|_s + \|\mathbf{y}\|_s). \end{aligned}$$

By (1), we have

$$\|\mathbf{x} \odot \mathbf{y}\|_s \leq \cos^{-1}(\cos(\|\mathbf{x}\|_s + \|\mathbf{y}\|_s)) = \pi - \|\mathbf{x}\|_s + \|\mathbf{y}\|_s - \pi,$$

since $0 \leq \|\mathbf{x}\|_s + \|\mathbf{y}\|_s \leq 2\pi$. If equality holds, then the Cauchy-Schwarz equality implies that \mathbf{x}_\perp and \mathbf{y}_\perp are parallel, which is equivalent to the desired result. The remaining inequality of (7.2) is clear. \square

Next we use the operation \odot and the norm to define a distance function

$$(7.3) \quad d_s(\mathbf{x}, \mathbf{y}) = \left\| \mathbf{x} \odot \mathbf{y}^{-1} \right\|_s$$

for $\mathbf{x}, \mathbf{y} \in \mathbb{S}$. This definition is analogous to the usual relationship between norms and distance functions in vector spaces. By (3.9), we have $\langle \mathbf{x} \odot \mathbf{y}^{-1}, \mathbf{e}_0 \rangle = \langle \mathbf{x}, J\mathbf{y}^{-1} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle$, and hence $d_s(\mathbf{x}, \mathbf{y}) = \cos^{-1}\langle \mathbf{x}, \mathbf{y} \rangle$. This shows that (7.3) is equivalent to the usual definition of the spherical distance function.

Before establishing properties of the distance function, we require a lemma. The following result was established by Ungar for Bruck loops [30]. Here we extend it to Kikkawa left loops.

Lemma 7.2. *For all x, y, z in a Kikkawa left loop,*

$$(7.4) \quad L_x y \cdot (L_x z)^{-1} = L(x, y)(y \cdot z^{-1}).$$

PROOF: In any LIP left loop, we have the identity

$$(7.5) \quad L(x, y)^{-1} = L(x^{-1}, x \cdot y)$$

for all x, y (see, e.g., [9, 3.2]). In any Kikkawa left loop, we have the identities

$$(7.6) \quad L(x, y)^{-1} = L(y, x)$$

$$(7.7) \quad L(x, y)(y \cdot x) = x \cdot y$$

for all x, y (see, e.g., [9, 3.5 (4) and 3.5 (2)]). We compute

$$\begin{aligned} (x \cdot y) \cdot (x \cdot z)^{-1} &= (x \cdot y) \cdot (x^{-1} \cdot z^{-1}) \\ &= ((x \cdot y) \cdot x^{-1}) \cdot L(x \cdot y, x^{-1})z^{-1}. \end{aligned}$$

Now

$$\begin{aligned} (x \cdot y) \cdot x^{-1} &= L(x \cdot y, x^{-1})(x^{-1} \cdot (x \cdot y)) \\ &= L(x \cdot y, x^{-1})y, \end{aligned}$$

using (7.5) and LIP. Thus

$$(x \cdot y) \cdot (x \cdot z)^{-1} = L(x \cdot y, x^{-1})(y \cdot z^{-1}),$$

using the A_l property. Finally by (7.6), (7.5), and (7.6) again,

$$L(x \cdot y, x^{-1}) = L(x^{-1}, x \cdot y)^{-1} = L(x, y)^{-1} = L(y, x).$$

This establishes the result. \square

Theorem 7.3. *The following hold.*

1. For all $\mathbf{x}, \mathbf{y} \in \mathbb{S}$, $0 \leq d_s(\mathbf{x}, \mathbf{y}) \leq \pi$. In addition, $d_s(\mathbf{x}, \mathbf{y}) = 0$ if and only if $\mathbf{x} = \mathbf{y}$, and $d_s(\mathbf{x}, \mathbf{y}) = \pi$ if and only if $\mathbf{x} = -\mathbf{y}$.
2. (Triangle inequality) For all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{S}$, $d_s(\mathbf{x}, \mathbf{y}) \leq d_s(\mathbf{x}, \mathbf{z}) + d_s(\mathbf{z}, \mathbf{y})$.
3. (Invariance under $O(\mathbf{V})$) For all $\mathbf{x}, \mathbf{y} \in \mathbb{S}$, $d_s(A\mathbf{x}, A\mathbf{y}) = d_s(\mathbf{x}, \mathbf{y})$.
4. (Invariance under $L(\mathbb{S})$) For all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{S}$, $d_s(L_{\mathbf{x}}\mathbf{y}, L_{\mathbf{x}}\mathbf{z}) = d_s(\mathbf{y}, \mathbf{z})$.

PROOF: (1) follows from Theorem 7.1 (1). (3) follows from Theorems 4.2 and 7.1 (4). For (2), we have

$$\mathbf{x} \odot \mathbf{y}^{-1} = \mathbf{x} \odot (\mathbf{z}^{-1} \odot (\mathbf{z} \odot \mathbf{y}^{-1})) = (\mathbf{x} \odot \mathbf{z}^{-1}) \odot L(\mathbf{x}, \mathbf{z}^{-1})(\mathbf{z} \odot \mathbf{y}^{-1}).$$

Taking norms of both sides, and then using Theorem 7.1 (3), (4), we obtain the desired result. For (4), we use (7.4) in (\mathbb{S}, \odot) , take norms, and apply Theorem 7.1 (4):

$$\begin{aligned} d_s(\mathbf{x} \odot \mathbf{y}, \mathbf{x} \odot \mathbf{z}) &= \left\| (\mathbf{x} \odot \mathbf{y}) \odot (\mathbf{x} \odot \mathbf{z})^{-1} \right\|_s \\ &= \left\| L(\mathbf{x}, \mathbf{y})(\mathbf{y} \odot \mathbf{z}^{-1}) \right\|_s \\ &= \left\| \mathbf{y} \odot \mathbf{z}^{-1} \right\|_s \\ &= d_s(\mathbf{y}, \mathbf{z}). \end{aligned}$$

This completes the proof. □

Theorem 7.1 (3), (4) and the factorization $O(\mathbf{H}) = L(\mathbb{S})O(\mathbf{V})$ show that the distance function $d_s : \mathbb{S} \times \mathbb{S} \rightarrow [0, \infty)$ is invariant under the action of the entire orthogonal group $O(\mathbf{H})$ on \mathbb{S} .

To conclude, we note that the \mathbb{R} -odule structure $(\mathbb{S}, \odot, \omega)$ can be used to parametrize some of the interesting curves of spherical geometry; cf. the remarks in [31] about curves in hyperbolic geometry. For instance, for $\mathbf{x}, \mathbf{y} \in \mathbb{S}$ with $\mathbf{x} \neq -\mathbf{y}$, the unique spherical line (great circle) through \mathbf{x} and \mathbf{y} turns out to be given by

$$(7.8) \quad \gamma(t) = \mathbf{x} \odot (\mathbf{x}^{-1} \odot \mathbf{y})^t,$$

$t \in \mathbb{R}$. (Note that $\gamma(0) = \mathbf{x}$ and $\gamma(1) = \mathbf{y}$ by LIP.) Another interesting curve through $\mathbf{x}, \mathbf{y} \in \mathbb{S}$ where $\mathbf{x} \neq -\mathbf{y}^{-1}$ and $(\mathbf{y} \odot \mathbf{x}) \odot \mathbf{y}^{-1} \neq -\mathbf{y}$ is given by

$$(7.9) \quad \eta(t) = (\mathbf{y} \odot ((\mathbf{y} \odot \mathbf{x})^{-1} \odot \mathbf{y}))^t \odot \mathbf{x},$$

$t \in \mathbb{R}$. (Note that $\eta(0) = \mathbf{x}$ and $\eta(1) = \mathbf{y}$ by Theorem 4.10. The conditions on \mathbf{x}, \mathbf{y} guarantee that Theorem 4.10 can be applied.) The curve η is also a circle on \mathbb{S} which is everywhere equidistant from the spherical line $\nu(t) = (\mathbf{y} \odot ((\mathbf{y} \odot \mathbf{x})^{-1} \odot \mathbf{y}))^t$. Indeed, $d_s(\eta(t), \nu(t)) = d_s(\mathbf{x}, \mathbf{e}_0) = \|\mathbf{x}\|_s$, by Theorem 7.3 (4).

8. Complex spheres

To conclude this paper, we mention that many of the preceding ideas can be extended to *complex* spheres. We illustrate this with the complex 1-sphere

$$(8.1) \quad S_{\mathbb{C}}^1 = \left\{ \mathbf{x} = (x_0, x_1) \in \mathbb{C}^2 : |x_0|^2 + |x_1|^2 = 1 \right\}.$$

For $\mathbf{x}, \mathbf{y} \in S_{\mathbb{C}}^1$, define

$$(8.2) \quad \mathbf{x} \odot \mathbf{y} = \begin{cases} \left(\frac{x_0}{\bar{x}_0}(\bar{x}_0 y_0 - \bar{x}_1 y_1), x_0 y_1 + x_1 y_0 \right) & \text{if } x_0 \neq 0 \\ (-\bar{x}_1 y_1, x_1 y_0) & \text{if } x_0 = 0 \end{cases}.$$

It turns out that $(S_{\mathbb{C}}^1, \odot)$ is an A_l , LIP left loop. However, $(S_{\mathbb{C}}^1, \odot)$ does *not* satisfy AIP. A fuller exploration of $(S_{\mathbb{C}}^1, \odot)$ will appear elsewhere.

REFERENCES

- [1] Bruck R., *A Survey of Binary Systems*, (3rd printing, corrected), Springer-Verlag, Berlin, 1971.
- [2] Chein O., Pflugfelder H.O., Smith J.D.H. (eds.), *Quasigroups and Loops: Theory and Applications*, Sigma Series in Pure Mathematics, Vol. 8, Heldermann Verlag, Berlin, 1990.
- [3] Gabrieli E., Karzel H., *Reflection geometries over loops*, Results Math. **32** (1997), 61–65.
- [4] Gabrieli E., Karzel H., *Point-reflection geometries, geometric K-loops and unitary geometries*, Results Math. **32** (1997), 66–72.
- [5] Gabrieli E., Karzel H., *The reflection structures of generalized co-Minkowski spaces leading to K-loops*, Results Math. **32** (1997), 73–79.
- [6] Glauberman G., *On loops of odd order*, J. Algebra **1** (1964), 374–396.
- [7] Helgason S., *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, New York, 1978.
- [8] Karzel H., *Zusammenhänge zwischen Fastbereichen, scharf zweifach transitiven Permutationsgruppen und 2-Strukturen mit Rechtecksaxiom*, Abh. Math. Sem. Univ. Hamburg **32** (1968), 191–206.
- [9] Kiechle H., *Theory of K-loops*, Habilitationsschrift, Universität Hamburg, Hamburg, Germany, 1998.
- [10] Kikkawa M., *On some quasigroups of algebraic models of symmetric spaces*, Mem. Fac. Lit. Sci. Shimane Univ. (Nat. Sci.) **6** (1973), 9–13.
- [11] Kikkawa M., *The geometry of homogeneous Lie loops*, Hiroshima Math. J. **5** (1975), 141–179.
- [12] Kinyon M.K., Jones O., *Loops and semidirect products*, to appear in Comm. Algebra.
- [13] Klingenberg W.P.A., *Riemannian Geometry*, 2nd ed., Studies in Mathematics I, de Gruyter, New York, 1995.
- [14] Kepka T., *A construction of Bruck loops*, Comment. Math. Univ. Carolinae **25** (1984), 591–595.
- [15] Kreuzer A., *Inner mappings of Bol loops*, Math. Proc. Cambridge Philos. Soc. **123** (1998), 53–57.
- [16] Loos O., *Symmetric Spaces*, vol. 1, Benjamin, New York, 1969.
- [17] Miheev P.O., Sabinin L.V., *Quasigroups and differential geometry*, Chapter XII in [2], pp. 357–430.

- [18] Nesterov A.I., Sabinin L.V., *Smooth loops, generalized coherent states and geometric phases*, Int. J. Theor. Phys. **36** (1997), 1981–1989.
- [19] Pflugfelder H.O., *Quasigroups and Loops: Introduction*, Sigma Series in Pure Mathematics, Vol. 7, Heldermann Verlag, Berlin, 1990.
- [20] Robinson D.A., *Bol loops*, Ph.D. Dissertation, University of Wisconsin, Madison, Wisconsin, 1964.
- [21] Robinson D.A., *A loop-theoretic study of right-sided quasigroups*, Ann. Soc. Sci. Bruxelles Sér. I **93** (1979), 7–16.
- [22] Sabinin L.V., *On the equivalence of categories of loops and homogeneous spaces*, Soviet Math. Dokl. **13** (1972), 970–974.
- [23] Sabinin L.V., *Methods of nonassociative algebra in differential geometry* (Russian), Supplement to the Russian transl. of S. Kobayashi, K. Nomizu, “Foundations of Differential Geometry”, vol. 1, Nauka, Moscow, 1981, pp. 293–339.
- [24] Sabinin L.V., *Smooth Quasigroups and Loops*, Kluwer Academic Press, Dordrecht, 1999.
- [25] Sabinin L.V., Sabinina L.L., Sbitneva L., *On the notion of gyrogroup*, Aequationes Math. **56** (1998), 11–17.
- [26] Smith J.D.H., *A left loop on the 15-sphere*, J. Algebra **176** (1995), 128–138.
- [27] Ungar A.A., *The relativistic noncommutative nonassociative group of velocities and the Thomas rotation*, Results Math. **16** (1989), 168–179.
- [28] Ungar A.A., *Weakly associative groups*, Results Math. **17** (1990), 149–168.
- [29] Ungar A.A., *Thomas precession and its associated grouplike structure*, Amer. J. Phys. **59** (1991), 824–834.
- [30] Ungar A.A., *The holomorphic automorphism group of the complex disk*, Aequationes Math. **47** (1994), 240–254.
- [31] Ungar A.A., *Thomas precession: its underlying gyrogroup axioms and their use in hyperbolic geometry and relativistic physics*, Found. Phys. **27** (1997), 881–951.

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, INDIANA UNIVERSITY SOUTH BEND,
SOUTH BEND, IN 46634, USA

E-mail: mkinyon@iusb.edu

<http://www.iusb.edu/~mkinyon>

(Received October 15, 1999)