

Yan-Kui Song

Closed subsets of absolutely star-Lindelöf spaces II

Commentationes Mathematicae Universitatis Carolinae, Vol. 44 (2003), No. 2, 329–334

Persistent URL: <http://dml.cz/dmlcz/119389>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Closed subsets of absolutely star-Lindelöf spaces II

YAN-KUI SONG

Abstract. In this paper, we prove the following two statements: (1) There exists a discretely absolutely star-Lindelöf Tychonoff space having a regular-closed subspace which is not CCC-Lindelöf. (2) Every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented in a Hausdorff (regular, Tychonoff) absolutely star-Lindelöf space as a closed G_δ subspace.

Keywords: star-Lindelöf space, absolutely star-Lindelöf space

Classification: 54D20, 54B10, 54D55

1. Introduction

By a space, we mean a topological space. A space X is *absolutely star-Lindelöf* (see [1]) (*discretely absolutely star-Lindelöf* (see [8])) if for every open cover \mathcal{U} of X and every dense subspace D of X , there exists a countable subset $F \subseteq D$ such that $\text{St}(F, \mathcal{U}) = X$ (respectively, F is discrete and closed in X and $\text{St}(F, \mathcal{U}) = X$), where $\text{St}(F, \mathcal{U}) = \bigcup \{U \in \mathcal{U} : U \cap F \neq \emptyset\}$.

A space X is *star-Lindelöf* (see [4], [8] – under different name) if for every open cover \mathcal{U} of X , there exists a countable subset F of X such that $\text{St}(F, \mathcal{U}) = X$. It is clear that very separable space is star-Lindelöf as well as every space of countable extent (in particular, every countably compact space or every Lindelöf space).

A space X is *centered-Lindelöf* (*linked-Lindelöf*, *CCC-Lindelöf*) (see [2], [3]) if every open cover has a σ -centered (σ -linked, CCC, respectively) subcover. A family of sets is *centered* (*linked*) if every finite subfamily (every two elements, respectively) has non-empty intersection and a family is σ -centered (σ -linked) if it can be represented as the union of countably many centered-subfamilies (linked-subfamilies, respectively). A family of nonempty sets is a *CCC-family* if there is no uncountable pairwise disjoint subfamily.

From the above definitions, it is not difficult to see that every discretely absolutely star-Lindelöf space is absolutely star-Lindelöf, every absolutely star-Lindelöf space is star-Lindelöf, every star-Lindelöf space is centered-Lindelöf, every centered-Lindelöf space is linked-Lindelöf and every linked-Lindelöf space is CCC-Lindelöf.

The author is supported by NSFC project 10271056.

Bonanzinga and Matveev [2] proved that every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented as a closed subspace in Hausdorff (regular, Tychonoff, respectively) star-Lindelöf space. They asked if every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented as a closed G_δ subspace in Hausdorff (regular, Tychonoff, respectively) star-Lindelöf space. Song [8] gave a positive answer to their question. Moreover, Song and Shi [9] showed that every centered-Lindelöf Tychonoff space can be represented as a closed G_δ subspace in a Tychonoff absolutely star-Lindelöf space. But their construction does not work in the classes of Hausdorff and regular spaces. Thus, it is natural for us to consider the following more general question:

Question. Is it true that every Tychonoff (Hausdorff, regular) star-Lindelöf (centered-Lindelöf, linkered-Lindelöf) space can be embedded into some Tychonoff (Hausdorff, regular, respectively) absolutely star-Lindelöf space as a closed subspace? And can it be embedded as a closed G_δ subspace?

Throughout the paper, the cardinality of a set A is denoted by $|A|$. For a cardinal κ , κ^+ denotes the smallest cardinal greater than κ . Let \mathfrak{c} denote the cardinality of the continuum and ω the first infinite cardinal. As usual, a cardinal is the initial ordinal and an ordinal is the set of smaller ordinals. When viewed as a space, every cardinal has the usual order topology. Other terms and symbols will be used as in [5].

2. Closed subspaces of absolutely star-Lindelöf spaces

In [1], Bonanzinga showed that a regular-closed subspace of a star-Lindelöf space need not be star-Lindelöf, and in [9], Song and Shi showed that a regular-closed subspace of an absolutely star-Lindelöf space need not be absolutely star-Lindelöf. In the following, we give a stronger example to show that a regular-closed subspace of a discretely absolutely star-Lindelöf space need not be CCC-Lindelöf.

Recall that the Alexandroff duplicate of a space X , denoted by $A(X)$, is constructed in the following way: the underlying set of $A(X)$ is $X \times \{0, 1\}$ and each point of $X \times \{1\}$ is isolated; a basic neighborhood of a point $\langle x, 0 \rangle \in X \times \{0\}$ is a set of the form $(U \times \{0\}) \cup ((U \times \{1\}) \setminus \{\langle x, 1 \rangle\})$, where U is a neighborhood of x in X . It is well-known that $A(X)$ is Hausdorff (regular, Tychonoff, normal) iff so is X and $A(X)$ is compact iff so is X .

Recall from [6] that a space X is *absolutely countably compact* if for every open cover \mathcal{U} of X and every dense subspace D of X , there exists a finite subset $F \subseteq D$ such that $\text{St}(F, \mathcal{U}) = X$. Vaughan [10] proved that every countably compact GO-space is absolutely countably compact. Thus, every cardinal with uncountable cofinality is absolutely countably compact. In the next example we use the following lemma from [11].

Lemma 2.1. *If X is countably compact, then $A(X)$ is acc.*

Given a Tychonoff space X , let βX denote the Čech-Stone compactification of X .

Example 2.2. *There exists a discretely absolutely star-Lindelöf Tychonoff space X having a regular-closed subspace which is not CCC-Lindelöf.*

PROOF: Let \mathcal{R} be a maximal almost disjoint family of infinite subsets of ω with $|\mathcal{R}| = \mathfrak{c}$. Define $S_1 = (\mathfrak{c}^+ \times \omega) \cup \mathcal{R}$. We topologize X as follows: $\mathfrak{c}^+ \times \omega$ has the usual product topology and is an open subspace of S_1 , and a basic neighborhood of $r \in \mathcal{R}$ takes the form

$$G_{\beta,K}(r) = (\{\alpha : \beta < \alpha < \mathfrak{c}^+\} \times (r \setminus K)) \cup \{r\}$$

for $\beta < \mathfrak{c}^+$ and a finite subset K of ω . Then, the space S_1 is Tychonoff and $e(S_1) = \mathfrak{c}$, because \mathcal{R} is discrete and closed in X . Now, we show that S_1 is discretely absolutely star-Lindelöf. For this end, let \mathcal{U} be an open cover of S_1 . Let S be the set of all isolated points of \mathfrak{c}^+ and let $T = S \times \omega$. Then, T is dense in X and every dense subspace of X includes T . Thus, it suffices to show that there exists a countable subset $F \subseteq T$ such that F is discrete closed in X and $\text{St}(F, \mathcal{U}) = S_1$. For each $n < \omega$, since $\mathfrak{c}^+ \times \{n\}$ is absolutely countably compact, there exists a finite subset $F_n \subseteq S \times \{n\}$ such that $\mathfrak{c}^+ \times \{n\} \subseteq \text{St}(F_n, \mathcal{U})$. Let $F' = \bigcup \{F_n : n \in \omega\}$. Then, $\mathfrak{c}^+ \times \omega \subseteq \text{St}(F', \mathcal{U})$. For each $x \in \mathcal{R}$, take $U_x \in \mathcal{U}$ with $x \in U_x$, and fix $\alpha_x < \mathfrak{c}^+$ and $n_x \in \omega$ such that $\{\langle n_x, \alpha \rangle : \alpha_x < \alpha < \mathfrak{c}^+\} \subseteq U_x$. For each $n \in \omega$, let $X_n = \{x \in \mathcal{R} : n_x = n\}$ and choose $\beta_n \in S$ with $\beta_n > \sup\{\alpha_x : x \in X_n\}$. Then, $X_n \subseteq \text{St}(\langle \beta_n, n \rangle, \mathcal{U})$. It is quicker to choose $\beta \in S$ such that $\beta_n < \beta$ for all $n \in \omega$. Thus, if we put $F'' = \{\langle \beta, n \rangle : n \in \omega\}$, then $\mathcal{R} \subseteq \text{St}(F'', \mathcal{U})$. Let $F = F' \cup F''$. Then, F is a countable subset of D such that $S_1 = \text{St}(F, \mathcal{U})$. Since $F \cap (\mathfrak{c}^+ \times \{n\})$ is finite for each $n < \omega$, F is discrete and closed in S_1 , which shows that S_1 is discretely absolutely star-Lindelöf.

Let D be a discrete space of cardinality \mathfrak{c} and let

$$S_2 = A((\beta D \times (\mathfrak{c}^+ + 1)) \setminus ((\beta D \setminus D) \times \{\mathfrak{c}^+\})) \setminus ((D \times \{\mathfrak{c}^+\}) \times \{1\}).$$

We show that S_2 is not CCC-Lindelöf. Since $|D| = \mathfrak{c}$, we can enumerate D as $\{d_\alpha : \alpha < \mathfrak{c}\}$. For each $\alpha < \mathfrak{c}$, let

$$U_\alpha = (\{d_\alpha\} \times [0, \mathfrak{c}^+]) \times \{0, 1\} \setminus \{\langle \langle d_\alpha, \mathfrak{c}^+ \rangle, 0 \rangle\}.$$

Let us consider the open cover

$$\mathcal{U} = \{U_\alpha : \alpha < \mathfrak{c}\} \cup \{[0, \mathfrak{c}] \times \beta D\} \times \{0, 1\}$$

of S_2 . This contains an uncountable pairwise disjoint subfamily $\{U_\alpha : \alpha < \mathfrak{c}\}$ and since U_α is the only element of \mathcal{U} containing $\langle \langle \mathfrak{c}^+, d_\alpha \rangle, 0 \rangle$, S_2 is not CCC-Lindelöf.

We assume that $S_1 \cap S_2 = \emptyset$. Let $\varphi : \mathcal{R} \rightarrow \{\mathfrak{c}^+\} \times D \times \{0\}$ be a bijection. Let X be the quotient space obtained from the discrete sum $S_1 \oplus S_2$ by identifying r with $\varphi(r)$ for each $r \in \mathcal{R}$. Let $\pi : S_1 \oplus S_2 \rightarrow X$ be the quotient map. It is easy to check that $\pi(S_2)$ is a regular-closed subset of X , however, it is not CCC-Lindelöf, since it is homeomorphic to S_2 .

Now, we show that X is discretely absolutely star-Lindelöf. For this end, let \mathcal{U} be an open cover of X . Let

$$S' = \pi((D \times S \times \{0\}) \cup (\beta D \times [0, \mathfrak{c}^+] \times \{1\})) \cup \pi(T),$$

where S is the set of all isolated point of \mathfrak{c}^+ . Then, S' is dense in X and every dense subspace of X includes S' . Thus, it suffices to show that there exists a countable subset $F \subseteq S'$ such that F is discrete and closed in X and $\text{St}(F, \mathcal{U}) = X$. Since $\pi(S_1)$ is homeomorphic to S_1 , $\pi(S_1)$ is absolutely discretely star-Lindelöf, and there is a countable subset F_1 of S' such that F_1 is discrete and closed in $\pi(S_1)$ and $\pi(S_1) \subseteq \text{St}(F_1, \mathcal{U})$. Since $\pi(S_1)$ is a closed subset of X , F_1 is discrete and closed in X . On the other hand, since \mathfrak{c}^+ is locally compact and countably compact, it follows from [4, Theorem 3.10.13] that $\beta D \times \mathfrak{c}^+$ is countably compact. Thus $\pi(A(\beta D \times \mathfrak{c}^+))$ is acc by Lemma 2.1. Hence, there exists a finite subset F_2 of S' such that

$$\pi(A(\beta D \times \mathfrak{c}^+)) \subseteq \text{St}(F_2, \mathcal{U}).$$

If we put $F = F_1 \cup F_2$, then F is a countable subset of S' such that $X = \text{St}(F, \mathcal{U})$. Since F_1 is discrete and closed in X and F_2 is finite, F is discrete and closed in X , which completes the proof. □

Remark 1. Bonanzinga and Matveev [2] proved that a regular-closed subspace of a star-Lindelöf space need not be CCC-Lindelöf. In fact, they used Example 2.32 from [4] under Continuum Hypothesis. Example 2.1 is stronger than theirs.

Remark 2. Example 2.2 also shows that regular-closed subspaces of discretely absolutely star-Lindelöf (absolutely star-Lindelöf, star-Lindelöf, centered-Lindelöf, linked-Lindelöf and CCC-Lindelöf) spaces need not be discretely absolutely star-Lindelöf (absolutely star-Lindelöf, star-Lindelöf, centered-Lindelöf and linked-Lindelöf and CCC-Lindelöf, respectively).

Song and Shi [9] proved that every centered-Lindelöf Tychonoff space is representable as a closed G_δ subspace of a star-Lindelöf Tychonoff space. The following theorem is a generalization of this fact.

Theorem 2.3. *Every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented as a closed G_δ subspace in a Hausdorff (regular, Tychonoff, respectively) absolutely star-Lindelöf space.*

PROOF: First, we state a construction from [8, Theorem 2.1] which is a minor improvement of [2, Theorem 1]. Let X be a linked-Lindelöf space, $\mathcal{T}(X)$ the

topology of X and \mathcal{L} the collection of all linked subfamilies of $\mathcal{T}(X)$, and consider \mathcal{L} as a discrete space. Let $\mathcal{A} = \beta\mathcal{L} \times \omega$ and define $S(X) = X \cup \mathcal{A}$. We topologize $S(X)$ as follows: the subspace \mathcal{A} has the usual product topology and is an open subspace of $S(X)$; and a basic neighborhood of $x \in X$ in $S(X)$ is a set of the form

$$G_{U,n} = U \cup (\text{cl}_{\beta\mathcal{L}} L(U) \times \{m : n < m < \omega\})$$

for an open neighborhood U of x in X and $n < \omega$, where

$$L(U) = \{\mathcal{U} \in \mathcal{L} : (\exists V \in \mathcal{U})(V \subseteq U)\}.$$

Then, it was proved in [2, Theorem 1] and [8, Theorem 2.1] that $S(X)$ is Hausdorff (regular, Tychonoff) iff so is X and $S(X)$ is star-Lindelöf if X is linked-Lindelöf.

Now, we modify the above construction. Let

$$\mathcal{R}(S(X)) = A(S(X)) \setminus (X \times \{1\}).$$

Let $X_\omega = X \times \{0\}$ and $X_n = (\beta\mathcal{L} \times \{n\}) \times \{0, 1\}$ for each $n < \omega$. Then

$$\mathcal{R}(S(X)) = X_\omega \cup \bigcup_{n < \omega} X_n.$$

Then, X can be represented as $\mathcal{R}(S(X))$ as a closed- G_δ subspace, since X is homeomorphic to X_ω and $A(S(X))$ is Hausdorff (regular, Tychonoff) if X is Hausdorff (regular, Tychonoff). Thus it suffices to show that $A(S(X))$ is absolutely star-Lindelöf. Let \mathcal{U} be an open cover of $\mathcal{R}(S(X))$. Without loss of generality, we can assume that \mathcal{U} consists of basic open sets. Let

$$D_n = ((\mathcal{L} \times \{n\}) \times \{0\}) \cup ((\beta\mathcal{L} \times \{n\}) \times \{1\})$$

for each $n < \omega$ and let

$$D = \bigcup_{n < \omega} D_n.$$

Then, every dense subspace of $\mathcal{R}(S(X))$ includes D . Thus, it suffices to show that there exists a countable subset $F \subseteq D$ such that $\text{St}(F, \mathcal{U}) = \mathcal{R}(S(X))$. Let $\mathcal{U}_X = \{U \cap X_\omega : U \in \mathcal{U}\}$. Then, \mathcal{U}_X is an open cover of X_ω . Since X_ω is homeomorphic to X and X is linked-Lindelöf, \mathcal{U}_X has a σ -linked open refinement $\bigcup\{\mathcal{U}_m : m < \omega\}$. Let

$$F' = \{\langle \mathcal{U}_m, n \rangle, i : m < \omega, n < \omega, i = 0, 1\} \subseteq D.$$

To show that $X_\omega \subseteq \text{St}(F', \mathcal{U})$, let $x \in X$ be fixed. Then there exist $m, n < \omega$, $V \in \mathcal{U}_m$ and $U \in \mathcal{T}(X)$ such that

$$\langle x, 0 \rangle \in V \subseteq G'_{U,n} = (U \times \{0\}) \cup (\text{cl}_{\beta\mathcal{L}} L(U) \times \{m : n < m < \omega\}) \times \{0, 1\} \in \mathcal{U}.$$

Since $\langle \langle \mathcal{U}_m, n+1 \rangle, 1 \rangle \in F' \cap G'_{\mathcal{U}, n}$, $\langle x, 0 \rangle \in \text{St}(F', \mathcal{U})$. Hence, $X_\omega \subseteq \text{St}(F', \mathcal{U})$. On the other hand, since X_n is compact for each $n < \omega$, it is not difficult to find a finite subset $F_n \subseteq D_n$ such that $X_n \subseteq \text{St}(F_n, \mathcal{U})$. If we put $F = F' \cup \bigcup_{n < \omega} F_n$, then F is a countable subset of D and $\mathcal{R}(S(X)) = \text{St}(F, \mathcal{U})$, which completes the proof. \square

Since every star-Lindelöf space is centered-Lindelöf and every centered-Lindelöf space is linked-Lindelöf, the next corollary follows from Theorem 2.3.

Corollary 2.4 (Song and Shi [9]). *Every Hausdorff (regular, Tychonoff) centered-Lindelöf (star-Lindelöf) space can be embedded in a Hausdorff (regular, Tychonoff, respectively) absolutely star-Lindelöf space as a closed- G_δ subspace.*

Acknowledgments. This paper was written while the author was studying at Nanjing University as a post-doctor. He would like to use the opportunity to express his gratitude to Prof. W.-X. Shi for his valuable suggestions. He is also most grateful to the referee for his helpful comments.

REFERENCES

- [1] Bonanzinga M., *Star-Lindelöf and absolutely star-Lindelöf spaces*, Questions Answers Gen. Topology **16** (1998), 79–104.
- [2] Bonanzinga M., Matveev M.V., *Closed subspaces of star-Lindelöf and related spaces*, to appear in East-West J. Math.
- [3] Bonanzinga M., Matveev M.V., *Products of star-Lindelöf and related spaces*, Houston J. Math. **27** (2001), 45–57.
- [4] van Douwen E.K., Reed G.M., Roscoe A.W., Tree I.J., *Star covering properties*, Topology Appl. **39** (1991), 71–103.
- [5] Engelking R., *General Topology, Revised and completed edition*, Heldermann Verlag, Berlin, 1989.
- [6] Matveev M.V., *Absolutely countably compact spaces*, Topology Appl. **58** (1994), 81–92.
- [7] Matveev M.V., *A survey on star-covering properties*, Topology Atlas preprint No. 330 (1998).
- [8] Song Y.-K., *Remarks on star-Lindelöf spaces*, Questions Answers Gen. Topology **20** (2002), 49–51.
- [9] Song Y.-K., Shi W.-X., *Closed subspaces of absolutely star-Lindelöf spaces*, to appear in Houston J. Math.
- [10] Vaughan J.E., *On the product of a compact space with an absolutely countably compact space*, Annals of the New York Acad. Sci. **788** (1996), 203–208.
- [11] Vaughan J.E., *Absolute countable compactness and property (a)*, Talk at 1996 Praha symposium on General Topology.

DEPARTMENT OF MATHEMATICS, NANJING NORMAL UNIVERSITY, NANJING, 210097,
P.R. CHINA

and

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING, 210093, P.R. CHINA

E-mail: songyankui@email.njnu.edu.cn

(Received October 8, 2002, revised January 13, 2003)