

Genadij O. Hakobyan; V. N. Margaryan

Gevrey hypoellipticity for a class of degenerated quasi-elliptic operators

Commentationes Mathematicae Universitatis Carolinae, Vol. 44 (2003), No. 4, 637--644

Persistent URL: <http://dml.cz/dmlcz/119418>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2003

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Gevrey hypoellipticity for a class of degenerated quasi-elliptic operators

G.O. HAKOBYAN, V.N. MARGARYAN

Abstract. The problems of Gevrey hypoellipticity for a class of degenerated quasi-elliptic operators are studied by several authors (see [1]–[5]). In this paper we obtain the Gevrey hypoellipticity for a degenerated quasi-elliptic operator in \mathbb{R}^2 , without any restriction on the characteristic polyhedron.

Keywords: Gevrey class, Gevrey hypoellipticity, hypoelliptic operator, degenerated quasi-elliptic operator

Classification: 35B05, 35H10, 35H35

1. Statement of the result

Let \mathbb{R}^n , or E^n , be the n -dimensional real Euclidean space of points $\xi = (\xi_1, \dots, \xi_n)$, $x = (x_1, \dots, x_n)$ with real components. Let \mathbb{N}_0^n be the set of multi-indexes $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ with nonnegative integer components. Denote

$$\mathbb{R}_0^n = \{\xi \in \mathbb{R}^n; \xi_1 \dots \xi_n \neq 0\}, \quad \mathbb{R}_+^n = \{\xi \in \mathbb{R}^n; \xi_j \geq 0, j = 1, \dots, n\}.$$

For $\xi \in \mathbb{R}^n$, $\alpha \in \mathbb{N}_0^n$ we set $\xi^\alpha = \xi_1^{\alpha_1} \dots \xi_n^{\alpha_n}$, $D^\alpha = D_1^{\alpha_1} \dots D_n^{\alpha_n}$, where $D_j = \frac{\partial}{\partial \xi_j}$ or $D_j = -i \frac{\partial}{\partial x_j}$, $j = 1, \dots, n$.

Let Ω be an open subset of \mathbb{R}^n , $\lambda \in \mathbb{R}_+^n$, $\lambda_i \geq 1$, $i = 1, \dots, n$. We denote by $G^\lambda(\Omega)$ the class of all functions $f \in C^\infty(\Omega)$ so that for any compactum $K \subset \subset \Omega$ there exists a constant $C = C(K, f)$ for which

$$\sup_{x \in K} |D^\alpha f(x)| \leq C^{|\alpha|+1} \alpha_1^{\alpha_1 \lambda_1} \dots \alpha_n^{\alpha_n \lambda_n}, \quad \forall \alpha \in \mathbb{N}_0^n.$$

Let in \mathbb{R}^2 with variables x, y ,

$$(1) \quad P(x, D) = \sum_{\alpha=(\alpha_1, \alpha_2, \alpha_3) \in (P)} C_\alpha x^{\alpha_3} D_x^{\alpha_1} D_y^{\alpha_2},$$

be a differential operator with constant coefficients C_α . Here the sum is over a finite set of multi-indexes $(P) = \{\alpha : \alpha \in \mathbb{N}_0^3, C_\alpha \neq 0\}$.

Definition 1. The characteristic polyhedron (C.P.) $\mathcal{N}(P)$ of $P(x, D)$ is the smallest convex polyhedron in \mathbb{R}_+^3 containing all points $\alpha \in (P) \cup \{0\}$.

The results of Gevrey regularity for a certain class of quasi-elliptic operators degenerate on a symplectic manifold and with some restrictions on $\mathcal{N}(P)$ were obtained by V.V. Grushin, L. Rodino, L.R. Volevich, C. Parenti and others.

Let $\lambda_1 \geq 1$, $h \geq 0$ be rational numbers, $\lambda = (\lambda_1, 1)$. We denote

$$(2) \quad \mathcal{N} = \{\nu : \nu \in \mathbb{R}_+^3, \lambda_1\nu_1 + \nu_2 \leq m, \lambda_1\nu_1 + (1+h)\nu_2 - \lambda_1\nu_3 \leq m, \lambda_1\nu_3 \leq hm\}.$$

We consider differential operator (1) for which C.P. have form (2). It is easy to show that $m, m/\lambda_1, hm/\lambda_1, m/(1+h)$ are naturals. After introducing some preliminary lemmas we will prove the following result, cf. Theorem 1.

Theorem. *Let the hypoelliptic differential operator $P(x, D)$ from (1) with $\mathcal{N}(P)$ in form (2) satisfy in some neighborhood U of 0*

$$(3) \quad \sum_{(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{N}(P) \cap \mathbb{N}_0^3} \|x^{\alpha_3} D_x^{\alpha_1} D_y^{\alpha_2} \psi\|_{L_2(\Omega)} \leq \|P(x, D)\psi\|_{L_2(\Omega)}, \quad \psi \in C_0^\infty(U).$$

Then all solutions of equation $P(x, D) = f$ belong to the class $G^{(\lambda_1, 1)}(V)$, with $V \subset\subset U, (0, 0) \in V$, where $f \in G^{(\lambda_1, 1)}(U)$.

We observe that (3) implies that $|P(x, \xi, \eta)| \neq 0$ for $x, \xi, \eta \neq 0$, analogously to the condition asked by Volevich [5] in order to ensure the hypoellipticity of $P(D)$ for $x \neq 0$, under suitable conditions Parenti-Rodino [2] that the hypoellipticity continues to hold for $x = 0$.

2. Preliminary lemmas

Let $h \geq 0$, $\lambda_1 \geq 1$ and m, j be naturals.

We denote

$$\mathcal{M}_1^j = \{\nu : \nu \in \mathbb{R}_+^2, 2\lambda_1\nu_1 + \nu_2 \leq j, \lambda_1\nu_1 \leq (1+h)m\},$$

$$\mathcal{M}_2^j = \{\nu : \nu \in \mathbb{R}_+^2, \lambda_1\nu_1 + \nu_2 \leq j - (1+h)m, \lambda_1\nu_1 \geq (1+h)m\},$$

if $j < (1-h)m$ then we take $\mathcal{M}_2^j = \emptyset$. We set $\mathcal{M}^j = \mathcal{M}_1^j \cup \mathcal{M}_2^j$,

$$\mathcal{A}_1^j = \{\nu : \nu \in \mathbb{R}_+^3, \lambda_1\nu_1 + \nu_2 \leq m + j, 2\lambda_1\nu_1 + \nu_2 \leq j + 2m, \nu_3\lambda_1 \leq hm,$$

$$\lambda_1\nu_1 + (1+h)\nu_2 - \lambda_1\nu_3 \leq m + (1+h)j, \lambda_1\nu_1 \leq (1+h)m + m\},$$

$$\mathcal{A}_2^j = \{\nu : \nu \in \mathbb{R}_+^3, \lambda_1\nu_1 + \nu_2 \leq m + j - (1+h)m,$$

$$\lambda_1\nu_1 + (1+h)\nu_2 - \lambda_1\nu_3 \leq m + (1+h)(j - (1+h)m),$$

$$\nu_3\lambda_1 \leq hm, \lambda_1\nu_1 \geq (1+h)m\},$$

if $j < (1+h)m - m/(1+h)$ then we take $\mathcal{A}_2^j = \emptyset$.

Lemma 1. Let $h \geq 0$, $\lambda_1 \geq 1$, $m, m/\lambda_1, j$ be naturals and \mathcal{N} the polyhedron in form (2). Then any multi-index $\nu \in (\mathcal{A}_1^j \setminus \mathcal{A}_1^{j-1}) \cap \mathbb{N}_0^3$ can be represented in the form $\nu = \alpha + (\beta, 0)$ where $\alpha \in \mathcal{N} \cap \mathbb{N}_0^3$, $\beta \in \mathcal{M}_1^j \cap \mathbb{N}_0^2$.

PROOF: Let $\nu \in (\mathcal{A}_1^j \setminus \mathcal{A}_1^{j-1}) \cap \mathbb{N}_0^3$. If $\nu_1 \geq m/\lambda_1$ then we take $\alpha = (m/\lambda_1, 0, \nu_3) \in \mathbb{N}_0^3$, $\beta = (\nu_1 - m/\lambda_1, \nu_2) \in \mathbb{N}_0^2$. For α and β we have

$$\lambda_1\alpha_1 + \alpha_2 = m, \lambda_1\alpha_1 + (1+h)\alpha_2 - \lambda_1\alpha_3 = m - \lambda_1\nu_3 \leq m, \lambda_1\alpha_3 = \lambda_1\nu_3 \leq hm,$$

i.e. $\alpha \in \mathcal{N}$, $2\lambda_1\beta_1 + \beta_2 = 2\lambda_1(\nu_1 - m/\lambda_1) + \nu_2 = 2\lambda_1\nu_1 + \nu_2 - 2m \leq j + 2m - 2m = j$, $\lambda_1\beta_1 = \lambda_1\nu_1 - m \leq (1+h)m + m - m = (1+h)m$, i.e. $\beta \in \mathcal{M}_1^j$. If $\nu_1 < m/\lambda_1$ (i.e. $\lambda_1\nu_1 \leq m - \lambda_1$) then we consider the following possible cases:

- I) $2\lambda_1\nu_1 + \nu_2 > j - 1 + 2m$ hence $\nu_2 > j - 1 + 2m - 2m + 2\lambda_1 > j$,
- II) $\lambda_1\nu_1 + \nu_2 > m + j - 1$ hence $\nu_2 > m + j - 1 - m + \lambda_1 \geq j$,
- III) $\lambda_1\nu_1 + (1+h)\nu_2 - \lambda_1\nu_3 > m + (1+h)(j - 1)$ hence $(1+h)\nu_2 > m + (1+h)(j - 1) - m + \lambda_1$ i.e. $\nu_2 > j - 1$.

Therefore $\nu_2 \geq j$.

We take $\alpha = (\nu_1, \nu_2 - j, \nu_3) \in \mathbb{N}_0^3$, $\beta = (0, j) \in \mathbb{N}_0^2 \cap \mathcal{M}_1^j$. We obtain $\lambda_1\alpha_1 + \alpha_2 = \lambda_1\nu_1 + \nu_2 - j \leq m + j - j = m$, $\lambda_1\alpha_1 + (1+h)\alpha_2 - \lambda_1\alpha_3 = \lambda_1\nu_1 + (1+h)\nu_2 - \lambda_1\nu_3 - (1+h)j \leq m + (1+h)j - (1+h)j = m$, $\lambda_1\alpha_3 = \lambda_1\nu_3 \leq hm$, i.e. $\alpha \in \mathcal{N}$. \square

Lemma 2. Let $h \geq 0$, $\lambda_1 \geq 1$, $m, m/\lambda_1, j$ be naturals and \mathcal{N} the polyhedron in form (2). Then any multi-index $\nu \in (\mathcal{A}_2^j \setminus \mathcal{A}_2^{j-1}) \cap \mathbb{N}_0^3$ can be represented in the form $\nu = \alpha + (\beta, 0)$ where $\alpha \in \mathcal{N} \cap \mathbb{N}_0^3$, $\beta \in \mathcal{M}_2^j \cap \mathbb{N}_0^2$.

PROOF: Since $\nu \in (\mathcal{A}_2^j \setminus \mathcal{A}_2^{j-1}) \cap \mathbb{N}_0^3$, we have $j \geq hm$ and $\nu_1 \geq m/\lambda_1$. If $j < (1+h)m$ then $\lambda_1\nu_1 + \nu_2 \leq m + j - (1+h)m < m$ and $\lambda_1\nu_1 + (1+h)\nu_2 - \nu_3\lambda_1 \leq m + j - (1+h)m < m$ i.e. $\nu \in \mathcal{N}$. Therefore, we can take $\alpha = \nu \in \mathcal{N} \cap \mathbb{N}_0^3$ and $\beta = 0 \in \mathcal{M}_2^j \cap \mathbb{N}_0^2$. We can write $\nu = \alpha + \beta$.

If $j \geq (1+h)m$ then we take $\alpha = (m/\lambda_1, 0, \nu_3) \in \mathbb{N}_0^3$, $\beta = (\nu_1 - m/\lambda_1, \nu_2) \in \mathbb{N}_0^2$. Since $\lambda_1\alpha_1 + \alpha_2 = m$, $\lambda_1\alpha_1 + (1+h)\alpha_2 - \lambda_1\alpha_3 = m - \lambda_1\nu_3 \leq m$ and $\lambda_1\alpha_3 = \lambda_2\nu_3 \leq hm$, it follows that $\alpha \in \mathcal{N}$.

Let us show that $\beta \in \mathcal{M}_2^j$. We will consider the following possible cases:

- I) $\lambda_1\nu_1 - m \geq (1+h)m$ hence $\lambda_1\beta_1 + \beta_2 = \lambda_1\nu_1 + \nu_2 - m \leq m + j - (1+h)m - m = j - (1+h)m$, i.e. $\beta \in \mathcal{M}_2^j$,
- II) $\lambda_1\nu_1 - m \leq (1+h)m$ hence $2\lambda_1\beta_1 + \beta_2 = 2\lambda_1\nu_1 + \nu_2 - 2m = \lambda_1\nu_1 + \nu_2 + \lambda_1\nu_1 - 2m \leq m + j - (1+h)m + \lambda_1(\nu_1 - m/\lambda_1) - m \leq m + j - (1+h)m + (1+h)m - m = j$, i.e. $\beta \in \mathcal{M}_2^j$.

For $\nu_1 \geq m/\lambda_1$ we have $\alpha = (m/\lambda_1, 0, \nu_3) \in \mathcal{N} \cap \mathbb{N}_0^3$, $\beta = (\nu_1 - m/\lambda_1, \nu_2) \in (\mathcal{M}_1^j \cup \mathcal{M}_2^j) \cap \mathbb{N}_0^2$. \square

Lemma 3. Let $h \geq 0$, $\lambda_1 \geq 1$, $m, m/\lambda_1, j$ be naturals and \mathcal{N} the polyhedron (2). If $\alpha \in \mathcal{N}$, $\beta \in \mathcal{M}_1^j \cap \mathbb{N}_0^2$ then $(\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) \in \mathcal{A}_1^{j-\gamma_1}$ for $0 \leq \gamma_1 \leq \min(\beta_1, \alpha_3)$.

PROOF: Since $\beta \in \mathcal{M}_1^j \cap \mathbb{N}_0^2$ and $0 \leq \gamma_1 \leq \min(\beta_1, \alpha_3)$ we have $j \geq 2\lambda_1\beta_1 + \beta_2 \geq 2\lambda_1\gamma_1 \geq \gamma_1$. Therefore

$$\begin{aligned} 2\lambda_1(\alpha_1 + \beta_1 - \gamma_1) + \alpha_2 + \beta_2 &= 2(\lambda_1\alpha_1 + \alpha_2) + (2\lambda_1\beta_1 + \beta_2) - 2\lambda_1\gamma_1 \\ &\leq 2m + j - 2\lambda_1\gamma_1 \leq 2m + j - \gamma_1, \end{aligned}$$

$$\begin{aligned} \lambda_1(\alpha_1 + \beta_1 - \gamma_1) + (1+h)(\alpha_2 + \beta_2) - \lambda_1(\alpha_3 - \gamma_1) \\ &= (\lambda_1\alpha_1 + (1+h)\alpha_2 - \lambda_1\alpha_3) + (\lambda_1\beta_1 + (1+h)\beta_2) \\ &\leq m + (1+h)(2\lambda_1\beta_1 + \beta_2) - (2h+1)\lambda_1\beta_1 \\ &\leq m + (1+h)j - (2h+1)\lambda_1\gamma_1 \leq m + (1+h)(j - \gamma_1), \end{aligned}$$

$$\begin{aligned} \lambda_1(\alpha_1 + \beta_1 - \gamma_1) &= \lambda_1\alpha_1 + \lambda_1\beta_1 - \lambda_1\gamma_1 \leq m + (1+h)m - \lambda_1\gamma_1 \leq m + (1+h)m, \\ \text{i.e. } (\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) &\in \mathcal{A}_1^{j-\gamma_1}. \end{aligned} \quad \square$$

Lemma 4. Let $h \geq 0$, $\lambda_1 \geq 1$, $m, m/\lambda_1, j$ be naturals and \mathcal{N} the polyhedron (2). If $\alpha \in \mathcal{N} \cap \mathbb{N}_0^3$, $\beta \in \mathcal{M}_2^j \cap \mathbb{N}_0^2$ then $(\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) \in \mathcal{A}_2^{j-\gamma_1} \cup \mathcal{A}_1^{j-\gamma_1}$ for $0 \leq \gamma_1 \leq \min(\beta_1, \alpha_3)$.

PROOF: Since $\beta \in \mathcal{M}_1^j \cap \mathbb{N}_0^2$ and $0 \leq \gamma_1 \leq \min(\beta_1, \alpha_3)$ we have $j \geq \lambda_1\beta_1 + (1+h)m > \gamma_1$. We consider the following possible cases:

I) for $\lambda_1(\alpha_1 + \beta_1 - \gamma_1) \geq (1+h)m$ we obtain

$$\begin{aligned} \text{a) } \lambda_1(\alpha_1 + \beta_1 - \gamma_1) + (\alpha_2 + \beta_2) &= (\lambda_1\alpha_1 + \alpha_2) + (\lambda_1\beta_1 + \beta_2) - \lambda_1\gamma_1 \\ &\leq m + j - (1+h)m - \lambda_1\gamma_1 \\ &\leq m + (j - \gamma_1) - (1+h)m, \end{aligned}$$

b) since $\alpha \in \mathcal{N}$, $\beta \in \mathcal{M}_2^j$ and $\gamma_1 \leq \alpha_3$ we have

$$\begin{aligned} \lambda_1(\alpha_1 + \beta_1 - \gamma_1) + (1+h)(\alpha_2 + \beta_2) - \lambda_1(\alpha_3 - \gamma_1) \\ &= (\lambda_1\alpha_1 + (1+h)\alpha_2 - \lambda_1\alpha_3) + (\lambda_1\beta_1 + (1+h)\beta_2) \\ &\leq m + (1+h)(\lambda_1\beta_1 + \beta_2) - h\lambda_1\beta_1 \\ &\leq m + (1+h)(j - (1+h)m) - h\lambda_1\beta_1 \\ &\leq m + (1+h)(j - (1+h)m) - h(1+h)m \\ &\leq m + (1+h)(j - (1+h)m) - (1+h)\lambda_1\alpha_3 \\ &\leq m + (1+h)(j - (1+h)m) - (1+h)\lambda_1\gamma_1 \\ &\leq m + (1+h)(j - \gamma_1 - (1+h)m), \end{aligned}$$

c) $\lambda_1(\alpha_3 - \gamma_1) \leq \lambda_1\alpha_3 \leq hm$.

From a), b), c) for case I) we obtain $(\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) \in \mathcal{A}_2^{j-\gamma_1}$ or $(\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) \in \mathcal{A}_2^{j-\gamma_1} \cup \mathcal{A}_1^{j-\gamma_1}$;

II) for $\lambda_1(\alpha_1 + \beta_1 - \gamma_1) \leq (1+h)m$ we obtain

$$\begin{aligned} \text{a)} \quad 2\lambda_1(\alpha_1 + \beta_1 - \gamma_1) + (\alpha_2 + \beta_2) &= (2\lambda_1\alpha_1 + \alpha_2) + (2\lambda_1\beta_1 + \beta_2) - 2\lambda_1\gamma_1 \\ &\leq 2m + j - 2\lambda_1\gamma_1 \leq 2m + (j - \gamma_1), \end{aligned}$$

$$\begin{aligned} \text{b)} \quad \lambda_1(\alpha_1 + \beta_1 - \gamma_1) + (\alpha_2 + \beta_2) &= (\lambda_1\alpha_1 + \alpha_2) + (\lambda_1\beta_1 + \beta_2) - \lambda_1\gamma_1 \\ &\leq m + j - (1+h)m - \lambda_1\gamma_1 \leq m + j - \gamma_1, \end{aligned}$$

c) since $\alpha \in \mathcal{N}$, $\beta \in \mathcal{M}_2^j$ and $\gamma_1 \leq \alpha_3$,

$$\begin{aligned} \lambda_1(\alpha_1 + \beta_1 - \gamma_1) + (1+h)(\alpha_2 + \beta_2) - \lambda_1(\alpha_3 - \gamma_1) \\ &= (\lambda_1\alpha_1 + (1+h)\alpha_2 - \lambda_1\alpha_3) + (\lambda_1\beta_1 + (1+h)\beta_2) \\ &\leq m + (1+h)(\lambda_1\beta_1 + \beta_2) - h\lambda_1\beta_1 \\ &\leq m + (1+h)(j - (1+h)m) - h(1+h)m \\ &\leq m + (1+h)(j - (1+h)m) - (1+h)\alpha_3 \\ &\leq m + (1+h)(j - (1+h)m) - (1+h)\gamma_1 \\ &= m + (1+h)(j - \gamma_1 - (1+h)m). \end{aligned}$$

Therefore $(\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) \in \mathcal{A}_1^{j-\gamma_1}$ or $(\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) \in \mathcal{A}_2^{j-\gamma_1} \cup \mathcal{A}_1^{j-\gamma_1}$. \square

3. Main results

Let $P(x, D) = \sum C_\alpha x^{\alpha_3} D_x^{\alpha_1} D_y^{\alpha_2}$ be a differential operator with C.P. in form (2). Since $m, m/\lambda_1, hm/\lambda_1, m/(1+h)$ are naturals, Lemmas 1–4 are valid for $\mathcal{N}(P)$.

For $t > 0$ we denote $B_t = \{(x, y) \in \mathbb{R}^2, |x|^2 + |y|^2 < t^2\}$.

We use a well known result (see for example Lemma 2.1 in [3]).

Lemma 5. *Let $\rho_1 > 0$, $\rho > 0$. Then there exists a function $\varphi \in C_0^\infty(\mathbb{R}^2)$ such that $\text{supp } \varphi \subset B_{\rho_1+\rho}$, $\varphi(x, y) = 1$, $(x, y) \in B_{\rho_1}$, $0 \leq \varphi(x, y) \leq 1$ and*

$$\max_{x,y} |D_x^{\alpha_1} D_y^{\alpha_2} \varphi(x, y)| \leq C_{\alpha_1, \alpha_2} \rho^{-(\alpha_1 + \alpha_2)},$$

where C_{α_1, α_2} is independent of ρ_1 and ρ .

For $u \in C^\infty$ we denote

$$\begin{aligned}\|u, \sigma\| &= \sum_{\alpha \in \mathcal{N} \cap \mathbb{N}_0^3} \|x^{\alpha_3} D_x^{\alpha_1} D_y^{\alpha_2}\|_{L_2(B_\sigma)}, \\ \|u, \sigma\|_t &= \max_{\beta_1, \beta_2 \in \mathcal{M}^t \cap \mathbb{N}_0^2} \|D_x^{\beta_1} D_y^{\beta_2} u, \sigma\| \quad \text{for } t > 0\end{aligned}$$

and

$$\|u, \sigma\|_t = \|u, \sigma\| \quad \text{for } t < 0.$$

Lemma 6. Let $u \in C^\infty$, $\alpha \in \mathcal{N} \cap \mathbb{N}_0^3$, $0 \leq \alpha'_1 \leq \alpha_1$, $0 \leq \alpha'_2 \leq \alpha_2$, $\alpha'_1, \alpha'_2 \in \mathcal{N}$, $\rho \in (0, 1)$. Then there exists a constant $C > 0$ so that for any $(\beta_1, \beta_2) \in \mathcal{M}^j \cap \mathbb{N}_0^2$, $j = 1, 2, \dots$,

$$\|x^{\alpha_3} (D_x^{\alpha'_1} D_y^{\alpha'_2} \varphi) D_x^{\alpha_1 - \alpha'_1 + \beta_1} D_y^{\alpha_2 - \alpha'_2 + \beta_2} u\|_{L_2} \leq C \rho^{-(\alpha'_1 + \alpha'_2)} \|u, \rho_1 + \rho\|_{j - (\alpha'_1 + \alpha'_2)},$$

where φ is from Lemma 5.

PROOF: We can assume without loss of generality that $j \geq \alpha'_1 + \alpha'_2$. Now we show that $\alpha = (\alpha_1 - \alpha'_1 + \beta_1, \alpha_2 - \alpha'_2 + \beta_2, \alpha_3) \in \mathcal{A}_1^{j - (\alpha'_1 + \alpha'_2)} \cup \mathcal{A}_2^{j - (\alpha'_1 + \alpha'_2)}$. From Lemmas 1, 2, α can be taken in form $\alpha = (\mu_1, \mu_2, \mu_3) + (\nu_1, \nu_2, 0)$ where $(\mu_1, \mu_2, \mu_3) \in \mathcal{N} \cap \mathbb{N}_0^3$, $(\nu_1, \nu_2) \in \mathcal{M}^{j - (\alpha'_1 + \alpha'_2)} \cap \mathbb{N}_0^2$. Then from Lemma 5 we obtain

$$\begin{aligned}&\|x^{\alpha_3} (D_x^{\alpha'_1} D_y^{\alpha'_2} \varphi) D_x^{\alpha_1 - \alpha'_1 + \beta_1} D_y^{\alpha_2 - \alpha'_2 + \beta_2} u\|_{L_2} \\ &\leq C_{\alpha'_1, \alpha'_2} \rho^{-(\alpha'_1 + \alpha'_2)} \|x^{\mu_3} (D_x^{\mu_1} D_y^{\mu_2} (D_x^{\nu_1} D_y^{\nu_2} u))\|_{L_2(B_{\rho_1 + \rho})} \\ &\leq C \rho^{-(\alpha'_1 + \alpha'_2)} \|u, \rho_1 + \rho\|_{j - (\alpha'_1 + \alpha'_2)},\end{aligned}$$

where $C = \max_{\alpha'_1 \leq m, \alpha'_2 \leq m} C_{\alpha'_1, \alpha'_2}$. \square

Corollary 1. Let $U \in C^\infty$. Then there exists $C > 0$ such that for all $(\beta_1, \beta_2) \in \mathcal{M}^j \cap \mathbb{N}_0^2$, $j \geq 1$,

$$\|[P, \varphi] D_x^{\beta_1} D_y^{\beta_2} u\|_{L_2} \leq C \sum_{i=1}^m \rho^{-i} \|u, \rho_1 + \rho\|_{j-i},$$

where φ is from Lemma 5.

PROOF: The proof follows from Lemma 6, if we note that $[P, \varphi]$ is representable by linear combination of terms in form

$$x^{\alpha_3} (D_x^{\alpha'_1} D_y^{\alpha'_2} \varphi) D_x^{\alpha_1 - \alpha'_1} D_y^{\alpha_2 - \alpha'_2},$$

where $(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{N} \cap \mathbb{N}_0^3$, $0 \leq \alpha'_1 \leq \alpha_1$, $0 \leq \alpha'_2 \leq \alpha_2$, and $\alpha'_1 + \alpha'_2 \geq 1$. \square

Lemma 7. Let $U \in C^\infty$ and $(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{N} \cap \mathbb{N}_+^3$, $(\beta_1, \beta_2) \in \mathcal{M}^j \cap \mathbb{N}_0^2$, $0 \leq \gamma_1 \geq \beta_1$, $\gamma_1 \in \mathcal{N}$. Then

$$(4) \quad \|(D_x^{\gamma_1} x^{\alpha_3}) D_x^{\alpha_1 + \beta_1 - \gamma_1} D_y^{\alpha_2 + \beta_2} u\|_{L_2(B_{\rho_1 + \rho})} \leq C \|u, \rho_1 + \rho\|_{j-\gamma_1}$$

with some constant $C > 0$.

PROOF: Inequality (4) is trivial for $\gamma_1 > \alpha_3$. Let $\gamma_1 \leq \alpha_3$. Then from Lemmas 3, 4 we obtain that $(\alpha_1 + \beta_1 - \gamma_1, \alpha_2 + \beta_2, \alpha_3 - \gamma_1) \in \mathcal{A}_2^{j-\gamma_1} \cup \mathcal{A}_1^{j-\gamma_1}$. From Lemmas 1, 2, this multi-index can be taken in form $(\alpha'_1, \alpha'_2, \alpha_3 - \gamma_1) + (\beta'_1, \beta'_2, 0)$, where $(\alpha'_1, \alpha'_2, \alpha_3 - \gamma_1) \in \mathcal{N} \cap \mathbb{N}_+^3$, $(\beta'_1, \beta'_2) \in \mathcal{M}^{j-\gamma_1}$. If we take $C \geq \alpha_3!/\gamma_1!$ then the proof is complete. \square

Corollary 2. Let $U \in C^\infty$. Then there exists a constant $C > 0$ so that for any multi-index $(\beta_1, \beta_2) \in \mathcal{M}^j \cap \mathbb{N}_0^2$, $j = 1, 2, \dots$,

$$\|[P, D_x^{\beta_1} D_y^{\beta_2}]u\|_{L_2(B_{\rho_1 + \rho})} \leq C \sum_{i=1}^j j!/(j-i)! \|u, \rho_1 + \rho\|_{j-i}.$$

PROOF: The proof follows from Lemma 7 if we note that $[P, D_x^{\beta_1} D_y^{\beta_2}]$ is representable by a linear combination of terms in form $(D_x^{\gamma_1} x^{\alpha_3}) D_x^{\alpha_1 + \beta_1 - \gamma_1} D_y^{\alpha_2 + \beta_2}$ where $(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{N} \cap \mathbb{N}_0^3$, $1 \leq \gamma_1 \leq \min(\beta_1, \alpha_3)$ and the number of the nonzero terms in which x^{α_3} is differentiated γ_1 times is less than $C_j^{\gamma_{\alpha_3}}$ ($C_j^{\gamma_{\alpha_3}}$ are binomial coefficients). \square

Theorem 1. Let the hypoelliptic differential operator $P(x, D)$ from (1) with $\mathcal{N}(P)$ in form (2) satisfy in any neighborhood U of 0

$$\sum_{(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{N}(P) \cap \mathbb{N}_0^3} \|x^{\alpha_3} D_x^{\alpha_1} D_y^{\alpha_2} \psi\|_{L_2} \leq \|P(x, D)\psi\|_{L_2}, \quad \psi \in C_0^\infty(U).$$

Then all solutions of equation $P(x, D) = f$ belong to the class $G^{(\lambda_1, 1)}(V)$, with $V \subset\subset U$, $(0, 0) \in V$, where $f \in G^{(\lambda_1, 1)}(U)$.

PROOF: We take $U = B_3$, $V = B_1$. Let $\rho > 0$, $\rho_1 > 1$, $\rho_1 + \rho < 2$, then for any multi-indices $\beta \in \mathbb{N}_0^2$ from (3) we obtain

$$\|D_x^{\beta_1} D_y^{\beta_2} u, \rho_1\| \leq \|\varphi D_x^{\beta_1} D_y^{\beta_2} u, 2\| \leq C \|P(x, D)(\varphi D_x^{\beta_1} D_y^{\beta_2} u)\|_{L_2},$$

where φ is from Lemma 5. Since

$$\begin{aligned} & \|P(x, D)(\varphi D_x^{\beta_1} D_y^{\beta_2} u)\|_{L_2} \\ &= \|\varphi D_x^{\beta_1} D_y^{\beta_2} f\|_{L_2} + \|[P, \varphi] D_x^{\beta_1} D_y^{\beta_2} u\|_{L_2} + \|[P, D_x^{\beta_1} D_y^{\beta_2}]u\|_{L_2(B_{\rho_1 + \rho})} \end{aligned}$$

then for any natural j

$$\begin{aligned} \|\|u, \rho\|\|_j &\leq \max_{(\beta_1, \beta_2) \in \mathcal{M}^j \cap \mathbb{N}_0^2} \{ \|D_x^{\beta_1} D_y^{\beta_2} f\|_{L_2(B_2)} + \| [P, \varphi] D_x^{\beta_1} D_y^{\beta_2} u \|_{L_2} \\ &\quad + \| [P, D_x^{\beta_1} D_y^{\beta_2}] u \|_{L_2(B_{\rho_1+\rho})} \}. \end{aligned}$$

From condition $f \in G^{(\lambda_1, 1)}(U)$ and from Corollary 1, 2 for $j \geq 1$ with some constant $C_1 = C_1(f)$ we obtain

$$(5) \quad \|u, \rho_1\|_j \leq C(C_1^{j+1} j! + \sum_{i=1}^m \rho^{-i} \|u, \rho_1 + \rho\|_{j-i} + \sum_{i=1}^j j!/(j-i)! \|u, \rho_1 + \rho\|_{j-i}).$$

For any natural $s, j \leq s$ we denote

$$\omega_{s,j} = s^{-j} \|u, 2 - (j+1)/s\|_j.$$

Applying (5) with $\rho = 1/s, \rho_1 = 2 - j/s$, we obtain

$$(6) \quad \omega_{s,j} \leq C(C_2^j + \sum_{i=1}^m \omega_{s,j-i} + \sum_{i=1}^j \omega_{s,j-i})$$

with some constant C_2 . From (6) we obtain by induction $\omega_{s,j} \leq C_3^{j+1}$ for $j \leq s$ with some constant $C_3 > 1$. For $j = s$ we obtain $\|u, 2 - (s+1)/s\|_s \leq C_3^{s+1} s^s$, $s = 1, 2, \dots$. Since

$$\{\nu : \nu \in \mathbb{R}_+^2, \lambda_1 \nu_1 + \nu_2 \leq k - (1+h)m\} \subset \mathcal{M}^k \subset \{\nu : \nu \in \mathbb{R}_+^2, \lambda_1 \nu_1 + \nu_2 \leq k\}$$

for any $k \geq (1+h)m$, the proof is complete. \square

REFERENCES

- [1] Grushin V.V., *On a class of elliptic pseudodifferential operators degenerate on a submanifold*, Mat. Sbornik **84** (1971), 163–1295; Math. USSR Sbornik **13** (1971), 155–185.
- [2] Parenti C., Rodino L., *Parametrices for a class of pseudo differential operators, I, II*, Ann. Mat. Pura Appl. **125** (1980), 221–278.
- [3] Rodino L., *Gevrey hypoellipticity for a class of operators with multiple characteristics*, Asterisque **89-90** (1981), 249–262.
- [4] Tartakoff D.S., *Elementary proofs of analytic hypoellipticity for Δ_b and δ -Neumann problem*, in Analytic Solution of Partial Differential Equations, Asterisque **89-90** (1981), 85–116.
- [5] Volevich L.R., *Local regularity of the solutions of the quasi-elliptic systems* (in Russian), Mat. Sbornik **59** (1962), 3–52.

DEPARTMENT OF PHYSICS, YSU, 375025, YEREVAN, ARMENIA

E-mail: gaghakob@ysu.am

(Received March 31, 2003, revised September 30, 2003)