

Aleksander V. Arhangel'skii

$G_\delta$ -modification of compacta and cardinal invariants

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 47 (2006), No. 1, 95--101

Persistent URL: <http://dml.cz/dmlcz/119576>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## $G_\delta$ -modification of compacta and cardinal invariants

A.V. ARHANGEL'SKII

*Abstract.* Given a space  $X$ , its  $G_\delta$ -subsets form a basis of a new space  $X_\omega$ , called the  $G_\delta$ -modification of  $X$ . We study how the assumption that the  $G_\delta$ -modification  $X_\omega$  is homogeneous influences properties of  $X$ . If  $X$  is first countable, then  $X_\omega$  is discrete and, hence, homogeneous. Thus,  $X_\omega$  is much more often homogeneous than  $X$  itself. We prove that if  $X$  is a compact Hausdorff space of countable tightness such that the  $G_\delta$ -modification of  $X$  is homogeneous, then the weight  $w(X)$  of  $X$  does not exceed  $2^\omega$  (Theorem 1). We also establish that if a compact Hausdorff space of countable tightness is covered by a family of  $G_\delta$ -subspaces of the weight  $\leq c = 2^\omega$ , then the weight of  $X$  is not greater than  $2^\omega$  (Theorem 4). Several other related results are obtained, a few new open questions are formulated. Fedorchuk's hereditarily separable compactum of the cardinality greater than  $c = 2^\omega$  is shown to be  $G_\delta$ -homogeneous under CH. Of course, it is not homogeneous when given its own topology.

*Keywords:* weight, tightness,  $G_\delta$ -modification, character, Lindelöf degree, homogeneous space

*Classification:* 54A25, 54B10

Let  $\mathcal{T}$  be a topology on a set  $X$ . Then the family of all  $G_\delta$ -subsets of  $X$  is a base of a new topology on  $X$ , denoted by  $\mathcal{T}_\omega$  and called the  $G_\delta$ -modification of  $\mathcal{T}$ . The space  $(X, \mathcal{T}_\omega)$  is also denoted by  $X_\omega$  and is called the  $G_\delta$ -modification of the space  $(X, \mathcal{T})$ . Clearly, the  $G_\delta$ -modification  $X_\omega$  of any topological space is a  $P$ -space, that is, every  $G_\delta$ -subset of  $X_\omega$  is open in  $X_\omega$ .

In general, the space  $(X, \mathcal{T}_\omega)$  is very different from the space  $(X, \mathcal{T})$ . Many properties of  $(X, \mathcal{T})$ , such as compactness, Lindelöfness, paracompactness are easily lost under  $G_\delta$ -modifications. On the other hand, properties of the space can greatly improve under the operation of  $G_\delta$ -modification. For example, if  $(X, \mathcal{T})$  is first countable, then the space  $(X, \mathcal{T}_\omega)$  is discrete. Thus, no matter which first countable space  $(X, \mathcal{T})$  we take, the resulting space  $(X, \mathcal{T}_\omega)$  will be metrizable, zero-dimensional, Čech-complete and homogeneous! We see that the difference in properties between the spaces  $(X, \mathcal{T})$  and  $(X, \mathcal{T}_\omega)$  can indeed be tremendous!

Some interesting facts on  $G_\delta$ -modifications and on  $P$ -spaces were established in [12], where also a survey of what is known in this direction is given. See also [11].

It is our goal in this article to show that homogeneity of  $G_\delta$ -modification has a deep influence on the structure of the space itself and on the relationship between its cardinal invariants. Our main result in this direction (Theorem 1 below) is inspired by R. de la Vega's recent result that the weight of any homogeneous compact Hausdorff space of countable tightness is  $\leq 2^\omega$ . We generalize de la Vega's theorem as follows:

**Theorem 1.** *Let  $X$  be a compact Hausdorff space of countable tightness such that the  $G_\delta$ -modification  $X_\omega$  of  $X$  is homogeneous. Then the weight  $w(X)$  of  $X$ , as well as the weight of  $X_\omega$ , is not greater than  $2^\omega$ .*

PROOF: We claim that there is a non-empty open subspace  $U$  of  $X_\omega$  such that  $w(U) \leq 2^\omega$ . Indeed, since  $X$  is a non-empty compact Hausdorff space of countable tightness, there exists a non-empty  $G_\delta$ -subset  $U$  of  $X$  such that the weight of the subspace  $U$  of  $X$  is not greater than  $2^\omega$  ([2], [1]). Then  $U$  is an open subspace of  $X_\omega$  and the weight of the subspace  $U$  of  $X_\omega$  is also not greater than  $2^\omega$ . Since  $X_\omega$  is homogeneous, it follows that every point in  $X_\omega$  has an open neighbourhood  $Ox$  in  $X_\omega$  such that  $w(Ox) \leq 2^\omega$ .

According to a result of E.G. Pytkeev [14], the Lindelöf degree of the  $G_\delta$ -modification of any compact Hausdorff space of countable tightness does not exceed  $2^\omega$  (see Theorem 4 in [14]). Therefore,  $l(X_\omega) \leq 2^\omega$ . Since the local weight of  $X_\omega$  does not exceed  $2^\omega$ , it follows that there exists an open covering  $\gamma$  of  $X_\omega$  such that  $w(U) \leq 2^\omega$ , for each  $U \in \gamma$ , and  $|\gamma| \leq 2^\omega$ . Fixing a base of cardinality  $\leq 2^\omega$  in each  $U \in \gamma$ , and taking the union of these bases, we obtain a base of cardinality  $\leq 2^\omega$  in  $X_\omega$ . Thus,  $w(X_\omega) \leq 2^\omega$ . Since,  $X$  is a continuous image of  $X_\omega$ , we have  $nw(X) \leq w(X_\omega) \leq 2^\omega$ . However, since  $X$  is compact,  $w(X) = nw(X) \leq 2^\omega$  ([9]).  $\square$

This theorem immediately implies that the cardinality of every first countable compact Hausdorff space does not exceed  $2^\omega$  [Arh2]. Indeed, the tightness of first countable spaces is countable, and, obviously, if the weight of a first countable Hausdorff space is  $\leq 2^\omega$ , then the cardinality of  $X$  is also not greater than  $2^\omega$ . Theorem 1 also implies de la Vega's result that the weight of any homogeneous compact Hausdorff space of countable tightness is  $\leq 2^\omega$ , since the  $G_\delta$ -modification of a homogeneous space is homogeneous.

A space  $Y$  is *power-homogeneous* if  $Y^\tau$  is homogeneous, for some  $\tau > 0$  (see [4]). Weakening one of the assumptions in Theorem 1, we arrive at a weaker conclusion:

**Theorem 2.** *Let  $X$  be a compact Hausdorff space of countable tightness such that the  $G_\delta$ -modification of  $X$  is power-homogeneous. Then the character of  $X$  is not greater than  $2^\omega$ .*

PROOF: Take any non-empty  $G_\delta$ -subset  $Y$  of  $X$ . There exists a non-empty  $G_\delta$ -subset  $U$  of  $Y$  such that the weight of the subspace  $U$  of the space  $X$  is not greater than  $2^\omega$  ([2], [1]). Then  $U$  is an open subspace of  $X_\omega$  and the weight

of the subspace  $U$  of  $X_\omega$  is also not greater than  $2^\omega$ . It follows that the set  $Z$  of all  $x \in X$  such that the character of  $x$  in  $X_\omega$  is not greater than  $2^\omega$  is dense in the space  $X_\omega$ . Since  $X_\omega$  is power-homogeneous and  $Z \neq \emptyset$ , it follows from Theorem 7 in [4] that the set  $M$  of all  $G_c$ -points in  $X_\omega$  is closed. Obviously,  $Z \subset M$ . Therefore,  $M = X$ ; thus, each  $x \in X$  is a  $G_c$ -point in  $X_\omega$ .

Fix an arbitrary  $a \in X$ . According to Pytkeev's theorem (see the proof of Theorem 1), the Lindelöf degree of  $X_\omega$  is not greater than  $c = 2^\omega$ . Put  $A = X \setminus \{a\}$ . Since  $a$  is a  $G_c$ -point in  $X_\omega$ , it follows that  $l(A) \leq 2^\omega$ , where  $A$  is considered as a subspace of  $X_\omega$ . Since the identity mapping of  $X_\omega$  onto  $X$  is continuous, we conclude that the Lindelöf degree of  $A$ , considered as a subspace of  $X$ , does not exceed  $2^\omega$  as well. This implies that  $a$  is a  $G_c$ -point in  $X$ . Since  $X$  is compact and Hausdorff, it follows that the character of  $X$  at  $a$  is not greater than  $2^\omega$  ([9]).  $\square$

**Theorem 3.** *Let  $X$  be a sequential Hausdorff compact space such that the  $G_\delta$ -modification of  $X$  is power-homogeneous. Then  $|X| \leq 2^\omega$ .*

PROOF: It follows from Theorem 2 that  $\chi(X) \leq 2^\omega$ . However, the cardinality of every sequential Hausdorff compact space such that  $\chi(X) \leq 2^\omega$  does not exceed  $2^\omega$  (see [2]).  $\square$

The last result generalizes Corollary 3.8 in [5] and an earlier result on the cardinality of homogeneous compact sequential spaces in [2].

The technique of  $G_\delta$ -modification can be used to obtain some addition theorems for the weight that do not involve the assumption of homogeneity. In particular, we have:

**Theorem 4.** *Let  $X$  be a compact Hausdorff space of countable tightness, and suppose that  $X$  is covered by a family  $\gamma$  of  $G_\delta$ -subsets such that the weight of  $P$  is not greater than  $2^\omega$ , for each  $P \in \gamma$ . Then the weight of  $X$  is not greater than  $2^\omega$ .*

PROOF: The proof is close to the proof of Theorem 1. Consider the  $G_\delta$ -modification  $X_\omega$  of  $X$ . The family  $\gamma$  is an open covering of  $X_\omega$ , and the weight of each  $P \in \gamma$ , interpreted as a subspace of  $X_\omega$ , is not greater than  $2^\omega$ . By Pytkeev's theorem (see the proof of Theorem 1), the Lindelöf degree of  $X_\omega$  is not greater than  $c = 2^\omega$ . Therefore, the weight of  $X_\omega$  is not greater than  $2^\omega$  (to get an appropriate base of  $X_\omega$ , just take the union of the bases of cardinality  $\leq 2^\omega$  of elements of  $\gamma$ ). Since  $X$  is a continuous image of  $X_\omega$ , we have  $nw(X) \leq w(X_\omega) \leq 2^\omega$ . However,  $X$  is compact. Hence,  $w(X) = nw(X) \leq 2^\omega$ .  $\square$

For some results related to Theorem 4 see [15] and [6].

The assumption of countable tightness in the last statement can be replaced by some other conditions.

**Theorem 5.** *Let  $X$  be a scattered compact Hausdorff space covered by a family  $\gamma$  of  $G_\delta$ -subsets such that the weight of  $P$  is not greater than  $2^\omega$ , for each  $P \in \gamma$ . Then the weight of  $X$  does not exceed  $2^\omega$ .*

PROOF: The Lindelöf degree of the  $G_\delta$ -modification  $X_\omega$  of the space  $X$  does not exceed  $\omega$  ([13]). Since  $\gamma$  is an open covering of  $X_\omega$ , we can assume that  $\gamma$  is countable. It follows that  $w(X_\omega) \leq 2^\omega$ , which implies that  $nw(X) \leq w(X_\omega) \leq 2^\omega$ . Finally, since  $X$  is compact, we have  $w(X) = nw(X) \leq 2^\omega$ .  $\square$

The proof of the next result should be clear by now:

**Theorem 6.** *Let  $X$  be a scattered space. Then the  $G_\delta$ -modification  $X_\omega$  of  $X$  is power-homogeneous if and only if the pseudocharacter of  $X$  is countable (that is, if and only if the  $G_\delta$ -modification of  $X$  is discrete).*

**Problem 7.** *Suppose that  $X$  is a compact Hausdorff space covered by a family  $\gamma$  of  $G_\delta$ -subsets  $P$  such that the weight of  $P$  is not greater than  $2^\omega$ , for each  $P \in \gamma$ . Is the weight of  $X$  not greater than  $2^\omega$ ?*

**Problem 8** (Arhangel'skii, Buzyakova). *Let  $X$  be a compact Hausdorff space of countable tightness such that the character of  $X$  does not exceed  $2^\omega$ . Is the weight of  $X$  not greater than  $2^\omega$ ?*

Consistently the answer to the last question is “yes”. Indeed, it was shown in [7] to be consistent with ZFC to assume that every compact Hausdorff space of countable tightness is sequential. It remains to apply the following result from [2]: the cardinality of every sequential Hausdorff compact space such that  $\chi(X) \leq 2^\omega$  does not exceed  $2^\omega$ .

Closely related to Problem 8 is the following question: Let  $X$  be a compact Hausdorff space of countable tightness such that the  $G_\delta$ -modification of  $X$  is homogeneous. Is  $|X| \leq 2^\omega$ ? The answer to this question is independent of ZFC. Under Proper Forcing Axiom (PFA) (for the discussion of (PFA) see [8]) the answer is “yes”. In fact, we can prove a stronger statement:

**Theorem 9.** *Assume (PFA), and let  $X$  be a Hausdorff compact space of countable tightness such that the  $G_\delta$ -modification of  $X$  is power-homogeneous. Then  $X$  is first countable (and hence,  $|X| \leq 2^\omega$  and  $w(X) \leq 2^\omega$ ).*

PROOF: A. Dow has shown in [Dow] that under (PFA) every non-empty compact Hausdorff space of countable tightness has a point of first countability. It follows easily from this result that, under (PFA), the set of isolated points is dense in the  $G_\delta$ -modification  $X_\omega$  of the compactum  $X$ .

Since  $X_\omega$  is power-homogeneous, it follows from Theorem 7 in [4] that the set  $M$  of all  $G_\delta$ -points in  $X_\omega$  is closed. Therefore,  $M = X$ , that is, each  $x \in X$  is a  $G_\delta$ -point in  $X_\omega$ . Since  $X_\omega$  is a  $P$ -space, we conclude that the space  $X_\omega$  is discrete. Hence, the pseudocharacter of the space  $X$  is countable. Since  $X$  is compact and Hausdorff, it follows that  $X$  is first countable.  $\square$

On the other hand, we have the following result:

**Theorem 10 (CH).** *Let  $X$  be a hereditarily separable compact Hausdorff space without points of first countability. Then the  $G_\delta$ -modification of  $X$  is homogeneous.*

This theorem will follow from a more general result below. Notice that Fedorchuk has constructed [10] a consistent example of a hereditarily separable, nowhere first countable, compact Hausdorff space  $X$  such that the cardinality of  $X$  is greater than  $2^\omega$ . In the model of Set-theory he considered (CH) was also satisfied.

**Theorem 11 (CH).** *Let  $X$  be a compact Hausdorff space of the weight  $\omega_1$  such that the character of  $X$  at each point is exactly  $\omega_1$ . Then the  $G_\delta$ -modification  $X_\omega$  of  $X$  is homeomorphic to the  $G_\delta$ -modification of the compactum  $D^{\omega_1}$ .*

Fix a set  $A$  of the cardinality  $\omega_1 = c = 2^\omega$ , give  $A$  the discrete topology, and let  $B$  be the  $G_\delta$ -modification of the product space  $A^{\omega_1}$ .

*Claim 1:* The  $G_\delta$ -modification of  $D^{\omega_1}$  is homeomorphic to the space  $B$ .

This is obvious.

By Claim 1, it is enough to prove that  $X_\omega$  is homeomorphic to  $B$ . For that, we need the following lemma:

**Lemma 12.** *Let  $X$  be a non-scattered compact Hausdorff space. Then there exists a disjoint covering  $\gamma$  of  $X$  by non-empty closed  $G_\delta$ -sets such that  $|\gamma| = 2^\omega$ .*

PROOF: Since  $X$  is not scattered, there exists a continuous mapping  $f$  of  $X$  onto the closed interval  $I = [0, 1]$  (see [9]). Then  $\gamma = \{f^{-1}(y) : 0 \leq y \leq 1\}$  is, clearly, the covering we are looking for.  $\square$

Below we will need the following slightly stronger version of Lemma 12:

**Lemma 13.** *Let  $X$  be a non-scattered compact Hausdorff space and  $F_0$  be a closed  $G_\delta$ -subset of  $X$ . Then there exists a disjoint covering  $\gamma_1$  of  $X$  by non-empty closed  $G_\delta$ -sets such that  $|\gamma_1| = 2^\omega$  and  $F_0 = \bigcup \eta$ , for some subfamily  $\eta$  of  $\gamma_1$ .*

PROOF: We can fix a continuous real-valued function  $g$  on  $X$  such that  $g^{-1}(0) = F_0$ , since  $X$  is normal. Take also a disjoint covering  $\gamma$  of  $X$  by closed  $G_\delta$ -subsets such that  $|\gamma| = 2^\omega$  (this is possible by Lemma 12). Now let  $\gamma_1$  be the family  $\{g^{-1}(a) \cap P : a \in \mathbb{R}, P \in \gamma\} \setminus \{\emptyset\}$ , where  $\mathbb{R}$  is the set of reals. Obviously,  $\gamma_1$  is the covering we are looking for.  $\square$

PROOF OF THEOREM 11: A standard construction by transfinite recursion along  $\omega_1$ , using (CH) and Lemmas 12 and 13, provides us with a transfinite sequence  $\{\gamma_\alpha : \alpha < \omega_1\}$  of disjoint coverings of  $X$  by closed non-empty  $G_\delta$ -subsets of  $X$  such that the following conditions are satisfied:

- 1)  $\gamma_\beta$  refines  $\gamma_\alpha$ , whenever  $\alpha < \beta < \omega_1$ ;

- 2) for each  $P \in \gamma_\alpha$ , the cardinality of the family  $\eta_P = \{F \in \gamma_{\alpha+1} : F \subset P\}$  is  $\omega_1$ ;
- 3) the family  $S = \bigcup \{\gamma_\alpha : \alpha < \omega_1\}$  is a network of the space  $X$ .

Observe that compactness of  $X$  and the above conditions ensure that the following condition is satisfied:

- 4) for every uncountable centered family  $\xi$  of elements of  $S$ , the intersection of  $\xi$  consists of exactly one point  $x_\xi$ ,  $\xi$  is a network of  $X$  at  $x$ , and  $\xi$  is a base of the  $G_\delta$ -modification  $X_\omega$  at  $x$ .

Note, that elements of  $S$  are open-closed subsets of  $X_\omega$ , and that if  $\xi \subset S$  is countable, then either  $\bigcap \xi = \emptyset$  or the cardinality of  $\bigcap \xi$  is  $c = \omega_1$ .

The above properties of the family  $\{\gamma_\alpha : \alpha < \omega_1\}$  allow to establish a homeomorphism between the space  $X_\omega$  and the space  $B$  in an obvious routine way.  $\square$

**Corollary 14 (CH).** *Let  $X$  be a compact Hausdorff space of the weight  $\omega_1$  such that the character of  $X$  at each point is exactly  $\omega_1$ . Then the  $G_\delta$ -modification  $X_\omega$  of  $X$  is homogeneous. Furthermore,  $X_\omega$  is homeomorphic to a topological group.*

PROOF: Indeed, by Theorem 11  $X_\omega$  is homeomorphic to the  $G_\delta$ -modification  $B$  of the compactum  $D^{\omega_1}$ . However, the space  $B$  is homogeneous, since  $D^{\omega_1}$  is homogeneous. Hence,  $X_\omega$  is homogeneous as well. In fact,  $B$  is homeomorphic to a topological group, since  $D^{\omega_1}$  is a topological group.  $\square$

**Problem 15.** *Can (CH) be dropped in the above statement?*

The following long standing problems posed in [3], [1], [2] remain open:

**Problem 16.** *Is it true in ZFC that every homogeneous compact sequential space is first countable?*

**Problem 17.** *Is it true in ZFC that every homogeneous compact space of countable tightness is first countable?*

**Acknowledgment.** I am grateful to Professor Raushan Z. Buzyakova for several very helpful remarks on the subject of this paper.

#### REFERENCES

- [1] Arhangel'skii A.V., *On cardinal invariants*, In: General Topology and its Relations to Modern Analysis and Algebra 3. Proceedings of the Third Prague Topological Symposium, 1971, 37–46. Academia Publishing House, Czechoslovak Academy of Sciences, Prague, 1972.
- [2] Arhangel'skii A.V., *Structure and classification of topological spaces and cardinal invariants*, Russian Math. Surveys **33** (1978), 33–96.
- [3] Arhangel'skii A.V., *Topological homogeneity, topological groups and their continuous images*, Russian Math. Surveys **42** (1987), 83–131.
- [4] Arhangel'skii A.V., *Homogeneity of powers of spaces and the character*, Proc. Amer. Math. Soc. **133** (2005), 2165–2172.

- [5] Arhangel'skii A.V., *A weak algebraic structure on topological spaces and cardinal invariants*, Topology Proc. **28** (2004), no. 1, 1–18.
- [6] Arhangel'skii A.V., van Mill J., Ridderbos G.J., *A new bound on the cardinality of power homogeneous compacta*, Houston J. Math., to appear.
- [7] Balogh Z., *On compact Hausdorff spaces of countable tightness*, Proc. Amer. Math. Soc. **105** (1989), 755–764.
- [8] Dow A., *An introduction to applications of elementary submodels to topology*, Topology Proc. **13** (1988), 17–72.
- [9] Engelking R., *General Topology*, PWN, Warszawa, 1977.
- [10] Fedorchuk V., *On the cardinality of hereditarily separable compact Hausdorff spaces*, Soviet Math. Dokl. **16** (1975), 651–655.
- [11] Kunen K., *Box products of compact spaces*, Trans. Amer. Math. Soc. **240** (1978), 307–316.
- [12] Levy R., Rice M.D., *Normal  $P$ -spaces and the  $G_\delta$ -topology* Colloq. Math. **44** (1981), 227–240.
- [13] Meyer P.R., *The Baire order problem for compact spaces*, Duke Math. J. **33** (1966), 33–40.
- [14] Pytkeev E.G., *About the  $G_\lambda$ -topology and the power of some families of subsets on compacta*, Colloq. Math. Soc. Janos Bolyai, 41. Topology and Applications, Eger (Hungary), 1983, pp. 517–522.
- [15] de la Vega R., *A new bound on the cardinality of homogeneous compacta*, Topology Appl., to appear.

DEPARTMENT OF MATHEMATICS, OHIO UNIVERSITY, ATHENS, OHIO 45701, USA

*E-mail:* arhangel@math.ohiou.edu

(Received September 13, 2005)