

Konrad Pióro

On a property of neighborhood hypergraphs

Commentationes Mathematicae Universitatis Carolinae, Vol. 47 (2006), No. 1, 149--154

Persistent URL: <http://dml.cz/dmlcz/119581>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On a property of neighborhood hypergraphs

KONRAD PIÓRO

Abstract. The aim of the paper is to show that no simple graph has a proper subgraph with the same neighborhood hypergraph. As a simple consequence of this result we infer that if a clique hypergraph \mathcal{G} and a hypergraph \mathcal{H} have the same neighborhood hypergraph and the neighborhood relation in \mathcal{G} is a subrelation of such a relation in \mathcal{H} , then \mathcal{H} is inscribed into \mathcal{G} (both seen as coverings). In particular, if \mathcal{H} is also a clique hypergraph, then $\mathcal{H} = \mathcal{G}$.

Keywords: graph, neighbor, neighborhood hypergraph, clique hypergraph

Classification: 05C99, 05C69, 05C65

Recall (see e.g. [1]) that a hypergraph $\mathcal{G} = (V, \mathcal{E})$ consists of a finite set V of vertices and a finite sequence \mathcal{E} of hyperedges, where each hyperedge is a non-empty subset of V , and the union of all hyperedges is V (note that a hypergraph may have multiple hyperedges). A hypergraph is *simple*, if no hyperedge is contained in another hyperedge.

With an ordinary graph G at least two hypergraphs can be associated. The first consists of all maximal cliques of G and is called the clique hypergraph. These hypergraphs form an important subclass of hypergraphs. For example, they are related with the Helly property (see [1]), and they also appear in the clique-transversal problem (see [4]), and consequently in graph coloring problems (see e.g. [5]).

The second hypergraph associated with G is formed by neighborhoods of vertices. Recall (see [1]) that two vertices of G are neighbors if they are adjacent or equal. The set of all neighbors of a vertex v is denoted by $N_G(v)$ and called the neighborhood of v . Next, we take all pairwise different neighborhoods in G to obtain a new hypergraph $\mathcal{N}(G)$ on the vertex set of G , called the neighborhood hypergraph of G . $\mathcal{N}(G)$ has no multiple hyperedges, but, in general, $\mathcal{N}(G)$ is not simple. Of course, hypergraphs $\mathcal{N}(H)_{\max}$ (consisting of all maximal hyperedges of $\mathcal{N}(G)$ with respect to inclusion) and $\mathcal{N}(G)_{\min}$ (consisting of all minimal hyperedges of $\mathcal{N}(G)$) are simple, but they play no important role here.

Theorem 1. *Let G be a simple graph and H its subgraph. If $\mathcal{N}(H) = \mathcal{N}(G)$, then $H = G$.*

PROOF: Take a vertex v of H such that $N_H(v)$ is maximal up to inclusion. Let X_H be the set of all vertices w of H such that $N_H(w) = N_H(v)$, and X_G be the set of all vertices w of G such that $N_G(w) = N_H(v)$.

Since $N_H(v)$ corresponds to a maximal (up to inclusion) hyperedge of $\mathcal{N}(H) = \mathcal{N}(G)$ and H is a subgraph of G , we infer that

$$X_H \subseteq X_G,$$

in particular $N_H(v) = N_G(v)$.

Assume that there is a vertex $w \in X_G \setminus X_H$. Since $N_G(w) = N_H(v)$ and $w \in N_G(w)$ (by the definition), the vertices v and w are adjacent in H , thus also in G .

Since $N_H(w) \subseteq N_G(w) = N_H(v)$ and $N_H(w) \neq N_H(v)$ (by the assumption), there exists a vertex u such that

$$u \in N_H(v) = N_G(w) \quad \text{and} \quad u \notin N_H(w).$$

Then

$$N_H(u) \neq N_G(u).$$

Hence and by the equality $\mathcal{N}(H) = \mathcal{N}(G)$, there is a vertex u' such that

$$N_H(u) = N_G(u').$$

Then $v \in N_H(u) = N_G(u')$, i.e. the vertices v and u' are adjacent in G .

On the other hand,

$$w \notin N_H(u) = N_G(u'),$$

so

$$u' \notin N_G(w) = N_H(v) = N_G(v),$$

i.e. the vertices u' and v are not adjacent in G . This contradiction implies

$$X_H = X_G.$$

Observe now that the vertices of $X = X_H = X_G$ form a clique in both the graphs H and G and the sets of neighbors of every vertex of X in the rest of the graphs H and G are the same. Thus to end the proof it is sufficient to apply the induction (on the order of graph) to the pair of graphs $H \setminus X$ and $G \setminus X$ (note that they may have isolated vertices, but it is not a problem). \square

Recall that a simple hypergraph \mathcal{G} is said to be a *clique hypergraph*, if it is the clique hypergraph of some graph G . Observe that neighborhoods of each vertex v in \mathcal{G} and G are the same, in particular $\mathcal{N}(\mathcal{G}) = \mathcal{N}(G)$ (where neighbors in a hypergraph are defined analogously as for a graph). Moreover, G is uniquely

determined (therefore it will be sometimes denoted by $G_{\mathcal{G}}$). Because two different vertices of G are adjacent if and only if they are both contained in a hyperedge of \mathcal{G} . Hence it also easy follows (see [1]) that a simple hypergraph $\mathcal{G} = (V, \mathcal{E})$ is a clique hypergraph if and only if for each subset A of V , the following condition holds:

(C_2) if every pair of vertices of A belongs to some hyperedge of \mathcal{G} , then A is contained in a hyperedge of \mathcal{G} .

(Hypergraphs satisfying (C_2), not necessarily simple, were called conformal by Berge in [1]. However, today the concept of conformality has a slightly different meaning (see e.g. [6]).)

(C_2) is related with the Helly property (see [1]). More precisely, a hypergraph $\mathcal{G} = (V, \mathcal{E})$ has the Helly property (i.e. for any $\mathcal{F} \subseteq \mathcal{E}$, if any two hyperedges in \mathcal{F} have a non-empty intersection, then the intersection of \mathcal{F} is also non-empty) if and only if its dual \mathcal{G}^* satisfies (C_2). $\mathcal{G}^* = (\mathcal{E}, V^*)$ is the hypergraph whose vertices are hyperedges of \mathcal{G} and the set of hyperedges is $V^* = \{\mathcal{G}(v) : v \in V\}$, where $\mathcal{G}(v) = \{E \in \mathcal{E} : v \in E\}$. Gilmore's Theorem (see Chapter 1, §7 in [1]) gives the following necessary and sufficient condition for a hypergraph \mathcal{G} to satisfy (C_2): for every three hyperedges E_1, E_2, E_3 of \mathcal{G} , there is a hyperedge of \mathcal{G} containing the set $(E_1 \cap E_2) \cup (E_2 \cap E_3) \cup (E_3 \cap E_1)$. The condition can be easily translated into the Helly property (see [1]). This result have been generalized by Berge and Duchet in [3] (see also [1]) to hypergraphs with the k -Helly property (i.e. for any family \mathcal{F} of hyperedges of \mathcal{G} , if every subfamily of \mathcal{F} with at most k elements has a non-empty intersection, then \mathcal{F} also has a non-empty intersection). The k -Helly property corresponds with the condition (C_k) obtained from (C_2) by replacing "every pair" with "every subset with at most k vertices".

We say that a hypergraph \mathcal{H} is *inscribed into* a hypergraph \mathcal{G} if for any hyperedge F of \mathcal{H} there is a hyperedge E of \mathcal{G} such that $F \subseteq E$. It is just a reformulation of the well-known notion for covering in the case of hypergraphs.

Theorem 2. *Let \mathcal{G} be a clique hypergraph and \mathcal{H} be an arbitrary hypergraph with the same vertex set such that*

- (*) $N_{\mathcal{G}}(v) \subseteq N_{\mathcal{H}}(v)$ for each vertex v ,
- (**) $\mathcal{N}(\mathcal{G}) = \mathcal{N}(\mathcal{H})$.

Then \mathcal{H} is inscribed into \mathcal{G} .

PROOF: Take an auxiliary graph H with the same vertex set as \mathcal{H} such that two different vertices of H are adjacent if and only if they are contained in a common hyperedge of \mathcal{H} . Then $N_H(v) = N_{\mathcal{H}}(v)$ for any vertex v . Hence and by (*) we first infer that the graph $G_{\mathcal{G}}$ is a subgraph of H . Secondly, $\mathcal{N}(G_{\mathcal{G}}) = \mathcal{N}(H)$ by (**). Thus by Theorem 1 we obtain $G_{\mathcal{G}} = H$, i.e. \mathcal{G} is the clique hypergraph of H . It easily implies that \mathcal{H} is inscribed into \mathcal{G} . \square

By the above proof we obtain in particular that for any hypergraph \mathcal{H} there exists exactly one clique hypergraph \mathcal{H}' with the same vertex set such that \mathcal{H} is

inscribed into \mathcal{H}' and $N_{\mathcal{H}'}(v) = N_{\mathcal{H}}(v)$ for each vertex v (it is sufficient to take the graph H for \mathcal{H} as above and its clique hypergraph).

This fact and Theorem 2 (because the relation “to be inscribed into” is a partial order for simple hypergraphs) imply that \mathcal{G} is a clique hypergraph if and only if for each simple hypergraph \mathcal{H} with the same vertex set, if \mathcal{G} is inscribed into \mathcal{H} and $\mathcal{N}(\mathcal{H}) = \mathcal{N}(\mathcal{G})$, then $\mathcal{H} = \mathcal{G}$. In particular

Corollary 3. *Let \mathcal{G} and \mathcal{H} be clique hypergraphs with the same vertex set satisfying (*) and (**). Then $\mathcal{G} = \mathcal{H}$.*

By Theorem 2 we obtain also that if a clique hypergraph \mathcal{G} is a subhypergraph of a hypergraph \mathcal{H} and $\mathcal{N}(\mathcal{G}) = \mathcal{N}(\mathcal{H})$, then \mathcal{H} is inscribed into \mathcal{G} . In particular, if \mathcal{H} is simple, then $\mathcal{G} = \mathcal{H}$.

Now we translate the above results for hypergraphs having the Helly property. Observe that Theorem 2 holds also for hypergraphs satisfying (C_2) . Because if \mathcal{G} is such a hypergraph, then \mathcal{G}_{\max} is a clique hypergraph, and also $N_{\mathcal{G}_{\max}}(v) = N_{\mathcal{G}}(v)$ for any vertex v .

For hypergraphs $\mathcal{G} = (V, (E_1, \dots, E_n))$ and $\mathcal{H} = (W, (E'_1, \dots, E'_n))$ we will “assume” in the results below that \mathcal{G}^* and \mathcal{H}^* (and also $\mathcal{N}(\mathcal{G}^*)$ and $\mathcal{N}(\mathcal{H}^*)$) have the same vertex set $\{E_1, \dots, E_n\}$. Say more formally, we identify hyperedges E_i and E'_i , i.e. the equality $\mathcal{G}^* = \mathcal{H}^*$ denotes that the natural correspondence $E_i \mapsto E'_i$ forms an isomorphism between these hypergraphs.

Corollary 4. *Let $\mathcal{G} = (V, (E_1, \dots, E_n))$ be a hypergraph with the Helly property. Let $\mathcal{H} = (W, (E'_1, \dots, E'_n))$ be a hypergraph satisfying*

- (*) for any $1 \leq i, j \leq n$, $E_i \cap E_j \neq \emptyset \implies E'_i \cap E'_j \neq \emptyset$,
- (**) $\mathcal{N}(\mathcal{H}^*) = \mathcal{N}(\mathcal{G}^*)$.

Then for each $w \in W$, there is $v \in V$ such that for any $1 \leq i \leq n$,

$$w \in E'_i \implies v \in E_i.$$

PROOF: (*) implies $N_{\mathcal{G}^*}(E_i) \subseteq N_{\mathcal{H}^*}(E'_i)$ for each $i = 1, 2, \dots, n$. Hence, \mathcal{H}^* is inscribed into \mathcal{G}^* . This implies the thesis. \square

The implication in the above result cannot be replaced by the equivalence. Take the following two hypergraphs $\mathcal{G} = (\{1, 2, 3, 4\}, (\{1, 2\}, \{2, 3\}, \{3, 4\}))$ and $\mathcal{H} = (\{1, 2, 3, 4\}, (\{1, 2\}, \{2, 3, 5\}, \{3, 4\}))$. Then \mathcal{G} and \mathcal{H} satisfy the conditions (*) and (**), and \mathcal{G} has the Helly property. On the other hand, $\mathcal{H}(5) = \{\{2, 3, 5\}\}$, and $\mathcal{G}(2) = \{\{1, 2\}, \{2, 3\}\}$, $\mathcal{G}(3) = \{\{2, 3\}, \{3, 4\}\}$.

Take a hypergraph $\mathcal{G} = (V, (E_1, \dots, E_n))$ and note that \mathcal{G}^* is simple if and only if for each vertices $v, w \in V$, the following condition holds:

$$(DS) \quad \{E_i: v \in E_i\} \subseteq \{E_j: w \in E_j\} \implies v = w.$$

Thus by Corollary 3 we obtain (because $(\mathcal{G}^*)^* = \mathcal{G}$):

Corollary 5. *Let hypergraphs with the Helly property $\mathcal{G} = (V, (E_1, \dots, E_n))$ and $\mathcal{H} = (W, (E'_1, \dots, E'_n))$ satisfy (DS) and (*), (**) of Corollary 4. Then $\mathcal{G} = \mathcal{H}$ (strictly formally, \mathcal{G} and \mathcal{H} are isomorphic).*

Using the last corollary of Theorem 1 (i.e. its modified version in which we assume that \mathcal{G} satisfies (C_2)) we can also show that if a hypergraph \mathcal{G} having the Helly property is a subhypergraph of a hypergraph \mathcal{H} and $\mathcal{N}(\mathcal{G}^*) = \mathcal{N}(\mathcal{H}^*)$, then \mathcal{H} has also the Helly property. If \mathcal{H} satisfies additionally (DS), then $\mathcal{H} = \mathcal{G}$.

Observe that to a given hypergraph $\mathcal{G} = (V, (E_1, \dots, E_n))$ new vertices can be added in such a way that the obtained hypergraph has the Helly property. More precisely, there is a hypergraph $\mathcal{G}' = (V', (E'_1, \dots, E'_n))$ such that

- (i) $E_i \subseteq E'_i$ for $i = 1, \dots, n$,
- (ii) for each $1 \leq i, j \leq n$, $E'_i \cap E'_j \neq \emptyset \iff E_i \cap E_j \neq \emptyset$,
- (iii) \mathcal{G}' has the Helly property.

Take the dual hypergraph \mathcal{G}^* , and the graph G with vertices E_1, \dots, E_n such that E_i and E_j ($i \neq j$) are adjacent if and only if they both belong to a hyperedge of \mathcal{G}^* . Next, take the hypergraph \mathcal{H} consisting of all maximal cliques of G and all hyperedges of \mathcal{G}^* . Then \mathcal{G}^* is inscribed into \mathcal{H} , so \mathcal{H}_{\max} is a clique hypergraph, which implies that \mathcal{H} satisfies (C_2) . Moreover, $N_{\mathcal{H}}(E_i) = N_G(E_i) = N_{\mathcal{G}^*}(E_i)$ for each $i = 1, \dots, n$. Thus it is sufficient to take $\mathcal{G}' = \mathcal{H}^*$.

Now we show that the assumptions of Theorems 1 and 2 (thus also their corollaries) are necessary. First, the following graphs $G = (\{1, 2\}, \{2, 3\}, \{3, 4\}, \{1, 4\})$ and $H = (\{1, 3\}, \{3, 4\}, \{2, 4\}, \{1, 2\})$ are different, but they have the same neighborhood hypergraph (because $\mathcal{N}(G)$ and $\mathcal{N}(H)$ consist of all three-element subsets of $\{1, 2, 3, 4\}$). Further, the clique hypergraphs of G and H are equal to G and H , respectively.

Secondly, take the following hypergraphs $\mathcal{G} = (\{1, 5, 6, 7\}, \{1, 4, 5, 7\}, \{2, 3, 4, 7\})$ and $\mathcal{H} = (\{1, 5, 6, 7\}, \{1, 2, 4, 5, 7\}, \{2, 3, 4, 7\})$. It is easy to see that they are clique hypergraphs. \mathcal{G} and \mathcal{H} satisfy (*) of Theorem 2, and (**) does not hold, since $N_{\mathcal{G}}(1) = \{1, 4, 5, 6, 7\} \notin \mathcal{N}(\mathcal{H})$. On the other hand, $\mathcal{N}(\mathcal{G})_{\max} = \mathcal{N}(\mathcal{H})_{\max}$ (because they have exactly one hyperedge $N_{\mathcal{G}}(7) = N_{\mathcal{H}}(7) = \{1, 2, \dots, 7\}$) and $\mathcal{N}(\mathcal{G})_{\min} = \mathcal{N}(\mathcal{H})_{\min}$ (because they have exactly two hyperedges $N_{\mathcal{G}}(3) = N_{\mathcal{H}}(3) = \{2, 3, 4, 7\}$ and $N_{\mathcal{G}}(6) = N_{\mathcal{H}}(6) = \{1, 5, 6, 7\}$). Observe also that $G_{\mathcal{G}}$ is a proper subgraph of $G_{\mathcal{H}}$ (where $G_{\mathcal{G}}$ and $G_{\mathcal{H}}$ are the graphs corresponding to \mathcal{G} and \mathcal{H}), although $\mathcal{N}(G_{\mathcal{H}})_{\max} = \mathcal{N}(G_{\mathcal{G}})_{\max}$ and $\mathcal{N}(G_{\mathcal{H}})_{\min} = \mathcal{N}(G_{\mathcal{G}})_{\min}$.

Finally observe that our results are not true for infinite graphs and hypergraphs. Let $A = \{a_i: i \in \mathbb{Z}\}$ and $B = \{b_i: i \in \mathbb{Z}\}$ be two infinite disjoint sets (where \mathbb{Z} is the set of all integers), and take

$$G_1 = \{\{a_i, a_j\}: i \neq j\} \cup \{\{b_i, b_j\}: i \neq j\} \cup \{\{a_i, b_j\}: j \leq i\},$$

$$G_2 = \{\{a_i, a_j\}: i \neq j\} \cup \{\{b_i, b_j\}: i \neq j\} \cup \{\{a_i, b_j\}: j \leq i - 1\}.$$

Then first G_2 is a proper subgraph of G_1 . Secondly, for each $i \in \mathbb{Z}$,

$$\begin{aligned} N_{G_1}(a_i) &= A \cup \{b_j: j \leq i\}, & N_{G_1}(b_i) &= B \cup \{a_j: j \geq i\}, \\ N_{G_2}(a_i) &= A \cup \{b_j: j \leq i-1\}, & N_{G_2}(b_i) &= B \cup \{a_j: j \geq i+1\}. \end{aligned}$$

Hence, $N_{G_2}(a_i) = N_{G_1}(a_{i-1}) \subseteq N_{G_1}(a_i)$ and $N_{G_2}(b_i) = N_{G_1}(b_{i+1}) \subseteq N_{G_1}(b_i)$. In particular, $\mathcal{N}(G_1) = \mathcal{N}(G_2)$.

By the above facts we have also that the clique hypergraphs \mathcal{G}_1 and \mathcal{G}_2 of the graphs G_1 and G_2 satisfy assumptions of Theorem 2. But they are not equal, because $G_1 \neq G_2$.

Acknowledgments. The author is indebted to the referee for suggestions that helped to improve the paper and simplify the proofs of the main results.

REFERENCES

- [1] Berge C., *Hypergraphs*, North-Holland, Amsterdam, 1989.
- [2] Berge C., *Graphs and Hypergraphs*, North-Holland, Amsterdam, 1973.
- [3] Berge C., Duchet P., *A generalization of Gilmore's theorem*, Recent Advances in Graph Theory, (Fiedler M., ed.), Academia, Prague, 1975, pp. 49–55.
- [4] Erdős P., Gallai T., Tuze Z., *Covering the cliques of a graph with vertices*, Discrete Math. **108** (1992), 279–289.
- [5] Jensen T.R., Toft B., *Graph Coloring Problems*, Wiley Interscience, New York, 1995.
- [6] Prisner E., *Intersection multigraphs of uniform hypergraphs*, Graphs Combin. **14** (1998), no. 4, 363–375.

INSTITUTE OF MATHEMATICS, WARSAW UNIVERSITY, UL. BANACHA 2, PL-02-097 WARSAW, POLAND

E-mail: kpioro@mimuw.edu.pl

(Received November 15, 2004, revised September 13, 2005)