

Shou Lin

A note on D-spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 47 (2006), No. 2, 313--316

Persistent URL: <http://dml.cz/dmlcz/119594>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A note on D-spaces

SHOU LIN

Abstract. Every semi-stratifiable space or strong Σ -space has a σ -cushioned $(\text{mod}k)$ -network. In this paper it is showed that every space with a σ -cushioned $(\text{mod}k)$ -network is a D-space, which is a common generalization of some results about D-spaces.

Keywords: D-spaces, $(\text{mod}k)$ -networks, cushioned collections, semi-stratifiable spaces

Classification: 54E18, 54D20

A *neighborhood assignment* for a topological space (X, τ) is a function $\phi : X \rightarrow \tau$ such that $x \in \phi(x)$ for each $x \in X$. A space X is said to be a D-space if, for each neighborhood assignment ϕ for X , there exists a closed discrete subset D of X such that $\{\phi(d) : d \in D\}$ covers X . The first published results on D-spaces appear in [5], where it is proved that finite products of Sorgenfrey lines are D-spaces. Several interesting questions on D-spaces were raised by E.K. van Douwen and W.F. Pfeffer in [6]. It is still an open problem whether every regular Lindelöf space is a D-space. It is also asked whether there exists a subparacompact or metacompact space which is not a D-space. These questions are still open.

In [1] the authors study D-property in classes of generalized metric spaces. It is known that semi-stratifiable spaces are D-spaces [2], and strong Σ -spaces are D-spaces [3]. Semi-stratifiable spaces need not be strong Σ -spaces, and strong Σ -spaces need not be semi-stratifiable spaces [8]. It is therefore natural to look for a “common denominator” to the results mentioned above.

Throughout this paper, all spaces are assumed to be T_1 . We refer the reader to [7] for notations and terminology not explicitly given here.

Definition 1. A space X is said to be a *semi-stratifiable space* [4] if, for each open set U of X , one can assign a sequence $\{F(n, U)\}_{n \in \mathbb{N}}$ of closed subsets of X such that

- (1) $U = \bigcup_{n \in \mathbb{N}} F(n, U)$;
- (2) $F(n, U) \subset F(n, V)$ whenever $U \subset V$.

A correspondence $U \rightarrow \{F(n, U)\}_{n \in \mathbb{N}}$ is a *semi-stratification* for X whenever it satisfies the conditions (1) and (2).

This project was supported by NNSF of China (No.10571151).

A collection \mathcal{P} of pairs of subsets of a space X is called a *pair-network* for X if whenever $x \in U$ with U open in X , $x \in P_1 \subset P_2 \subset U$ for some $(P_1, P_2) \in \mathcal{P}$. A collection \mathcal{P} of pairs of subsets of a space X is called *cushioned* if $\overline{\bigcup\{P_1 : (P_1, P_2) \in \mathcal{P}'\}} \subset \bigcup\{P_2 : (P_1, P_2) \in \mathcal{P}'\}$ for each $\mathcal{P}' \subset \mathcal{P}$.

Lemma 2. *A space is a semi-stratifiable space if and only if it is a space with a σ -cushioned network.*

PROOF: Let (X, τ) be a semi-stratifiable space, and $U \rightarrow \{F(n, U)\}_{n \in \mathbb{N}}$ be a semi-stratification for X . For each $n \in \mathbb{N}$, put $\mathcal{P}_n = \{(F(n, U), U) : U \in \tau\}$. If $\tau' \subset \tau$, then $\bigcup_{U \in \tau'} F(n, U) \subset F(n, \bigcup \tau') \subset \bigcup \tau'$. Thus \mathcal{P}_n is a cushioned collection of X . Hence, $\bigcup_{n \in \mathbb{N}} \mathcal{P}_n$ is a σ -cushioned network for X .

Conversely, suppose that a space X has a σ -cushioned network $\bigcup_{n \in \mathbb{N}} \mathcal{P}_n$, where each \mathcal{P}_n is cushioned in X . For each $n \in \mathbb{N}$, $U \in \tau$, put $F(n, U) = \overline{\bigcup\{P_1 : (P_1, P_2) \in \bigcup_{i \leq n} \mathcal{P}_i, P_2 \subset U\}}$. It is obvious that $F(n, U) \subset U$ and $F(n, U) \subset F(n, V)$ whenever $U \subset V$. Let $x \in U \in \tau$. There exist $n \in \mathbb{N}$ and $(P_1, P_2) \in \mathcal{P}_n$ such that $x \in P_1 \subset P_2 \subset U$, thus $x \in F(n, U)$. Hence $U = \bigcup_{n \in \mathbb{N}} F(n, U)$. Therefore, $U \rightarrow \{F(n, U)\}_{n \in \mathbb{N}}$ is a semi-stratification for X . □

Definition 3. A collection \mathcal{P} of pairs of subsets of a space X is called a *(modk)-network* for X if, there is a cover \mathcal{K} of compact subsets of X such that, whenever $K \in \mathcal{K}$ and $K \subset U$ with U open in X , then $K \subset P_1 \subset P_2 \subset U$ for some $(P_1, P_2) \in \mathcal{P}$.

Theorem 4. *Spaces with a σ -cushioned (modk)-network are D-spaces.*

PROOF: Suppose a space X has a σ -cushioned (modk)-network $\bigcup_{n \in \mathbb{N}} \mathcal{P}_n$ with respect to a cover \mathcal{K} of compact subsets, where each \mathcal{P}_n is cushioned and $\mathcal{P}_n \subset \mathcal{P}_{n+1}$ for each $n \in \mathbb{N}$. Assume each \mathcal{P}_n is well-ordered.

Let ϕ be an arbitrary neighborhood assignment for X . First, a subset D_n of X is defined inductively as follows for each $n \in \mathbb{N}$.

Set $D_0 = \emptyset$. Assume $D_m \subset X$ is defined for all $0 < m < n$.

A finite subset D_α^n of X is defined inductively as follows for order numbers α .

Set $D_0^n = \emptyset$. Assume D_β^n is defined for all $0 < \beta < \alpha$. Put

$$U = \bigcup \left\{ \phi(d) : d \in \left(\bigcup \{D_\beta^n : \beta < \alpha\} \right) \cup D_{n-1} \right\}.$$

Denote R_α^n by the following requirement: There exist $K \in \mathcal{K}$, $(P_1, P_2) \in \mathcal{P}_n$ and $\{x_1, x_2, \dots, x_k\} \subset K \setminus U$ such that

$$K \subset P_1 \subset P_2 \subset U \cup \phi(x_1) \cup \phi(x_2) \cup \dots \cup \phi(x_k).$$

If R_α^n does not hold, the induction on α stops. Otherwise, let (P_1, P_2) be the first pair in \mathcal{P}_n that satisfies R_α^n , and put $D_\alpha^n = \{x_1, x_2, \dots, x_k\}$.

Let $D_n = (\bigcup_{\alpha < \gamma_n} D_\alpha^n) \cup D_{n-1}$ for some order number γ_n , where γ_n is the first ordinal number for which R_α^n does not hold.

Secondly, put $D = \bigcup \{D_n : n \in \mathbb{N}\}$, and we shall prove that X is a D-space by D .

Claim 1. $X = \bigcup_{d \in D} \phi(d)$.

If not, then $K \setminus \bigcup_{d \in D} \phi(d) \neq \emptyset$ for some $K \in \mathcal{K}$. Put $L = K \setminus \bigcup_{d \in D} \phi(d)$. L is a non-empty, compact subset of X . There is a finite subset $\{x_1, x_2, \dots, x_k\}$ of L such that $L \subset \phi(x_1) \cup \phi(x_2) \cup \dots \cup \phi(x_k)$ because ϕ is a neighborhood assignment for X . Denote $M = K \setminus (\phi(x_1) \cup \phi(x_2) \cup \dots \cup \phi(x_k))$. Then $M \subset K \setminus L \subset \bigcup_{d \in D} \phi(d)$. By the compactness of M , $M \subset \bigcup_{d \in D_j} \phi(d)$ for some $j \in \mathbb{N}$, thus $K \subset \phi(x_1) \cup \phi(x_2) \cup \dots \cup \phi(x_k) \cup (\bigcup_{d \in D_j} \phi(d))$. Since \mathcal{P} is a (modk)-network for X with respect to \mathcal{K} , there are $m \in \mathbb{N}$ and $(P_1, P_2) \in \mathcal{P}_m$ such that

$$K \subset P_1 \subset P_2 \subset \phi(x_1) \cup \phi(x_2) \cup \dots \cup \phi(x_k) \cup \left(\bigcup_{d \in D_j} \phi(d) \right).$$

Put $n = \max\{j + 1, m\}$, $U = \bigcup \{\phi(d) : d \in D_{n-1}\}$. Then $\bigcup_{d \in D_j} \phi(d) \subset U$, $(P_1, P_2) \in \mathcal{P}_n$ and $\{x_1, x_2, \dots, x_k\} \subset K \setminus U$. Hence R_1^n holds, thus $D_1^n = \{x_1, x_2, \dots, x_k\}$, and $K \subset P_1 \subset P_2 \subset \bigcup_{d \in D} \phi(d)$, a contradiction.

Claim 2. Each D_n is a closed discrete subset of X .

D_0 is closed discrete in X . Suppose that D_{n-1} is closed discrete in X . It suffices to prove that $\bigcup_{\alpha < \gamma_n} D_\alpha^n$ is closed discrete in X in order to show that D_n is closed discrete in X . For each $\alpha < \gamma_n$, let $(P_{1\alpha}^n, P_{2\alpha}^n)$ be the first element in \mathcal{P}_n satisfying R_α^n . Assume that $x \in \overline{\bigcup_{\alpha < \gamma_n} D_\alpha^n}$. Since $\bigcup_{\alpha < \gamma_n} D_\alpha^n \subset \bigcup_{\alpha < \gamma_n} P_{1\alpha}^n$, then $x \in \bigcup_{\alpha < \gamma_n} P_{2\alpha}^n \subset \bigcup \{\phi(d) : d \in (\bigcup \{D_\alpha^n : \alpha < \gamma_n\}) \cup D_{n-1}\}$. And since each $D_\alpha^n \cap \bigcup_{d \in D_{n-1}} \phi(d) = \emptyset$, then $x \in \bigcup \{\phi(d) : d \in \bigcup_{\alpha < \gamma_n} D_\alpha^n\}$. Let α_x be the minimal element α in γ_n satisfying $x \in \bigcup \{\phi(d) : d \in D_\alpha^n\}$. Put $V = \bigcup \{\phi(d) : d \in D_{\alpha_x}^n\} \setminus \overline{\bigcup_{\alpha < \alpha_x} P_{1\alpha}^n}$. Then V is an open neighborhood of x in X , and $V \cap (\bigcup_{\alpha < \gamma_n} D_\alpha^n) \subset D_{\alpha_x}^n$ is finite. Hence, $\bigcup_{\alpha < \gamma_n} D_\alpha^n$ is closed discrete in X .

Claim 3. D is closed discrete in X .

For each $x \in X$, there is $n \in \mathbb{N}$ such that $x \in \bigcup_{d \in D_n} \phi(d)$ by Claim 1. Since D_n is closed discrete in X , there is an open neighborhood W of x in X such that $W \subset \bigcup_{d \in D_n} \phi(d)$ and W contains at most a point in D_n . Thus $W \cap D \subset ((\bigcup_{d \in D_n} \phi(d)) \cap (D \setminus D_n)) \cup (W \cap D_n)$. For each $y \in D \setminus D_n$, $y \in D_\alpha^m$ for some $m > n$ and some $\alpha < \gamma_m$. Put $U = \bigcup \{\phi(d) : d \in (\bigcup \{D_\beta^m : \beta < \alpha\}) \cup D_{m-1}\}$. Then $\bigcup_{d \in D_n} \phi(d) \subset U$, and $D_\alpha^m \cap U = \emptyset$, thus $y \notin \bigcup_{d \in D_n} \phi(d)$. Hence, $(D \setminus D_n) \cap (\bigcup_{d \in D_n} \phi(d)) = \emptyset$. This shows that W contains at most a point in D . Therefore, D is closed discrete in X . \square

A collection \mathcal{F} of closed subsets of a space X is called *closure-preserved* if $\bigcup \mathcal{F}'$ is closed for each $\mathcal{F}' \subset \mathcal{F}$. A space is called a *strong Σ -space* (*strong $\Sigma^\#$ -space*) [8] if there exists a σ -locally finite (σ -closure-preserved) collection \mathcal{F} of closed subsets, and a cover \mathcal{K} of X by compact subsets, such that, whenever $K \in \mathcal{K}$ and $K \subset U$ with U open in X , then $K \subset F \subset U$ for some $F \in \mathcal{F}$. Every strong Σ -space is a strong $\Sigma^\#$ -space. If a collection \mathcal{F} of closed subsets of a space X is closure-preserved, then a collection $\{(F, F) : F \in \mathcal{F}\}$ of pairs of subsets of X is cushioned. Hence every strong $\Sigma^\#$ -space or semi-stratifiable space is a space with a σ -cushioned (mod k)-network. Thus Theorem 4 is a common generalization of Buzyakova's result [3] and Borges-Wehrly's result [2]. The following corollary is an unpublished result obtained by Liang-xue Peng.

Corollary 5. *Every strong $\Sigma^\#$ -space is a D-space.*

Acknowledgment. The author would like to thank Liang-xue Peng for useful information about D-spaces.

REFERENCES

- [1] Arhangel'skii A.V., Buzyakova R.Z., *Addition theorems and D-spaces*, Comment. Math. Univ. Carolin. **43** (2002), 653–663.
- [2] Borges C.R., Wehrly A.C., *A study of D-spaces*, Topology Proc. **16** (1991), 7–15.
- [3] Buzyakova R.Z., *On D-property of strong Σ -spaces*, Comment. Math. Univ. Carolin. **43** (2002), 493–495.
- [4] Creede G., *Concerning semi-stratifiable spaces*, Pacific J. Math. **32** (1970), 47–54.
- [5] van Douwen E.K., *Simultaneous extension of continuous functions*, Thesis, Free University, Amsterdam, 1975.
- [6] van Douwen E.K., Pfeffer W.F., *Some properties of the Sorgenfrey line and related spaces*, Pacific J. Math. **81** (1979), 371–377.
- [7] Engelking R., *General Topology* (Revised and completed edition), Heldermann Verlag, Berlin, 1989.
- [8] Gruenhage G., *Generalized metric spaces*, in Handbook of Set-Theoretic Topology, K. Kunen, J.E. Vaughan (Eds.), Elsevier Science Publishers B.V., Amsterdam, 1984, pp. 423–501.

DEPARTMENT OF MATHEMATICS, ZHANGZHOU NORMAL UNIVERSITY, FUJIAN 363000, P.R. CHINA

INSTITUTE OF MATHEMATICS, NINGDE TEACHERS' COLLEGE, FUJIAN 352100, P.R. CHINA
E-mail: linshou@public.ndptt.fj.cn

(Received October 10, 2005, revised January 13, 2006)