

Zuo Dong Yang; Junli Yuan

Non-existence result for quasi-linear elliptic equations with supercritical growth

Commentationes Mathematicae Universitatis Carolinae, Vol. 48 (2007), No. 3, 417--430

Persistent URL: <http://dml.cz/dmlcz/119669>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2007

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Non-existence result for quasi-linear elliptic equations with supercritical growth

ZUODONG YANG, JUNLI YUAN

Abstract. We obtain a non-existence result for a class of quasi-linear eigenvalue problems when a parameter is small. By using Pohozaev identity and some comparison arguments, non-existence theorems are established for quasi-linear eigenvalue problems under supercritical growth condition.

Keywords: quasi-linear elliptic equations, non-existence, large solution, small solution

Classification: 35J65, 35B25

1. Introduction

In this paper we are concerned with the non-existence of positive solutions of a class of quasi-linear eigenvalue problems

$$(1.1) \quad -\operatorname{div}(|\nabla u|^{p-2}\nabla u) = \lambda f(u(x)) \quad \text{in } \Omega,$$

$$(1.2) \quad u = 0 \quad \text{on } \partial\Omega,$$

where $f \in C^1(0, \infty) \cap C^0([0, \infty))$, $f(s) > 0$ for $s \geq 0$; $\lambda > 0$, $\Omega = B_1 = \{x \in \mathbb{R}^N : |x| < 1\}$ is the unit ball, and $1 < p < N$. By a positive solution u of (1.1)–(1.2) we mean that $u \in C_0^1(\Omega)$, $u > 0$ in Ω , and satisfies

$$\int_{\Omega} |\nabla u|^{p-2}\nabla u \nabla v = \lambda \int_{\Omega} f(u)v$$

for any $v \in C_0^\infty(\Omega)$. Thus, solutions are considered in a weak sense. By a small solution u_λ of (1.1)–(1.2) we mean that $\lim_{\lambda \rightarrow 0^+} \|u_\lambda\|_\infty = 0$. By a positive large solution $u_\lambda(r)$ of (1.1)–(1.2) we mean that $\lim_{\lambda \rightarrow 0^+} \|u_\lambda\|_\infty = \infty$.

Project supported by the National Natural Science Foundation of China (Grant No. 10571022).

Project supported by the Natural Science Foundation of the Jiangsu Higher Education Institutions of China (Grant No. 04KJB110062) and the Science Foundation of Nanjing Normal University (Grant No. 2003SXXXGQ2B37).

Equations of the above form are mathematical models occurring in studies of the p -Laplace equation, generalized reaction-diffusion theory, non-Newtonian fluid theory ([1], [2]), non-Newtonian filtration ([3]) and the turbulent flow of a gas in a porous medium ([4]). In the non-Newtonian fluid theory, the quantity p is a characteristic of the medium. Media with $p > 2$ are called dilatant fluids and those with $p < 2$ are called pseudo-plastics. If $p = 2$, they are Newtonian fluids.

For $p = 2$, the problem (1.1)–(1.2) has been studied by many authors, such as Ni and Serrin [5], Gelfand [6], Keller and Cohen [7], Amann [8], Crandall and Rabinowitz [9], Lions [10], Brezis and Nirenberg [11], to name just a few. For $p > 1$, the existence and uniqueness of the positive solutions of (1.1)–(1.2) have been studied by many authors, for example [12]–[17], [20]–[21] and the references therein. When f is strictly increasing on \mathbb{R}^+ , $f(0) = 0$, $\lim_{s \rightarrow 0^+} f(s)/s^{p-1} = 0$ and $f(s) \leq \alpha_1 + \alpha_2 s^\mu$, $0 < \mu < p - 1$, $\alpha_1, \alpha_2 > 0$, it was shown in [12] that there exist at least two positive solutions for equations (1.1)–(1.2) when λ is sufficiently large. If $\liminf_{s \rightarrow 0^+} f(s)/s^{p-1} > 0$, $f(0) = 0$ and the monotonicity hypothesis $(f(s)/s^{p-1})' < 0$ holds for all $s > 0$, it was proved in [13] that the problem (1.1)–(1.2) has a unique positive solution when λ is sufficiently large. Moreover, it was also shown in [14] that problem (1.1)–(1.2) has a unique positive large solution and at least one positive small solution when λ is large if f is nondecreasing, and there exist $\alpha_1, \alpha_2 > 0$ such that $f(s) \leq \alpha_1 + \alpha_2 s^\beta$, $0 < \beta < p - 1$; $\lim_{s \rightarrow 0^+} \frac{f(s)}{s^{p-1}} = 0$, and there exist $T, Y > 0$ with $Y \geq T$ such that

$$(f(s)/s^{p-1})' > 0 \quad \text{for } s \in (0, T)$$

and

$$(f(s)/s^{p-1})' < 0 \quad \text{for } s > Y.$$

In contrast to these cases, it seems that very little is known about existence and non-existence of positive solutions and non-small solutions for the problem (1.1)–(1.2) when λ is sufficiently small. Hai [18] considered the case when Ω is an annular domain, and obtained the existence of positive large solutions for the problem (1.1)–(1.2) when λ is sufficiently small. Guo and Yang [22] considered the case when Ω is a bounded smooth domain, and obtained the existence of positive large solutions and small solutions for the problem (1.1)–(1.2) when λ is sufficiently small. In this paper, we shall consider the case when $\Omega = B_1 = \{x \in \mathbb{R}^N : |x| < 1\}$ is the unit ball, and establish the non-existence of positive solutions and non-small solutions for the problem (1.1)–(1.2) when λ is sufficiently small.

Our approach depends heavily upon the special properties of the positive radial solutions for the problem (1.1)–(1.2). We expect that such non-existence result of (1.1)–(1.2) are still true for the general domain Ω .

We can find the related non-existence results for $p = 2$ in [19]. When $p = 2$, it is well known that all the positive solutions in $C^2(B_R)$ of the problem

$$\begin{aligned} \Delta u + f(u) &= 0 \text{ in } B_R, \\ u(x) &= 0 \text{ on } \partial B_R \end{aligned}$$

are radially symmetric solutions for very general f (see [25]). Unfortunately, this result does not apply to the case $p \neq 2$. Kichenassary and Smoller showed that there exist many positive nonradial solutions of the above problem for some f (see [26]). The major stumbling block in the case of $p \neq 2$ is that certain nice features inherent to the case $p = 2$ seem to be lost or at least difficult to verify. The main differences between $p = 2$ and $p \neq 2$ can be found in [12], [13].

2. Non-existence result

In this section we study the non-existence of positive solutions of the problems (1.1)–(1.2). The nonlinear function $f \in C^1(\mathbb{R})$ (or f is in general locally Lipschitz continuous) satisfies the supercritical condition as $u \rightarrow \infty$; that is, f satisfies the following conditions:

(H_1) When $p \geq 2$, there are $q > \frac{N(p-1)+p}{N-p}$, $A > 0$ such that $(q + 1)F(u) \leq uf(u)$ for $u \geq A$, where $F(u) = \int_0^u f(v) dv$ and A is a positive constant with $F(A) > 0$.

(H_1)' When $1 < p < 2$, there are $q + 1 > \frac{2^{(2-p)/(p-1)}Np}{N-p}$, $A > 0$ such that $(q + 1)F(u) \leq uf(u)$ for $u \geq A$, where $F(u) = \int_0^u f(v) dv$ and A is a positive constant with $F(A) > 0$.

To prove the main theorem, we consider the following initial value problems

$$(2.1) \quad (\Phi_p(u'))' + \frac{(N-1)}{r} \Phi_p(u') + f(u(r)) = 0, \quad r > 0,$$

$$(2.2) \quad u(0, \alpha) = \alpha > 0, \quad u'(0, \alpha) = 0,$$

where $\Phi_p(s) = |s|^{p-2}s$, $p > 1$.

We first recall a Pohozaev identity which was obtained by Ni and Serrin [5], or Mitidieri and Pohozaev [23].

Lemma 2.1. *Let $u(r)$ be a solution of equation (2.1) in $(r_1, r_2) \subset (0, \infty)$ and a be an arbitrary constant. Then, for each $r \in (r_1, r_2)$ we have*

$$\begin{aligned} (2.3) \quad \frac{d}{dr} [r^N \{ (1 - 1/p)|u'|^p + F(u) + \frac{a}{r}uu'|u'|^{p-2} \}] \\ = r^{N-1} [NF(u) - au f(u) + (a + 1 - N/p)|u'|^p], \end{aligned}$$

where $F(u) = \int_0^u f(s) ds$.

Definition 2.2. For each $\alpha \in (0, \infty)$ and $B \geq 0$, let $R(\alpha, B)$ be the first r such that $u(r, \alpha) = B$. If there is no such r , we shall adopt the convention that $R(\alpha, B) = \infty$. We also stipulate that $R(\alpha) = R(\alpha, 0)$ and $R_1(\alpha) = R(\alpha, A)$, where A is given in (H_1) or $(H_1)'$.

Definition 2.3. For $p \geq 2$, let $\gamma = \frac{1}{(q+1)(N-p)}[(N-p)(q+1) - Np] > 0$; for $1 < p < 2$, let $\gamma_1 = \frac{1}{(q+1)(N-p)}[(N-p)(q+1) - 2^{(2-p)/(p-1)}Np] > 0$. Define two positive functions $R_*(B)$ and $R^*(B)$ on $[A, \infty]$ by

$$R_*(B)^{p/(p-1)} = M(\overline{B})^{-1/(p-1)}B$$

and

$$R^*(B)^p = \left(\frac{p}{p-1}\right)^{p-1} \left(\frac{NB}{q+1}\right)^p (F(B))^{-1},$$

where $\overline{B} = [N^{-1/(p-1)}\frac{(p-1)}{p} + 1]\gamma^{-1}B$ for $p \geq 2$; $\overline{B} = [2^{\frac{2-p}{p-1}}N^{-\frac{1}{p-1}}\frac{(p-1)}{p} + 1]\gamma_1^{-1}B$ for $1 < p < 2$, and $M(\overline{B}) = \max\{f(u) : u \in [0, \overline{B}]\}$.

We shall first prove that for a fixed $B \geq A$, there exist an upper bound and a lower bound for $R(\alpha, B)$.

Lemma 2.4. Let f satisfy (H_1) for $p \geq 2$ or $(H_1)'$ for $1 < p < 2$. Then for any $B \geq A$ and $\alpha \in (\overline{B}, \infty)$, we have

$$(2.4) \quad R_*(B) \leq R(\alpha, B) \leq R^*(B),$$

and

$$(2.5) \quad \left(\frac{(q+1)F(B)}{N} \frac{F(B)}{B}\right)^{1/(p-1)} R_*(B)^{1/(p-1)} \leq -u'(R(\alpha, B), \alpha) \leq \frac{pN}{(p-1)(q+1)} BR_*(B)^{-1}.$$

PROOF: Letting $u(r) = u(r, \alpha)$ and $a = N/(q+1)$ in equation (2.3) and integrating equation (2.3) from 0 to r , from (H_1) or $(H_1)'$ we have

$$(2.6) \quad \frac{(p-1)}{p}|u'|^p + F(u(r, \alpha)) + \frac{N}{(q+1)} \frac{u(r, \alpha)u'(r, \alpha)|u'(r, \alpha)|^{p-2}}{r} < 0$$

if $u(s, \alpha) > A$ for all $s \in [0, r]$. It is clear that (H_1) or $(H_1)'$ implies $F(u) > 0$ for all $u > A$. Hence, for any $\alpha \in (A, \infty)$, by (2.6) we have $u'(r, \alpha) < 0$ in $(0, R_1(\alpha))$. Furthermore, we have $R_1(\alpha) < \infty$ for all $\alpha \in (A, \infty)$. Indeed, by (H_1) or $(H_1)'$ there is a positive constant m such that

$$(2.7) \quad f(u) \geq m \text{ for all } u \geq A.$$

From (2.1)–(2.2) and (2.7), for $r \in (0, R_1(\alpha))$ and $\alpha \geq A$, we have

$$(2.8) \quad r^{N-1} \Phi_p(u'(r, \alpha)) = - \int_0^r s^{N-1} f(u(s, \alpha)) ds \leq -\frac{m}{N} r^N,$$

which implies that

$$R_1(\alpha)^{p/(p-1)} \leq \left(\frac{N}{m}\right)^{1/(p-1)} \left[\frac{p}{(p-1)}(\alpha - A)\right].$$

Therefore, by (H_1) , $(H_1)'$ and (2.6) we obtain

$$(2.9) \quad \frac{(p-1)}{p} |u'(R(\alpha, B), \alpha)|^p < \frac{N}{(q+1)} \frac{B}{R(\alpha, B)} |u'(R(\alpha, B), \alpha)|^{p-1}$$

and

$$(2.10) \quad F(B) < \frac{N}{(q+1)} \frac{B}{R(\alpha, B)} |u'(R(\alpha, B), \alpha)|^{p-1}.$$

Now, (2.9) implies

$$(2.11) \quad (-u'(R(\alpha, B), \alpha))R(\alpha, B) < \frac{pN}{(p-1)(q+1)} B.$$

From (2.10) and (2.11), we obtain an upper bound for $R(\alpha, B)$, that is,

$$(2.12) \quad R(\alpha, B)^p \leq \left[\left(\frac{p}{p-1}\right)^{p-1} \left(\frac{NB}{q+1}\right)^p\right] F(B)^{-1}$$

for all $\alpha \in (B, \infty)$. This proves the second inequality of (2.4). To prove the first inequality of (2.4), there are two cases to be considered:

- (a) $R(\alpha, \overline{B}) \geq R_*(B)$,
- (b) $R(\alpha, \overline{B}) < R_*(B)$.

In case (a), since $R(\alpha, B) > R(\alpha, \overline{B})$ we have $R(\alpha, B) > R_*(B)$. In case (b), we need a comparison argument.

Let $v_\alpha(r) \equiv v(r, \alpha, \overline{B})$ be the solution of the initial value problem

$$(2.13) \quad (\Phi_p(v'))' + \frac{N-1}{r} \Phi_p(v') + \overline{C} = 0 \quad \text{for } r > R(\alpha, \overline{B}),$$

$$(2.14) \quad v(R(\alpha, \overline{B})) = \overline{B},$$

$$(2.15) \quad v'(R(\alpha, \overline{B})) = u'(R(\alpha, \overline{B}), \alpha),$$

where $\overline{C} = M(\overline{B})$.

Then $v_\alpha(r)$ can be solved explicitly as

$$(2.16) \quad v_\alpha(r) = \bar{B} - \int_{\bar{R}}^r \left[\left(\frac{\bar{R}}{s}\right)^{N-1} |u'(\bar{R})|^{p-1} + \frac{\bar{C}}{N} \left(s - \frac{\bar{R}^N}{s^{N-1}}\right) \right]^{1/(p-1)} ds,$$

where $\bar{R} = R(\alpha, \bar{B})$. We further consider two subcases here: (i) $p \geq 2$ and (ii) $1 < p < 2$.

In subcase (i), it is obvious that $1/(p-1) \leq 1$. Using the inequalities $(1+x)^{1/(p-1)} \leq 1+x^{1/(p-1)}$ for $x \geq 0$ and (2.11), we have

$$\begin{aligned} v_\alpha(r) &\geq \bar{B} - \int_{\bar{R}}^r \left[\left(\frac{\bar{R}}{s}\right)^{N-1} |u'(\bar{R})|^{p-1} + \frac{\bar{C}}{N} s \right]^{1/(p-1)} ds \\ &\geq \bar{B} - \int_{\bar{R}}^r \left(\frac{\bar{R}}{s}\right)^{(N-1)/(p-1)} |u'(\bar{R})| \left[1 + \frac{((\bar{C}/N)s)^{1/(p-1)}}{(\bar{R}/s)^{(N-1)/(p-1)} |u'(\bar{R})|} \right] ds \\ &= \bar{B} - \int_{\bar{R}}^r \left[\left(\frac{\bar{R}}{s}\right)^{(N-1)/(p-1)} |u'(\bar{R})| + \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} s^{1/(p-1)} \right] ds \\ &\geq \bar{B} - \frac{(p-1)}{(N-p)} \bar{R} |u'(\bar{R})| - \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} \int_{\bar{R}}^r s^{1/(p-1)} ds \\ &\geq \bar{B} - \frac{(p-1)}{(N-p)} \frac{Np}{(p-1)(q+1)} \bar{B} - \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} \frac{(p-1)}{p} r^{p/(p-1)} \\ &= \gamma \bar{B} - \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} \frac{(p-1)}{p} r^{p/(p-1)} \\ &\geq B \end{aligned}$$

for all $r \in [R(\alpha, \bar{B}), R_*(B)]$.

In subcase (ii), we have $1/(p-1) > 1$. Let $q+1 > 2^{(2-p)/(p-1)}(Np/(N-p))$. Using the inequalities $(1+x)^{1/(p-1)} \leq 2^{(2-p)/(p-1)}(1+x^{1/(p-1)})$ for $x \geq 0$ and (2.13), we have

$$\begin{aligned} v_\alpha(r) &\geq \bar{B} - \int_{\bar{R}}^r \left[\left(\frac{\bar{R}}{s}\right)^{N-1} |u'(\bar{R})|^{p-1} + \frac{\bar{C}}{N} s \right]^{1/(p-1)} ds \\ &\geq \bar{B} - \int_{\bar{R}}^r \left(\frac{\bar{R}}{s}\right)^{(N-1)/(p-1)} |u'(\bar{R})| 2^{(2-p)/(p-1)} \\ &\quad \left[1 + \frac{((\bar{C}/N)s)^{1/(p-1)}}{(\bar{R}/s)^{(N-1)/(p-1)} |u'(\bar{R})|} \right] ds \\ &= \bar{B} - \int_{\bar{R}}^r 2^{(2-p)/(p-1)} \left[\left(\frac{\bar{R}}{s}\right)^{(N-1)/(p-1)} |u'(\bar{R})| + \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} s^{1/(p-1)} \right] ds \\ &\geq \bar{B} - 2^{(2-p)/(p-1)} \frac{(p-1)}{(N-p)} \bar{R} |u'(\bar{R})| - 2^{(2-p)/(p-1)} \end{aligned}$$

$$\begin{aligned}
 & \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} \int_{\bar{R}}^r s^{1/(p-1)} ds \\
 \geq & \bar{B} - 2^{(2-p)/(p-1)} \frac{(p-1)}{(N-p)} \frac{Np}{(p-1)(q+1)} \bar{B} - 2^{(2-p)/(p-1)} \\
 & \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} \frac{(p-1)}{p} r^{p/(p-1)} \\
 = & \gamma_1 \bar{B} - 2^{(2-p)/(p-1)} \left(\frac{\bar{C}}{N}\right)^{1/(p-1)} \frac{(p-1)}{p} r^{p/(p-1)} \\
 \geq & B
 \end{aligned}$$

for all $r \in [R(\alpha, \bar{B}), R_*(B)]$. Therefore, (2.4) follows if we can prove that $u(r, \alpha) \geq v_\alpha(r)$ on $[R(\alpha, \bar{B}), R_*(B)]$.

In fact, we have

$$(2.17) \quad (r^{N-1}\Phi_p(u'))' - (r^{N-1}\Phi_p(v'_\alpha))' = r^{N-1}\{\bar{C} - f(u(r, \alpha))\} \geq 0$$

as long as $u(r, \alpha) > 0$. That is,

$$(2.18) \quad (p-1)(r^{N-1}|\xi(r)|^{p-2}(u-v_\alpha)')' \geq 0$$

as long as $u(r, \alpha) > 0$. Here $\xi(r)$ is between $u'(r)$ and $v'_\alpha(r)$. Integrating (2.18) twice and using (2.14)–(2.15), we obtain $u(r, \alpha) \geq v_\alpha(r)$ on $[R(\alpha, \bar{B}), R_*(B)]$. This proves the first inequality of (2.4).

Finally, (2.5) follows from (2.4), (2.10) and (2.11). The proof is complete. \square

Remark 2.5. If the growth of f is critical, then $R(\alpha)$ may tend to 0 as $\alpha \rightarrow \infty$. Indeed, let us consider

$$f(u) = \begin{cases} \frac{N(N-p)^{p-1}}{p-1} \varepsilon^{p(2-p)} u^{(N(p-1)+p)/(N-p)} & \text{if } u \geq 1, p \geq 2 \\ \frac{N(N-p)^{p-1}}{p-1} \varepsilon^{p(2-p)} u^{(N(p2^{(2-p)/(p-1)}-1)+p)/(N-p)} & \text{if } u \geq 1, 1 < p < 2 \\ \frac{N(N-p)^{p-1}}{p-1} & \text{if } u \leq 1. \end{cases}$$

Then it is well known for any $\varepsilon \in (0, 1)$ that

$$U_\varepsilon(r) = \left(\frac{\varepsilon}{\varepsilon^2 + r^{p/(p-1)}}\right)^{(N-p)/p}$$

is a solution of (2.3)–(2.4) for $U_\varepsilon(r) > 1, p \geq 2$. Note that $U_\varepsilon(0) = \varepsilon^{-(N-p)/p} \equiv \alpha$ which tends to ∞ as $\varepsilon \rightarrow 0^+$. Let $A = 1$ in (H_1) . Then it is easy to verify that

$$R_1(\alpha)^{p/(p-1)} = \varepsilon - \varepsilon^2$$

and

$$-u'(R_1(\alpha), \alpha) = \frac{N-p}{p-1}(\varepsilon - \varepsilon^2)^{1/p}\varepsilon^{-1},$$

and so

$$\lim_{\varepsilon \rightarrow 0^+} -u'(R_1(\alpha), \alpha)R_1(\alpha) = \frac{N-p}{p-1}$$

which is the contrary of (2.11). Using (2.16), it is easy to see that $R(\alpha)$ behaves like $\alpha^{-\frac{1}{(p-1)N}}$, which tends to 0 as $\alpha \rightarrow +\infty$.

Lemma 2.6 ([22]). *Let f be nondecreasing for $0 < s < 1$, and f satisfies*

- (i) $f \in C^1(0, \infty) \cup C^0([0, \infty))$;
- (ii) $f(s) > 0$ for $s \geq 0$ and $|f'(s)|$ is bounded in $[0, 1]$;
- (iii) there exists $\mu > p - 1$ such that

$$s^{-\mu}f(s) \rightarrow \beta \text{ as } s \rightarrow \infty;$$

- (iv) $\limsup_{s \rightarrow 0^+} (f(s)/s^{p-1})' < 0$.

Then problem (1.1)–(1.2) has only one positive small solution for λ sufficiently small.

Lemma 2.7 (Weak comparison principle) [20], [21]. *Let Ω be a bounded domain in \mathbb{R}^N ($N \geq 2$) with smooth boundary $\partial\Omega$ and $\varphi : (0, \infty) \rightarrow (0, \infty)$ is continuous and non-decreasing. Let $u_1, u_2 \in W^{1,p}(\Omega)$ satisfy*

$$\int_{\Omega} |\nabla u_1|^{p-2} \nabla u_1 \nabla \psi \, dx + \int_{\Omega} \varphi u_1 \psi \, dx \leq \int_{\Omega} |\nabla u_2|^{p-2} \nabla u_2 \nabla \psi \, dx + \int_{\Omega} \varphi u_2 \psi \, dx$$

for all non-negative $\psi \in W^{1,p}(\Omega)$. Then the inequality

$$u_1 \leq u_2 \text{ on } \partial\Omega$$

implies that

$$u_1 \leq u_2 \text{ in } \Omega.$$

Lemma 2.8. *Assume that f satisfies (H_1) for $p \geq 2$ or $(H_1)'$ for $1 < p < 2$, and*

- (H_2) $f(u) > 0$ for $u > 0$;
- (H_3) (i) $f(0) > 0$;
- (ii) $f(0) = 0$ and $\lim_{s \rightarrow 0^+} f(s)/s^{p-1} > 0$.

Then $R(\alpha) < \infty$, for all $\alpha > 0$.

PROOF: The hypothesis of the Theorem implies there is an $\epsilon > 0$ such that

$$(2.19) \quad f(u) \geq \epsilon u^{p-1} \text{ for all } u \geq 0.$$

It is easy to see that $R(\alpha) < \infty$ for all $\alpha > 0$. In fact, consider the problem

$$\begin{aligned} \operatorname{div} (|\nabla u|^{p-2} \nabla u) + f(u) &= 0 \quad \text{in } B_R, \\ u &= 0 \quad \text{on } \partial B_R. \end{aligned}$$

Let $R = R(\alpha)$, consider the transformation $r = Rs$ and denote $v(s, \alpha) = u(r, \alpha)$. Then v satisfies the problem

$$(2.20) \quad \operatorname{div} (|\nabla v|^{p-2} \nabla v) + R^p f(v) = 0 \quad \text{in } B_1,$$

$$(2.21) \quad v = 0 \quad \text{on } \partial B_1.$$

Suppose that there exists a sequence $\{(R_n, v_n)\}$ (where $R_n = R(\alpha_n)$, $v_n(s) = v(s, \alpha_n)$) satisfying $R_n \rightarrow \infty$ as $n \rightarrow \infty$ and v_n is a positive solution of (2.20)–(2.21) for $R = R_n$. Then, $\omega_n(s) = v_n / \|v_n\|_\infty$ solves the problem

$$\begin{aligned} -\operatorname{div} (|\nabla \omega_n|^{p-2} \nabla \omega_n) &= R_n^p \frac{f(v_n)}{\|v_n\|_\infty^{p-1}} \quad \text{in } B_1, \\ \omega_n(s) &= 0 \quad \text{on } \partial B_1. \end{aligned}$$

It follows from the above problem that

$$\omega_n(s) = R_n^{p/(p-1)} G_p^1 \left(\frac{f(v_n)}{\|v_n\|_\infty^{p-1}} \right),$$

where G_p^1 is the inverse of $A_p^1 = -\operatorname{div} (|\nabla \cdot|^{p-2} \nabla \cdot)$ under the Dirichlet boundary condition. By Lemma 2.7 and (2.19) imply that

$$(2.22) \quad \omega_n(s) \geq (\epsilon R_n^p)^{1/(p-1)} G_p^1(\omega_n^{p-1}) = (\epsilon R_n^p)^{1/(p-1)} \eta_n(s).$$

Here η_n satisfies

$$\begin{aligned} -\operatorname{div} (|\nabla \eta_n|^{p-2} \nabla \eta_n) &= \omega_n^{p-1} \quad \text{in } B_1, \\ \eta_n &= 0 \quad \text{on } \partial B_1. \end{aligned}$$

Since $\omega_n > 0$ and $\|\omega_n\|_\infty = 1$ for any n , the compactness of G_p^1 from $C^0(B_1)$ to $C^1(\overline{B_1})$ implies that there exists a subsequence of $\{\eta_n(s)\}$ (still denoted by $\{\eta_n(s)\}$ later) such that $\eta_n \rightarrow \eta$ in $C^1(\overline{B_1})$ as $n \rightarrow \infty$ and $\eta(s) > 0$ in B_1 . Now we easily obtain a contradiction from (2.22) since $R_n \rightarrow \infty$ as $n \rightarrow \infty$. The proof is complete. \square

Theorem 2.9. *Assume that f satisfies (H_1) for $p \geq 2$ or $(H_1)'$ for $1 < p < 2$. If $f(s) > 0$ for $s \geq 0$, then there exists $\lambda_* > 0$ such that there is no positive non-small radially symmetric solution of equations (1.1)–(1.2) for any $\lambda \in (0, \lambda_*)$. If $f(0) \leq 0$, then there exists $\lambda_* > 0$ such that there is no positive radially symmetric solution of the problem (1.1)–(1.2) for any $\lambda \in (0, \lambda_*)$.*

PROOF: It is easy to see that $(u(\cdot), \lambda)$ is a positive radial solution of equations (1.1)–(1.2) if and only if $u(\cdot, \alpha)$ is a positive solution of equations (2.1)–(2.2) with $u(r) = u(\lambda^{1/p}r, \alpha)$ and $\lambda = R^p(\alpha)$, where $R(\alpha)$ is the first zero of $u(\cdot, \alpha)$. By Lemma 2.8, we have $R(\alpha) < \infty$ for all $\alpha > 0$. Therefore the solution set of (2.1)–(2.2) can be written as $\{(u(\cdot, \alpha), \lambda(\alpha)) : \alpha \in (0, \infty)\}$ with $\lambda(\alpha) = R^p(\alpha)$. Therefore, it is sufficient to study $R(\alpha)$ for $\alpha \in (0, \infty)$.

It is clear that $R(\alpha) > 0$ for $\forall \alpha \in (0, \infty)$. It is also easy to see that $\alpha_k \rightarrow \alpha_0 \in (0, \infty)$ and then $R(\alpha_0) > 0$. Hence, by Lemma 2.4, the only possibility for the case where $R(\alpha)$ tends to 0 as $\alpha \rightarrow 0^+$. We shall rule out this possibility by considering the following cases: (i) $f(0) = 0, \lim_{s \rightarrow 0^+} f(s)/s^{p-1} > 0$; (ii) $f(0) = 0, \lim_{s \rightarrow 0^+} f(s)/s^{p-1} = 0$; (iii) $f(0) = 0$ and $\lim_{s \rightarrow 0^+} f(s)/s^{p-1} < 0$ and (iv) $f(0) < 0$. For the case where $f(0) > 0$ and f is nondecreasing for $0 < s < 1$, we know from Lemma 2.6 that there exists a unique positive small solution $u(r, \lambda)$ which will tend to zero uniformly in Ω as $\lambda \rightarrow 0^+$. This implies that $u(\cdot, \alpha)$ is a positive small solution if $R(\alpha)$ is sufficiently small.

Case (i). In this case, we shall prove that problem (1.1)–(1.2) has no positive radially symmetric solution u_λ with $\|u_\lambda\|_\infty \rightarrow 0$ when λ is sufficiently small.

If $\lim_{s \rightarrow 0^+} f(s)/s^{p-1} = \alpha > 0$, suppose that there exists a sequence $\{(\lambda_n, u_n)\}$ satisfying $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$ and u_n is a radially symmetric positive solution of equations (1.1)–(1.2) for $\lambda = \lambda_n$ such that $\|u_n\|_\infty \rightarrow 0$ as $n \rightarrow \infty$. Then, $\omega_n(x) = u_n/\|u_n\|_\infty$ satisfies

$$(2.23) \quad -\operatorname{div}(|\nabla \omega_n|^{p-2} \nabla \omega_n) = \lambda_n \frac{f(\|u_n\|_\infty \omega_n)}{\|u_n\|_\infty^{p-1}} \omega_n^{p-1} \text{ in } B_1,$$

$$(2.24) \quad \omega_n(x) = 0 \text{ on } \partial B_1.$$

Since $\omega_n > 0, \|\omega_n\|_\infty = 1$ for any n and $\frac{f(\|u_n\|_\infty \omega_n)}{(\|u_n\|_\infty \omega_n)^{p-1}} \rightarrow \alpha$ as $n \rightarrow \infty$, the compactness of G_p^1 from $C^0(B_1)$ to $C_0^1(\overline{B_1})$ (see [12]) implies that there exists a subsequence of $\{\omega_n\}$ (still denoted by $\{\omega_n\}$ later) and $\bar{\omega} \in C_0^1(\overline{B_1})$ such that $\omega_n \rightarrow \bar{\omega}$ in $C^1(\overline{B_1})$. Thus, $\bar{\omega}$ is a bounded solution of

$$\begin{aligned} -\operatorname{div}(|\nabla \bar{\omega}|^{p-2} \nabla \bar{\omega}) &= 0 \text{ in } B_1, \\ \bar{\omega} &= 0 \text{ on } \partial B_1. \end{aligned}$$

This implies that $\bar{\omega} \equiv 0$ in B_1 . This contradicts the facts that $\omega_n \rightarrow \bar{\omega}$ in $C^1(\overline{B_1})$ and $\|\omega_n\|_\infty = 1$.

If $\lim_{s \rightarrow 0^+} f(s)/s^{p-1} = +\infty$, suppose that there exists a sequence $\{(\lambda_n, u_n)\}$ satisfying $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$ and u_n is a radial positive solution of equations (1.1)–(1.2) for $\lambda = \lambda_n$ such that $\|u_n\|_\infty \rightarrow 0$ as $n \rightarrow \infty$. Then $\omega_n(x) = u_n/\|u_n\|_\infty$ satisfies

$$(2.25) \quad \begin{aligned} -(r^{N-1}\Phi_p(\omega'_n))' &= \lambda_n r^{N-1} \|u_n\|_\infty^{(p-1)} f(\|u_n\|_\infty \omega_n) \text{ in } (0, 1), \\ \omega'_n(0) &= 0, \omega_n(1) = 0 \end{aligned}$$

and $\omega_n(0) = 1$. First, we shall prove that $\tau_n = \lambda_n \|u_n\|_\infty^{(p-1)}$ is uniformly bounded. Suppose that $\tau_n \rightarrow \infty$ as $n \rightarrow \infty$. Let $y_n = \tau_n^{1/p} r, \tilde{\omega}_n(y_n) = \omega_n(r)$. Then $\tilde{\omega}_n$ satisfies

$$\begin{aligned} -\operatorname{div}(|\nabla \tilde{\omega}_n|^{p-2} \nabla \tilde{\omega}_n) &= f(\|u_n\|_\infty \tilde{\omega}_n) \text{ in } B_n, \\ \tilde{\omega}_n &= 0 \text{ on } \partial B_n. \end{aligned}$$

Here B_n is B_1 under the change of variables. Since $\|u_n\|_\infty \rightarrow 0$ as $n \rightarrow \infty$ and $f(0) = 0$, we have that $\tilde{\omega}_n \rightarrow \tilde{\omega}$ in $C^1_{\text{loc}}(0, \infty)$ as $n \rightarrow \infty$ and $\tilde{\omega}(r)$ is a bounded solution of

$$-\operatorname{div}(|\nabla \tilde{\omega}|^{p-2} \nabla \tilde{\omega}) = 0 \text{ in } \mathbb{R}^N$$

with $\|\tilde{\omega}\|_\infty = 1$. This implies that $\tilde{\omega} \equiv 0$ in \mathbb{R}^N . This contradicts the fact that $\|\tilde{\omega}\|_\infty = 1$. Thus, $\{\tau_n\}$ is uniformly bounded. Then, equation (2.25) and $\|u_n\|_\infty = 1$ imply that there exists a subsequence of $\{\omega_n\}$ and $\omega \in C^1_0(\overline{B_1})$ such that $\omega_n \rightarrow \omega$ in $C^1(\overline{B_1})$. Then ω is a bounded solution of the problem

$$\begin{aligned} -\operatorname{div}(|\nabla \omega|^{p-2} \nabla \omega) &= 0 \text{ in } B_1, \\ \omega &= 0 \text{ on } \partial B_1 \end{aligned}$$

with $\|\omega\|_\infty = 1$. This implies that $\omega \equiv 0$. This contradicts the fact that $\|\omega\|_\infty = 1$.

Case (ii). In this case, we shall prove that $\lim_{\alpha \rightarrow 0^+} R(\alpha) = \infty$. We observe that $u(\cdot, \alpha)$ satisfies the following equation:

$$(2.26) \quad u(r, \alpha) = \alpha - \int_0^r \left(\int_0^s \left(\frac{z}{s}\right)^{N-1} f(u(z)) dz \right)^{1/(p-1)} ds.$$

Since $f(0) = 0, \lim_{s \rightarrow 0^+} f(s)/s^{p-1} = 0$, for any $\epsilon > 0$ there exists $\delta > 0$ such that $f(u) \leq \epsilon u^{p-1}$ for $u \in (0, \delta)$. Therefore, if $u(r, \alpha) \in (0, 2\alpha) \subset (0, \delta)$ then

$|f(u(r, \alpha))| \leq 2^{p-1}\epsilon\alpha^{p-1}$. Now, it is easy to verify that

$$\begin{aligned}
 & \left| \int_0^r \left(\int_0^s \left(\frac{z}{s}\right)^{N-1} f(u(z, \alpha)) dz \right)^{1/(p-1)} ds \right| \\
 & \leq \int_0^r \left(\int_0^s \left(\frac{z}{s}\right)^{N-1} |f(u(z, \alpha))| dz \right)^{1/(p-1)} ds \\
 (2.27) \quad & \leq 2\alpha\epsilon^{1/(p-1)} \left(\int_0^r s^{(1-N)/(p-1)} \left(\int_0^s z^{N-1} dz \right)^{1/(p-1)} ds \right) \\
 & = 2\alpha\epsilon^{1/(p-1)} \left(\frac{1}{N}\right)^{1/(p-1)} \left(\int_0^r s^{1/(p-1)} ds \right) \\
 & = \left(\frac{1}{N}\right)^{1/(p-1)} 2\alpha\epsilon^{1/(p-1)} \frac{(p-1)}{p} r^{p/(p-1)}
 \end{aligned}$$

as far as $u(s, \alpha) \in (0, 2\alpha)$ for all $s \in (0, r)$. Hence, by (2.26)–(2.27), and for $\alpha \in (0, \delta/2)$ and $r \in (0, (\frac{p}{2(p-1)})^{(p-1)/p} (N/\epsilon)^{1/p})$, we have

$$|u(r, \alpha)| \leq \alpha + \left| \int_0^r \left(\int_0^s \left(\frac{z}{s}\right)^{N-1} f(u(z)) dz \right)^{1/(p-1)} ds \right| \leq 2\alpha,$$

so $u(r, \alpha) \in (0, 2\alpha)$. This implies $\lim_{\alpha \rightarrow 0^+} R(\alpha) = \infty$.

Case (iii). In this case, there are positive constants m and δ such that $-mu^{p-1} \leq f(u) \leq 0$ on $[0, \delta]$. Therefore, if $u(s, \alpha) \in [0, \delta]$ for all $s \in (0, r)$, then by (2.26) we have

$$\begin{aligned}
 (2.28) \quad u(r, \alpha) & \leq \alpha + m^{1/(p-1)} \int_0^r \left(\int_0^s \left(\frac{z}{s}\right)^{N-1} u^{p-1}(z, \alpha) dz \right)^{1/(p-1)} ds \\
 & \leq \alpha + m^{1/(p-1)} u(r, \alpha) \frac{(p-1)}{p} \left(\frac{1}{N}\right)^{1/(p-1)} r^{p/(p-1)}.
 \end{aligned}$$

Hence, if $u(R(\alpha, \delta), \alpha) = \delta$, then (2.26) implies that

$$R^{p/(p-1)}(\alpha, \delta) \geq \frac{p(\delta-\alpha)N^{1/(p-1)}}{\delta(p-1)m^{1/(p-1)}} \text{ and so } R(\alpha) \text{ has a positive lower bound for } \alpha \in (0, \delta/2).$$

Case (iv). In this case, there are $\epsilon > 0$ and $\delta > 0$ such that $f(u) \leq -\epsilon$ on $[0, \delta]$. Let $\overline{C} = -\epsilon$ in (2.13), $R(\alpha, \overline{B}) = 0$, $\overline{B} = \alpha$ in (2.14), and $u'(0, \alpha) = 0$ in (2.15). Then (2.18) becomes $v_\alpha(r) = \alpha + (\frac{\epsilon}{N})^{1/(p-1)} (\frac{p-1}{p}) r^{p/(p-1)}$ which implies that

$$(2.29) \quad u(r, \alpha) \geq v_\alpha(r) = \alpha + \left(\frac{\epsilon}{N}\right)^{1/(p-1)} \left(\frac{p-1}{p}\right) r^{p/(p-1)}$$

as long as $u(r, \alpha) \in [0, \delta]$. In particular, $R(0) > 0$. The continuous dependence of $u(\cdot, \alpha)$ in α and (2.29) imply that there is a positive lower bound for $R(\alpha)$ for all $\alpha \in [0, \delta]$. The proof of Theorem 2.9 is complete. \square

Remark 2.10. It is worth remarking that the validity of Theorem 2.9 relies on the topology of the domain Ω . Indeed when Ω is an annular domain, i.e., $\Omega = \{x \in \mathbb{R}^N : a < |x| < b\}$, $N \geq 2$, and $f(u)$ is continuous and $\lim_{u \rightarrow \infty} \frac{f(u)}{|u|^{p-2}u} = \infty$ (f is superlinear) uniformly for $t \in [a, b]$, there is at least one positive non-small solution for each $\lambda \in (0, \lambda^*)$, see [18], [24] and the reference therein.

REFERENCES

- [1] Astarita G., Marrucci G., *Principles of Non-Newtonian Fluid Mechanics*, McGraw-Hill, 1974.
- [2] Martinson L.K., Pavlov K.B., *Unsteady shear flows of a conducting fluid with a rheological power law*, *Magnitnaya Gidrodinamika* **2** (1971), 50–58.
- [3] Kalashnikov A.S., *On a nonlinear equation arising in the theory of nonlinear filtration* (in Russian), *Trudy Sem. Petrovsk.* **4** (1978), 137–146.
- [4] Esteban J.R., Vazquez J.L., *On the equation of turbulent filtration in one-dimensional porous media*, *Nonlinear Anal.* **10** (1982), 1303–1325.
- [5] Ni W.M., Serrin J., *Nonexistence theorems for singular solutions of quasilinear partial differential equations*, *Comm. Pure Appl. Math.* **39** (1986), 379–399.
- [6] Gelfand I.M., *Some problems in the theory of quasilinear equations*, *Amer. Math. Soc. Transl. Ser. 2* **29** (1963), 295–381.
- [7] Keller H.B., Cohen D.S., *Some positive problems suggested by nonlinear heat generation*, *J. Math. Mech.* **16** (1967), 1361–1376.
- [8] Amann H., *Fixed point equations and nonlinear eigenvalue problems in ordered Banach space*, *SIAM Rev.* **18** (1976), 620–709.
- [9] Crandall M.G., Rabinowitz P.H., *Some continuation and variational methods for positive solutions of nonlinear elliptic eigenvalue problems*, *Arch. Rational Mech. Anal.* **58** (1975), 207–218.
- [10] Lions P.L., *On the existence of positive solutions of semilinear elliptic equations*, *SIAM Rev.* **24** (1982), 441–467.
- [11] Brezis H., Nirenberg L., *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, *Comm. Pure Appl. Math.* **36** (1983), 437–477.
- [12] Guo Z., *Some existence and multiplicity results for a class of quasilinear elliptic eigenvalue problems*, *Nonlinear Anal.* **18** (1992), 957–971.
- [13] Guo Z., Webb J.R.L., *Uniqueness of positive solutions for quasilinear elliptic equations when a parameter is large*, *Proc. Royal Soc. Edinburgh, Sect. A* **124** (1994), 189–198.
- [14] Guo Z., *On the number of positive solutions for quasilinear elliptic eigenvalue problems*, *Nonlinear Anal.* **27** (1996), no. 2, 229–247.
- [15] Guo Z., *Existence and uniqueness of positive radial solutions for a class of quasilinear elliptic equations*, *Appl. Anal.* **47** (1992), 173–190.
- [16] Yang Z., Yang H., *A priori estimates for a quasilinear elliptic partial differential equations non-positive problems*, *Nonlinear Anal.* **43** (2001), no. 2, 173–181.
- [17] Yang Z., *Existence of entire explosive positive radial solutions for a class of quasilinear elliptic systems*, *J. Math. Anal. Appl.* **288** (2003), 768–783.
- [18] Hai D.D., *Positive solutions of quasilinear boundary value problems*, *J. Math. Anal. Appl.* **217** (1998), 672–686.
- [19] Lin S.S., *Positive singular solution for semilinear elliptic equations with supercritical growth*, *J. Differential Equations* **114** (1994), 57–76.
- [20] Yang Z., *Existence of positive entire solutions for singular and non-singular quasi-linear elliptic equation*, *J. Comput. Appl. Math.* **197** (2006), 355–364.

- [21] Lu Q., Yang Z., Twizell E.H., *Existence of entire explosive positive solutions of quasilinear elliptic equations*, Appl. Math. Comput. **148** (2004), 359–372.
- [22] Guo Z., Yang Z., *The structure of positive solutions to a class of quasilinear elliptic equations with a small parameter* (in Chinese), Chinese Ann. Math. Ser. A **19** (1998), 385–392.
- [23] Mitidieri E., Pohozaev S.I., *Nonexistence of positive solutions for quasilinear elliptic problem on \mathbf{R}^N* , Proc. Steklov Inst. Math. **227** (1999) 186–216).
- [24] Manásevich R., Schmitt K., *Boundary value problems for quasilinear second order differential equations*, Nonlinear Analysis and Boundary Value Problems (Udine), CISM Courses and Lectures, 371, Springer, Vienna, 1996, pp. 79–119.
- [25] Gidas B., Ni W.M., Nirenberg L., *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), 209–243.
- [26] Tolksdorf P., *On the Dirichlet problem for quasilinear equations in domains with conical boundary point*, Comm. Partial Differential Equations **8** (1983), no. 7, 773–817.

Zuodong Yang:

INSTITUTE OF MATHEMATICS, SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE, NANJING NORMAL UNIVERSITY, JIANGSU NANJING 210097, CHINA

and

COLLEGE OF ZHONGBEI, NANJING NORMAL UNIVERSITY, JIANGSU NANJING 210046, CHINA

Junli Yuan:

INSTITUTE OF MATHEMATICS, SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE, NANJING NORMAL UNIVERSITY, JIANGSU NANJING 210097, CHINA

(Received March 27, 2006, revised March 11, 2007)