

Jaroslav Ježek; Václav Slavík
Incomparably continuable sets of semilattices

Mathematica Bohemica, Vol. 125 (2000), No. 2, 135–137

Persistent URL: <http://dml.cz/dmlcz/125962>

Terms of use:

© Institute of Mathematics AS CR, 2000

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

INCOMPARABLY CONTINUABLE SETS OF SEMILATTICES

JAROSLAV JEŽEK, VÁCLAV SLAVÍK, Praha

(Received January 6, 1998)

Abstract. A finite set of finite semilattices is said to be incomparably continuable if it can be extended to an infinite set of pairwise incomparable (with respect to embeddability) finite semilattices. After giving some simple examples we show that the set consisting of the four-element Boolean algebra and the four-element fork is incomparably continuable.

Keywords: semilattice, embedding

MSC 1991: 06A12

By a semilattice we mean a meet semilattice. Two semilattices A and B are said to be comparable if either A can be embedded into B or B can be embedded into A . A finite set S of pairwise incomparable finite semilattices is called incomparably continuable if there exists an infinite set of pairwise incomparable finite semilattices containing S .

This paper does not attempt to develop a general theory of incomparably continuable sets. We just bring a few examples, showing that the general theory might be quite intricate. Of course, the concept could be introduced in the more general framework of universal algebra.

For the terminology and basic concepts of universal algebra, the reader is referred to [1].

For every $n \geq 1$ denote by \mathcal{C}_n the n -element chain. Denote by \mathcal{B}_4 the four-element Boolean algebra and by \mathcal{F} the fork, i.e., the semilattice with elements a, b, c, d and relations $a < b < c$ and $b < d$.

While working on this paper both authors were partially supported by the Grant Agency of the Czech Republic, Grant No 201/96/0312.

Theorem 1. *If $n \leq 3$, then $\{\mathcal{C}_n\}$ is not incomparably continuable. If $n \geq 4$, then $\{\mathcal{C}_n\}$ is incomparably continuable.*

Proof. Every semilattice incomparable with \mathcal{C}_3 consists of the zero element and a set of atoms. Any two such semilattices are comparable. Consequently, $\{\mathcal{C}_3\}$ is not incomparably continuable.

For every $n \geq 3$ let S_n be the semilattice with elements $0, a_1, \dots, a_n, b_1, \dots, b_n$ such that $0 < a_i < b_i$ for all i , $a_i < b_{i+1}$ for $i < n$, $a_n < b_1$ (and there are no other relations $x < y$). Then $\{\mathcal{C}_4, S_3, S_4, S_5, \dots\}$ is an infinite set of pairwise incomparable finite semilattices. \square

Lemma 2. *Let f_0, f_1, \dots be an infinite sequence of non-increasing functions from n^ω (where $n \in \omega$ is fixed). Then there exists an increasing sequence $i_0 < i_1 < i_2 < \dots$ of nonnegative integers with $f_{i_0} \leq f_{i_1} \leq \dots$*

Proof. By induction on n . For $n = 0$ it is clear. Let $n > 0$.

Suppose first that for some $k \geq 0$, there are infinitely many numbers i with $f_i(j) < n - 1$ for all $j \geq k$. Then for some $k \geq 0$, there are infinitely many numbers i with $f_i(j) = n - 1$ for $j < k$ and $f_i(j) < n - 1$ for $j \geq k$. Evidently, we can use induction.

Now suppose that there is no such k . Then there is an infinite increasing sequence $j_0 < j_1 < j_2 < \dots$ such that whenever $f_{j_i}(i) = n - 1$ then $f_{j_{i+1}}(i) = n - 1$. We can assume $j_i = i$, and hence: whenever $f_i(i) = n - 1$ then $f_{i+1}(i) = f_{i+2}(i) = \dots = n - 1$. Define g_k by $g_k(i) = f_k(i)$ if $f_k(i) < n - 1$ and $g_k(i) = n - 2$ if $f_k(i) = n - 1$. By induction, there is an infinite sequence $k_0 < k_1 < k_2 < \dots$ with $g_{k_0} \leq g_{k_1} \leq g_{k_2} \leq \dots$. But then evidently $f_{k_0} \leq f_{k_1} \leq f_{k_2} \leq \dots$ \square

Lemma 3. *Let a_0, a_1, \dots be an infinite sequence of nonnegative integers. Then there is a sequence $i_0 < i_1 < i_2 < \dots$ such that $a_{i_0} \leq a_{i_1} \leq a_{i_2} \leq \dots$*

Proof. If for every k there is an i with $a_i \geq k$, it is evident. In the opposite case there is a number k with $a_i \leq k$ for all i . Then there is a number k with $a_i = k$ for infinitely many numbers i and everything is evident. \square

Lemma 4. *Let f_0, f_1, f_2, \dots be an infinite sequence of non-increasing functions from ω^n (where $n \in \omega$ is fixed). Then there exists an infinite sequence $i_0 < i_1 < i_2 < \dots$ with $f_{i_0} \leq f_{i_1} \leq f_{i_2} \leq \dots$*

Proof. By induction on n . For $n = 0$ it is evident. Let $n > 0$. By induction, there is a sequence $j_0 < j_1 < \dots$ such that $f_{j_0} \upharpoonright n - 1 \leq f_{j_1} \upharpoonright n - 1 \leq f_{j_2} \upharpoonright n - 1 \leq \dots$. By Lemma 3 there is a sequence $k_0 < k_1 < k_2 < \dots$ such that $f_{j_{k_0}}(n - 1) \leq f_{j_{k_1}}(n - 1) \leq f_{j_{k_2}}(n - 1) \leq \dots$. Then evidently $f_{j_{k_0}} \leq f_{j_{k_1}} \leq f_{j_{k_2}} \leq \dots$ \square

Lemma 5. *Let f_0, f_1, f_2, \dots be an infinite sequence of non-increasing functions from ω^ω . Then there exist i, j with $i \neq j$ and $f_i \leq f_j$.*

Proof. Suppose that there are no such i, j . Denote by m the minimum of all $f_i(j)$. Let us fix a pair c, d with $f_c(d) = m$. For every $i \neq c$ there exists a number j with $f_i(j) < f_c(j)$; necessarily, $j < d$. Hence there is a number $j_0 < d$ such that $f_i(j_0) < f_c(j_0)$ for infinitely many numbers i . It follows from Lemma 2 that there is a sequence $k_0 < k_1 < k_2 < \dots$ such that $f_{k_0} \upharpoonright (\omega - j_0) \leq f_{k_1} \upharpoonright (\omega - j_0) \leq f_{k_2} \upharpoonright (\omega - j_0) \leq \dots$. It follows from Lemma 4 that there is a sequence $l_0 < l_1 < l_2 < \dots$ such that $f_{k_{l_0}} \upharpoonright j_0 \leq f_{k_{l_1}} \upharpoonright j_0 \leq f_{k_{l_2}} \upharpoonright j_0 \leq \dots$. Now, $f_{k_{l_0}} \leq f_{k_{l_1}}$ and $k_{l_0} \neq k_{l_1}$. \square

Theorem 6. *The set $\{\mathcal{F}, \mathcal{B}_A\}$ is not incomparably continuable.*

Proof. Suppose that there is an infinite sequence $\mathcal{F}, \mathcal{B}_A, A_1, A_2, \dots$ of pairwise incomparable finite semilattices. For every nonincreasing sequence f from ω^ω which is non-zero on only a finite subset of ω define a finite semilattice S_f with zero 0 and elements $0 < a_{0,1} < \dots < a_{0,f(0)}, 0 < a_{1,1} < \dots < a_{1,f(1)}, \dots, 0 < a_{k,1} < \dots < a_{k,f(k)}$ where k is the greatest number with $f(k) \neq 0$. Then every A_i is some S_f . Since S_f is embeddable into S_g if $f \leq g$, the result follows from Lemma 5. \square

It would be possible to formulate various open problems concerning incomparably continuable sets of semilattices (or other kinds of algebraic systems). Let us point out just one open problem:

Problem. Is there an algorithm deciding, for any finite set of finite semilattices, whether the set is incomparably continuable?

References

- [1] *R. McKenzie, G. McNulty, W. Taylor: Algebras, Lattices, Varieties, Vol. I. Wadsworth & Brooks/Cole, Monterey, CA, 1987.*

Authors' addresses: Jaroslav Ježek, Charles University, Sokolovská 83, 186 75 Praha 8, Czech Republic, e-mail: jezek@karlin.mff.cuni.cz; Václav Slavík, Czech Agricultural University, Kamýčká 129, 165 21 Praha 6, Czech Republic, e-mail: slavik@tf.czu.cz.