

Róbert Vrábel'

Quasilinear and quadratic singularly perturbed Neumann's problem

Mathematica Bohemica, Vol. 123 (1998), No. 4, 405–410

Persistent URL: <http://dml.cz/dmlcz/125970>

Terms of use:

© Institute of Mathematics AS CR, 1998

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

QUASILINEAR AND QUADRATIC SINGULARLY PERTURBED
NEUMANN'S PROBLEM

RÓBERT VRÁBEL, Trnava

(Received June 6, 1997)

Abstract. The problem of existence and asymptotic behaviour of solutions of the quasilinear and quadratic singularly perturbed Neumann's problem as a small parameter at the highest derivative tends to zero is studied.

Keywords: singularly perturbed Neumann's problem

MSC 1991: 05C10, 05C75

1. INTRODUCTION

In the paper [2] the author established sufficient conditions for the existence and uniform convergence of the solutions of a semilinear singularly perturbed differential equation $\varepsilon y'' + ky = f(t, y)$ to a solution of the reduced problem $ku = f(t, u)$ as the small positive parameter ε tends to zero. The purpose of this paper is an extension of Theorem 1 of the above cited paper to more general cases. We will consider Neumann's problem

$$(NP_0) \quad \begin{aligned} \varepsilon y'' &= F(t, y, y'), & a < t < b, \\ y'(a, \varepsilon) &= 0, & y'(b, \varepsilon) &= 0, \end{aligned}$$

where $F \in C^1([a, b] \times \mathbb{R}^2)$ and ε is a small positive parameter. The proofs of the theorems are based upon the method of lower and upper solutions.

As usual, we say that $\alpha \in C^2([a, b])$ is a lower solution for (NP_0) if $\alpha'(a, \varepsilon) \geq 0$, $\alpha'(b, \varepsilon) \leq 0$, and $\varepsilon \alpha''(t, \varepsilon) \geq F(t, \alpha(t, \varepsilon), \alpha'(t, \varepsilon))$ for every $t \in [a, b]$. An upper solution $\beta \in C^2([a, b])$ satisfies $\beta'(a, \varepsilon) \leq 0$, $\beta'(b, \varepsilon) \geq 0$, and $\varepsilon \beta''(t, \varepsilon) \leq F(t, \beta(t, \varepsilon), \beta'(t, \varepsilon))$ for every $t \in [a, b]$.

Definition 1. We say that a function F satisfies the Bernstein-Nagumo condition if for each $M > 0$ there exists a continuous function $h_M : [0, \infty) \rightarrow [a_M, \infty)$ with $a_M > 0$ and $\int_0^\infty \frac{s}{h_M(s)} ds = \infty$ such that for all $y, |y| \leq M$, all $t \in [a, b]$ and all $z \in \mathbb{R}$ we have

$$|F(t, y, z)| \leq h_M(|z|).$$

Remark. As a remark we conclude that the functions of the form $F(t, y, y') = f(t, y)y' + g(t, y)$ and $F(t, y, y') = f(t, y)y'^2 + g(t, y)$ satisfy the Bernstein-Nagumo condition.

Lemma 1. If α, β are lower and upper solutions for (NP_0) such that $\alpha(t, \varepsilon) \leq \beta(t, \varepsilon)$ on $[a, b]$ and F satisfies the Bernstein-Nagumo condition, then there exists a solution y of (NP_0) with $\alpha(t, \varepsilon) \leq y(t, \varepsilon) \leq \beta(t, \varepsilon)$, $a \leq t \leq b$.

Notation. Let

$$D_\delta(u) = \{(t, y) \in \mathbb{R}^2 : a \leq t \leq b, |y - u(t)| < \delta\},$$

$$D_{\delta, a}(u) = \{(t, y) \in \mathbb{R}^2 : a \leq t \leq a + \delta, y \in \mathbb{R}\} \cap D_\delta(u),$$

and

$$D_{\delta, b}(u) = \{(t, y) \in \mathbb{R}^2 : b - \delta \leq t \leq b, y \in \mathbb{R}\} \cap D_\delta(u),$$

where $\delta \leq b - a$ is a positive constant and $u = u(t)$ is a solution of the reduced problem $F(t, u, u') = 0$ defined on $[a, b]$ such that $u \in C^2([a, b])$.

Let $h(t, y)$ denote $F(t, y, u'(t))$.

2. QUASILINEAR NEUMANN'S PROBLEM

In this section we consider the quasilinear Neumann's problem

$$(NP_1) \quad \begin{aligned} \varepsilon y'' &= f(t, y)y' + g(t, y), & a < t < b, \\ y'(a, \varepsilon) &= 0, & y'(b, \varepsilon) &= 0, \end{aligned}$$

where $f, g \in C^1(D_\delta(u))$. Concerning the behaviour of solutions of (NP_1) for $\varepsilon \rightarrow 0^+$ we have the following result.

Theorem 1. Consider the problem (NP_1) . Let there exist a solution $u \in C^2([a, b])$ of the reduced problem. Let δ, m be positive constants such that $\frac{\partial h(t, y)}{\partial y} \geq m$ for every $(t, y) \in D_\delta(u)$. Let $f(t, y) \leq 0$ and $f(t, y) \geq 0$ for every $(t, y) \in D_{\delta, a}(u)$

and $(t, y) \in D_{\delta, b}(u)$, respectively. Then there exists ε_0 such that for any $\varepsilon \in (0, \varepsilon_0]$ the problem (NP₁) has a solution satisfying the inequality

$$|y(t, \varepsilon) - u(t)| \leq v_1(t, \varepsilon) + v_2(t, \varepsilon) + C\varepsilon$$

on $[a, b]$, where

$$v_1(t, \varepsilon) = |u'(a)| \frac{\exp[-\sqrt{\frac{m}{\varepsilon}}(b-t)] + \exp[-\sqrt{\frac{m}{\varepsilon}}(t-b)]}{\sqrt{\frac{m}{\varepsilon}} (\exp[\sqrt{\frac{m}{\varepsilon}}(b-a)] - \exp[-\sqrt{\frac{m}{\varepsilon}}(b-a)])},$$

$$v_2(t, \varepsilon) = |u'(b)| \frac{\exp[-\sqrt{\frac{m}{\varepsilon}}(a-t)] + \exp[-\sqrt{\frac{m}{\varepsilon}}(t-a)]}{\sqrt{\frac{m}{\varepsilon}} (\exp[\sqrt{\frac{m}{\varepsilon}}(b-a)] - \exp[-\sqrt{\frac{m}{\varepsilon}}(b-a)])}$$

and C is a positive constant.

Proof. We define the lower solutions by

$$\alpha(t, \varepsilon) = u(t) - v_1(t, \varepsilon) - v_2(t, \varepsilon) - \Gamma(\varepsilon)$$

and the upper solutions by

$$\beta(t, \varepsilon) = u(t) + v_1(t, \varepsilon) + v_2(t, \varepsilon) + \Gamma(\varepsilon).$$

Here $\Gamma(\varepsilon) = \frac{\varepsilon\gamma}{m}$, where γ is a constant which will be defined below. One can easily check that the functions α, β satisfy the boundary conditions required for the lower and upper solutions of (NP₁) and $\alpha \leq \beta$ on $[a, b]$. Now we show that $\varepsilon\alpha''(t, \varepsilon) \geq f(t, \alpha(t, \varepsilon))\alpha'(t, \varepsilon) + g(t, \alpha(t, \varepsilon))$ and $\varepsilon\beta''(t, \varepsilon) \leq f(t, \beta(t, \varepsilon))\beta'(t, \varepsilon) + g(t, \beta(t, \varepsilon))$ on $[a, b]$. By the Taylor theorem we obtain

$$\begin{aligned} \varepsilon\alpha'' - F(t, \alpha, \alpha') &= \varepsilon\alpha'' - (F(t, \alpha, \alpha') - F(t, u, u')) \\ &= \varepsilon\alpha'' - [(F(t, \alpha, u') - F(t, u, u')) + (F(t, \alpha, \alpha') - F(t, \alpha, u'))] \\ &= \varepsilon\alpha'' - \left[\frac{\partial h(t, \eta(t, \varepsilon))}{\partial y} (\alpha - u) + f(t, \alpha) (\alpha' - u') \right] \\ &= \varepsilon u'' - \varepsilon v_1'' - \varepsilon v_2'' + \frac{\partial h(t, \eta)}{\partial y} (v_1 + v_2 + \Gamma) + f(t, \alpha) (v_1' + v_2') \\ &\geq \varepsilon u'' - \varepsilon v_1'' - \varepsilon v_2'' + m(v_1 + v_2 + \Gamma) + f(t, \alpha) (v_1' + v_2') \\ &= \varepsilon u'' + \varepsilon\gamma + f(t, \alpha) (v_1' + v_2') \\ &\geq -\varepsilon |u''| + \varepsilon\gamma + f(t, \alpha) (v_1' + v_2') \end{aligned}$$

and

$$F(t, \beta, \beta') - \varepsilon\beta'' \geq -\varepsilon |u''| + \varepsilon\gamma + f(t, \beta) (v_1' + v_2'),$$

where $(t, \eta(t, \varepsilon))$ is a point between $(t, \alpha(t, \varepsilon))$ and $(t, u(t))$, $(t, \eta(t, \varepsilon)) \in D_\delta(u)$ for sufficiently small ε .

Let $u'(a) \neq 0$, $u'(b) \neq 0$ (if $u'(a) = 0$ or $u'(b) = 0$, we proceed analogously). From the above assumptions we obtain that $f(t, \alpha)(v'_1 + v'_2) \geq 0$ and $f(t, \beta)(v'_1 + v'_2) \geq 0$ on $[a, a + \bar{\delta}] \cup [b - \bar{\delta}, b]$ for $\varepsilon \in (0, \varepsilon_1]$ where $\bar{\delta} = \min\{\delta, \delta_1\}$, and δ_1, ε_1 are such that $v'_1 + v'_2 < 0$ ($v'_1 + v'_2 > 0$) on $[a, a + \delta_1]$ ($[b - \delta_1, b]$) and $(t, \alpha) \subset D_{\bar{\delta}}(u)$, $(t, \beta) \subset D_{\bar{\delta}}(u)$ for $\varepsilon \in (0, \varepsilon_1]$. On the interval $[a + \bar{\delta}, b - \bar{\delta}]$ we have $|f(t, \alpha)(v'_1 + v'_2)| \leq c_1\varepsilon$ and $|f(t, \beta)(v'_1 + v'_2)| \leq c_1\varepsilon$ for sufficiently small ε , for instance if $\varepsilon \in (0, \varepsilon_0]$, $\varepsilon_0 \leq \varepsilon_1$ and c_1 is a suitable positive constant (if $u'(a) = 0$ ($u'(b) = 0$) then $|f(t, \alpha)(v'_1 + v'_2)| \leq c_1\varepsilon$ and $|f(t, \beta)(v'_1 + v'_2)| \leq c_1\varepsilon$ on $[a, b - \bar{\delta}]$ ($[a + \bar{\delta}, b]$)).

Thus if we choose a constant $\gamma \geq c_1 + \max\{|u''(t)|, t \in [a, b]\}$ then $\varepsilon\alpha''(t, \varepsilon) \geq f(t, \alpha(t, \varepsilon))\alpha'(t, \varepsilon) + g(t, \alpha(t, \varepsilon))$ and $\varepsilon\beta''(t, \varepsilon) \leq f(t, \beta(t, \varepsilon))\beta'(t, \varepsilon) + g(t, \beta(t, \varepsilon))$ on $[a, b]$. The existence of a solution of (NP₁) satisfying the above inequalities follows from Lemma. This completes the proof. \square

Example 1. As an illustrative example we consider the (NP₁) for the differential equation $\varepsilon y'' = yy' - (t - \frac{1}{2})$ on $[0, 1]$. General solution of the reduced problem $yy' - (t - \frac{1}{2}) = 0$ is $u^2 = t^2 - t + k$, $k \in \mathbb{R}$; however, only $u(t) = t - \frac{1}{2}$ satisfies the assumptions asked on the solution of the reduced problem. On the basis of Theorem 1, there is ε_0 such that for every $\varepsilon \in (0, \varepsilon_0]$ the problem has a solution satisfying $|y(t, \varepsilon) - (t - \frac{1}{2})| \leq v_1 + v_2 + c_1\varepsilon$ on $[0, 1]$.

3. QUADRATIC NEUMANN'S PROBLEM

Now we will consider the quadratic Neumann's problem

$$(NP_2) \quad \begin{aligned} \varepsilon y'' &= f(t, y)y'^2 + g(t, y), & a < t < b, \\ y'(a, \varepsilon) &= 0, & y'(b, \varepsilon) &= 0, \end{aligned}$$

where $f, g \in C^1(D_\delta(u))$.

Theorem 2. Consider the problem (NP₂). Let there exist a solution $u \in C^2([a, b])$ of the reduced problem. Let δ, m be positive constants such that $\frac{\partial h(t, y)}{\partial y} \geq m$ for every $(t, y) \in D_\delta(u)$. Let $f(t, y) \leq 0$ ($f(t, y) \geq 0$) for $(t, y) \in D_{\delta, a}(u)$ when $u'(a) > 0$ ($u'(a) < 0$) and $f(t, y) \leq 0$ ($f(t, y) \geq 0$) for $(t, y) \in D_{\delta, b}(u)$ when $u'(b) < 0$ ($u'(b) > 0$). Then there exists ε_0 such that for any $\varepsilon \in (0, \varepsilon_0]$ the problem (NP₂) has a solution satisfying the inequality

$$|y(t, \varepsilon) - u(t)| \leq v_1(t, \varepsilon) + v_2(t, \varepsilon) + C\varepsilon$$

on $[a, b]$ where v_1, v_2 are the functions from Theorem 1 and C is a positive constant.

P r o o f. The idea of the proof is essentially the same as in the proof of Theorem 1. Let us define the lower solutions by

$$\alpha(t, \varepsilon) = u(t) - v_1(t, \varepsilon) - v_2(t, \varepsilon) - \Gamma(\varepsilon)$$

and the upper solutions by

$$\beta(t, \varepsilon) = u(t) + v_1(t, \varepsilon) + v_2(t, \varepsilon) + \Gamma(\varepsilon).$$

Analogously as in Theorem 1 we obtain

$$\begin{aligned} \varepsilon \alpha'' - F(t, \alpha, \alpha') &\geq -\varepsilon |u''| + \gamma \varepsilon - f(t, \alpha)(\alpha^2 - u^2) \\ &= -\varepsilon |u''| + \gamma \varepsilon + f(t, \alpha)(v'_1 + v'_2)(2u' - v'_1 - v'_2) \end{aligned}$$

and

$$\begin{aligned} F(t, \beta, \beta') - \varepsilon \beta'' &\geq -\varepsilon |u''| + \gamma \varepsilon + f(t, \beta)(\beta^2 - u^2) \\ &= -\varepsilon |u''| + \gamma \varepsilon + f(t, \beta)(v'_1 + v'_2)(2u' + v'_1 + v'_2). \end{aligned}$$

Similarly as in the previous theorem we conclude (for $u'(a) \neq 0, u'(b) \neq 0$) that

$$f(t, \alpha)(v'_1 + v'_2)(2u' - v'_1 - v'_2) \geq 0, \quad f(t, \beta)(v'_1 + v'_2)(2u' + v'_1 + v'_2) \geq 0$$

on $[a, a + \delta] \cup [b - \delta, b]$ and

$$\begin{aligned} |f(t, \alpha)(v'_1 + v'_2)(2u' - v'_1 - v'_2)| &\leq c_2 \varepsilon, \\ |f(t, \beta)(v'_1 + v'_2)(2u' + v'_1 + v'_2)| &\leq c_2 \varepsilon \end{aligned}$$

on $[a + \delta, b - \delta]$ for $\varepsilon \in (0, \varepsilon_0]$, sufficiently small $\delta > 0$ and a suitable positive constant c_2 . Therefore, for $\gamma \geq c_2 + \max\{|u''(t)|, t \in [a, b]\}$ we have

$$\varepsilon \alpha''(t, \varepsilon) \geq f(t, \alpha(t, \varepsilon)) \alpha'^2(t, \varepsilon) + g(t, \alpha(t, \varepsilon))$$

and

$$\varepsilon \beta''(t, \varepsilon) \leq f(t, \beta(t, \varepsilon)) \beta'^2(t, \varepsilon) + g(t, \beta(t, \varepsilon))$$

on $[a, b]$. Hence Theorem 2 is proved. □

Example 2. Consider problem (NP_2) for differential equation $\varepsilon y'' = yy'^2 - (t+1)$ on $[-2, 1]$. Obviously $u(t) = t+1$ is the only solution of the reduced problem $uu'^2 - (t+1) = 0$ satisfying the assumptions of Theorem 2. Hence, there is ε_0 such that for every $\varepsilon \in (0, \varepsilon_0]$ the problem has a solution satisfying

$$|y(t, \varepsilon) - (t+1)| \leq v_1 + v_2 + c_2\varepsilon$$

on $[-2, 1]$.

References

- [1] V. Šeda: On some non-linear boundary value problems for ordinary differential equations. Arch. Math. (Brno) 25 (1989), 207–222.
- [2] R. Vrábek: Upper and lower solutions for singularly perturbed Neumann's problem. Math. Bohem. 122 (1997), 175–180.

Author's address: Róbert Vrábek, Department of Mathematics, Slovak Technical University, 917 24 Trnava, Slovakia, e-mail: vrabel@sun.mtf.stuba.sk.