

Jozef Džurina

Property (A) of n -th order ODE's

Mathematica Bohemica, Vol. 122 (1997), No. 4, 349–356

Persistent URL: <http://dml.cz/dmlcz/126218>

Terms of use:

© Institute of Mathematics AS CR, 1997

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

PROPERTY (A) OF n -TH ORDER ODE'S

JOZEF DŽURINA, Košice

(Received April 9, 1996)

Abstract. The aim of this paper is to deduce oscillatory and asymptotic behavior of the solutions of the ordinary differential equation

$$L_n u(t) + p(t)u(t) = 0.$$

Keywords: property (A) of ODE's

MSC 1991: 34C10

Consider the n -th order ($n \geq 2$) differential equation

$$(1) \quad L_n u(t) + p(t)u(t) = 0,$$

where

$$L_n u(t) = \left(\frac{1}{r_{n-1}(t)} \left(\frac{1}{r_{n-2}(t)} \cdots \left(\frac{1}{r_1(t)} u'(t) \right)' \cdots \right)' \right)',$$

p and $r_i : (t_0, \infty) \rightarrow \mathbb{R}^+ = (0, \infty)$ are continuous, $1 \leq i \leq n-1$. In the sequel we will suppose that $\int_{t_0}^{\infty} r_i(s) ds = \infty$ for $1 \leq i \leq n-1$. It is usual to denote

$$(2) \quad \begin{aligned} D_0 u(t) &= u(t), \\ D_i u(t) &= \frac{1}{r_i(t)} \frac{d}{dt} D_{i-1} u(t), \quad 1 \leq i \leq n-1, \\ D_n u(t) &= \frac{d}{dt} D_{n-1} u(t). \end{aligned}$$

By a solution of Eq. (1) we mean a function $u : (T_u, \infty) \rightarrow \mathbb{R}$ such that

- (i) $D_i u(t)$, $0 \leq i \leq n$ exist and are continuous on $[T_u, \infty)$;

- (ii) $u(t)$ satisfies (1);
- (iii) $\sup \{|u(s)| : t \leq s < \infty\} > 0$ for any $t \geq T_u$.

Such a solution of (1) is called oscillatory if it has arbitrarily large zeros; otherwise it is called nonoscillatory. An equation is said to be oscillatory if all its solutions are oscillatory.

It is well known (see e.g. [2] or [3]) that the set \mathcal{N} of all nonoscillatory solutions of (1) can be divided into the following classes:

$$\begin{aligned} \mathcal{N} &= \mathcal{N}_0 \cup \mathcal{N}_2 \cup \dots \cup \mathcal{N}_{n-1} \quad \text{for } n \text{ odd,} \\ \mathcal{N} &= \mathcal{N}_1 \cup \mathcal{N}_3 \cup \dots \cup \mathcal{N}_{n-1} \quad \text{for } n \text{ even,} \end{aligned}$$

where $u(t) \in \mathcal{N}_\ell$ if and only if

$$(3) \quad \begin{aligned} u(t)D_i u(t) &> 0, & 0 \leq i \leq \ell, \\ (-1)^{i-\ell} u(t)D_i u(t) &> 0, & \ell \leq i \leq n \end{aligned}$$

for all large t . Following Foster and Grimmer [3] we say that $u(t)$ is a function of degree ℓ if $u(t)$ satisfies (3).

For the class \mathcal{N}_0 of (1), it is shown in [4] that $\mathcal{N}_0 \neq \emptyset$ if n is odd. Therefore, we are interested in the following particular situation:

Definition 1. Equation (1) is said to have property (A) if for n even $\mathcal{N} = \emptyset$ (i.e. (1) is oscillatory) and for n odd $\mathcal{N} = \mathcal{N}_0$.

This definition can be found in [6]. There is much literature regarding property (A) of (1) (see enclosed references). Integral conditions have been given under which (1) enjoys property (A). The following result is due to Trench [18].

Define for $1 \leq k \leq n-1$ and $t, s \in [t_0, \infty)$

$$\begin{aligned} I_0 &= 1, \\ I_k(t, s; r_1, \dots, r_k) &= \int_s^t r_1(x) I_{k-1}(x, s; r_2, \dots, r_k) dx, \\ J_k(t) &= I_k(t_0, t; r_1, \dots, r_k), \\ N_k(t) &= I_k(t_0, t; r_{n-1}, \dots, r_{n-k}). \end{aligned}$$

Theorem A. Let n be even. Assume that for all $i \in \{1, 3, \dots, n-1\}$

$$(4) \quad \int_{t_0}^{\infty} J_{i-1}(t) N_{n-i-1}(t) p(t) dt = \infty.$$

Then (1) has property (A).

A question naturally arises what will happen when conditions (4) are violated. In fact, Theorem A cannot cover an important class of Euler's equation of the form

$$(5) \quad \frac{d^m}{dt^m} t^{\alpha+m} \frac{d^m x}{dt^m} + ct^{\alpha-m} x = 0, \quad t \geq 1,$$

where α and $c > 0$ are constants with $\alpha + m \leq 1$, since in this case the integrals in (4) converge.

Trench's result has been later improved by Kusano, Naito and Tanaka in [6] and [7], where (1) is compared with a set of second order differential equations and property (A) of (1) is reduced to the oscillation of a set of second order differential equations. On the other hand, Chanturia and Kiguradze [1] have improved (4) for the particular case of (1), namely for the differential equation

$$(6) \quad y^{(n)}(t) + p(t)y(t) = 0.$$

They have compared (1) with Euler's equation $t^n y^{(n)} + cy = 0$ to obtain the integral criterion

$$\liminf_{t \rightarrow \infty} t^{n-1} \int_t^\infty p(s) ds = \frac{M^*}{n-1}$$

for property (A) of (6).

Our concern in this paper is to replace condition (4) by a similar one that is applicable also to (5). Our results complement and extend the above-mentioned results and also some other ones given in [16], [14], [10] and [8].

We consider a set of ℓ -th order ($n-1 \geq \ell \geq 1$) differential inequalities

$$(E_{\ell+1}) \quad \{M_{\ell+1}u(t) + q_{\ell+1}(t)u(t)\} \operatorname{sgn} u(t) \leq 0,$$

where $q_{\ell+1}$ is positive and continuous and

$$M_{\ell+1}u(t) = \left(\frac{1}{r_\ell(t)} \left(\frac{1}{r_{\ell-1}(t)} \dots \left(\frac{1}{r_1(t)} u'(t) \right)' \dots \right)' \right)',$$

that is $M_{\ell+1}u(t) = r_{\ell+1}(t)D_{\ell+1}u(t)$ for $\ell < n$, and $M_n u(t) = D_n u(t)$.

Let us put

$$J_{1,\ell}(t) = J_\ell(t) \quad \text{and} \quad J_{2,\ell}(t) = I_{\ell-1}(t, t_0; r_2, \dots, r_\ell).$$

Our main results are based on the following theorem:

Theorem 1. Let $1 \leq \ell \leq n - 1$. Assume that

$$(7_\ell) \quad \int^\infty \left(J_{1,\ell}(t)q_{\ell+1}(t) - \frac{r_1(t)J_{2,\ell}(t)}{4J_{1,\ell}(t)} \right) dt = \infty.$$

Then $(E_{\ell+1})$ has no solutions of degree ℓ .

P r o o f. Assume that $(E_{\ell+1})$ possesses a positive nonoscillatory solution $u(t)$ such that $u(t)$ is of degree ℓ , that is

$$D_0u(t) > 0, D_1u(t) > 0, \dots, D_\ell u(t) > 0, (D_{\ell+1}u(t))' < 0, \quad t \geq t_0.$$

Let

$$z(t) = \frac{J_{1,\ell}(t)D_\ell u(t)}{u(t)}, \quad t \geq t_0.$$

Then $z(t) > 0$ and

$$(8) \quad z'(t) = \frac{r_1(t)J_{2,\ell}(t)}{J_{1,\ell}(t)}z(t) + \frac{J_{1,\ell}(t)(D_\ell u(t))'}{u(t)} - z(t)\frac{r_1(t)D_1u(t)}{u(t)}.$$

Assume that $\ell > 1$. The identity $D_\ell u(t) = \frac{1}{r_\ell(t)}(D_{\ell-1}u(t))'$ implies that

$$\begin{aligned} D_{\ell-1}u(t) &= D_{\ell-1}u(t_0) + \int_{t_0}^t r_\ell(s)D_\ell u(s) ds \\ &\geq D_\ell u(t) \int_{t_0}^t r_\ell(s) ds. \end{aligned}$$

Hence, after $(\ell - 3)$ -fold integration, we arrive at

$$D_1u(t) \geq J_{2,\ell}(t)D_\ell u(t), \quad t \geq t_0.$$

Therefore, combining (8) with the last inequality, one gets

$$(9) \quad \frac{J_{1,\ell}(t)(D_\ell u(t))'}{u(t)} \geq z'(t) + \frac{r_1(t)J_{2,\ell}(t)}{J_{1,\ell}(t)}(z^2(t) - z(t)).$$

Note that $z^2(t) - z(t) \geq -\frac{1}{4}$. Multiplying $(E_{\ell+1})$ by $J_{1,\ell}(t)$ and dividing the resulting equality by $u(t)$, we see in view of (9) that $z(t)$ is a positive solution of the differential inequality

$$(10) \quad z'(t) - \frac{r_1(t)J_{2,\ell}(t)}{4J_{1,\ell}(t)} + J_{1,\ell}(t)q_{\ell+1}(t) \leq 0.$$

That (10) also holds for $\ell = 1$ follows from (8) and (M_2) (note that $J_{2,1}(t) \equiv 1$). An integration of (10) yields

$$z(t) + \int_{t_0}^t \left(J_{1,\ell}(s)q_{\ell+1}(s) - \frac{r_1(s)J_{2,\ell}(s)}{4J_{1,\ell}(s)} \right) ds \leq z(t_0).$$

Letting $t \rightarrow \infty$, we get a contradiction with (7_ℓ) . The proof is complete. \square

The following result can be found in [5, Corollary 1].

Theorem B. *The equation (1) has a solution of degree $n - 1$ if and only if the inequality (E_n) has a solution of degree $n - 1$.*

For the particular case of (1) with $n = 2$ and $n = 3$ we have the following corollaries.

Corollary 1. *Denote $R(t) = \int_{t_0}^t r(s) ds$. Assume that*

$$(11) \quad \int^{\infty} \left(R(s)p(s) - \frac{r(s)}{4R(s)} \right) ds = \infty.$$

Then the second order differential equation

$$(12) \quad \left(\frac{1}{r(t)} u' \right)' + p(t)u = 0$$

is oscillatory.

Proof. By Theorem B, Eq. (12) is oscillatory if and only if (E_2) with $q_2 = p$ and $r_1 = r$ has no solution of degree 1. Since (7_1) reduces to (11), the assertion of this corollary follows from Theorem 1. \square

Corollary 2. *Assume that*

$$(13) \quad \int^{\infty} \left(J_{1,2}(s)p(s) - \frac{r_1(s)J_{2,2}(s)}{4J_{1,2}(s)} \right) ds = \infty.$$

Then the third order differential equation

$$\left(\frac{1}{r_2(t)} \left(\frac{1}{r_1(t)} u' \right)' \right)' + p(t)u = 0$$

has property (A).

Proof. The proof of this corollary is analogous to that of Corollary 1 (noting that (7_2) reduces to (13)) and can be omitted. \square

Example 1. Consider the equation

$$\left(\frac{1}{t} u'' \right)' + \frac{a}{t^4} u = 0, \quad a > 0, \quad t \geq 1.$$

By Corollary 2, this equation has property (A) provided $a > 4.5$.

Now we extend our previous results to (1) with $n > 3$. For all large t and $i \in \{1, \dots, n-1\}$ define

$$\begin{aligned} K_1(t; p) &= \int_t^\infty p(s) \, ds, \\ K_2(t; r_{n-1}, p) &= \int_t^\infty r_{n-1}(x) K_1(x; p) \, dx, \\ K_i(t; r_{n-i+1}, \dots, r_{n-1}, p) &= \int_t^\infty r_{n-i+1}(x) K_{i-1}(x; r_{n-i+2}, \dots, r_{n-1}, p) \, dx, \\ q_n(t) &= p(t), \\ q_i(t) &= r_i(t) K_{n-i}(t; r_{i+1}, \dots, r_{n-1}, p). \end{aligned}$$

Theorem 2. Assume that for all $\ell \in \{1, \dots, n-1\}$ with $n + \ell$ odd, conditions (7_ℓ) are satisfied. Then (1) has property (A).

Proof. Since (7₁) with $n = 2$ reduces to (11) and (7₂) with $n = 3$ reduces to (13) the assertion of the theorem for $n = 2$ and $n = 3$ follows from Corollaries 1 and 2.

Now assume that $n > 3$. We want to show that $\mathcal{N}_\ell = \emptyset$ for all $\ell \in \{1, \dots, n-1\}$ with $n + \ell$ odd. Note that by Theorem 1, condition (7_{n-1}) implies that differential inequality (E_n) has no solution of degree $n-1$. By Theorem (B), Eq. (1) has no solution of degree $n-1$, either (i.e. $\mathcal{N}_{n-1} = \emptyset$).

Let $1 \leq \ell \leq n-2$. Assume that (1) has a positive nonoscillatory solution $u(t)$ and $u(t)$ is of degree ℓ . From (1) and $u'(t) > 0$ it follows that

$$D_{n-1}u(\infty) - D_{n-1}u(t) + \int_t^\infty p(s)u(s) \, ds = 0, \quad t \geq t_0.$$

That is,

$$-D_{n-1}u(t) + u(t) \int_t^\infty p(s) \, ds \leq 0.$$

Hence, after $(n - \ell - 2)$ -fold integration we arrive at

$$M_{\ell+1}u(t) + q_{\ell+1}(t)u(t) \leq 0.$$

That is, $u(t)$ is a solution of ($E_{\ell+1}$), but as $u(t)$ is of degree ℓ , it contradicts the assertions of Theorem 1. The proof is complete. \square

Corollary 3. Assume that

$$(14) \quad \int^\infty \left((t - t_0)^{n-1} p(t) - \frac{(n-1)(n-1)!}{4(t-t_0)} \right) dt = \infty.$$

Then the n -th order differential equation

$$(15) \quad u^{(n)} + p(t)u = 0$$

has property (A).

Proof. To prove that (15) has property (A), it suffices (see Theorem 1.1 in [1]) to show that (15) has no solution of degree $n - 1$. This fact follows from Theorems A and 1. \square

It is interesting to compare Corollary 3 with the following result which is due to Chanturia and Kiguradze [1].

Lemma A. *The condition*

$$(16) \quad \int_{t_0}^{\infty} t^{n-1}p(t) dt = \infty$$

is necessary for (15) to have property (A).

Note that the stronger condition (13) guarantees property (A) of (14), while (16) is not enough.

Remark. Using suitable comparison theorems, our results can be extended to more general differential equations. In fact, it is known [5] that the delay differential equation

$$(17) \quad L_n u(t) + p(t)u(\tau(t)) = 0,$$

where L_n and p are the same as in (1) and τ satisfies

$$(18) \quad \tau \in C^1, \quad \tau(t) \leq t, \quad \tau(t) \rightarrow \infty \text{ as } t \rightarrow \infty,$$

has property (A) if so does the differential equation without delay

$$(19) \quad L_n u(t) + \frac{p(\tau^{-1}(t))}{\tau'(\tau^{-1}(t))} u(t) = 0,$$

where τ^{-1} is the inverse function to τ . Applying Theorem 2 to (19) we immediately have a sufficient condition for (17) to have property (A). We illustrate this by the following result.

Corollary 4. *Assume that (18) holds. Further assume that*

$$\int_{t_0}^{\infty} \left((\tau(t) - t_0)^{n-1} p(t) - \frac{(n-1)(n-1)!}{4(t-t_0)} \tau'(t) \right) dt = \infty.$$

Then the delay differential equation

$$u^{(n)}(t) + p(t)u(\tau(t)) = 0$$

has property (A).

References

- [1] *T. A. Chanturia and I. T. Kiguradze*: Asymptotic Properties of Solutions of Nonautonomous Ordinary Differential Equations. Nauka, Moscow, 1990. (In Russian.)
- [2] *J. Džurina*: Comparison theorems for functional differential equations. *Math. Nachrichten* 164 (1993), 13–22.
- [3] *K. E. Foster and R. C. Grimmer*: Nonoscillatory solutions of higher order differential equations. *J. Math. Anal. Appl.* 71 (1979), 1–17.
- [4] *P. Hartman and A. Wintner*: Linear differential and difference equations with monotone solutions. *Amer. J. Math.* 75 (1953), 731–743.
- [5] *T. Kusano and M. Naito*: Comparison theorems for functional differential equations with deviating arguments. *J. Math. Soc. Japan* 3 (1981), 509–532.
- [6] *T. Kusano, M. Naito and K. Tanaka*: Oscillatory and asymptotic behavior of solutions of a class of linear ordinary differential equations. *Proc. Roy. Soc. Edinburg.* 90 (1981), 25–40.
- [7] *T. Kusano and M. Naito*: Oscillation criteria for general ordinary differential equations. *Pacific J. Math.* 92 (1981), 345–355.
- [8] *D. Knežo and V. Šoltés*: Existence and properties of nonoscillation solutions of third order differential equations. *Fasciculi Math.* 25 (1995), 63–74.
- [9] *S. Kulcsár*: Boundedness convergence and global stability of solution of a nonlinear differential equations of the second order. *Publ. Math.* 37 (1990), 193–201.
- [10] *G. S. Ladde, V. Lakshmikantham, B. G. Zhang*: Oscillation Theory of Differential Equations with Deviating Arguments. Dekker, New York, 1987.
- [11] *D. L. Lovelady*: An asymptotic analysis of an odd order linear differential equation. *Pacific J. Math.* 57 (1975), 475–480.
- [12] *D. L. Lovelady*: Oscillation of a class of odd order linear differential equations. *Hiroshima Math. J.* 5 (1975), 371–383.
- [13] *W. E. Mahfoud*: Comparison theorems for delay differential equations. *Pacific J. Math.* 83 (1979), 187–197.
- [14] *W. E. Mahfoud*: Characterization of oscillation of solutions of the delay equation $x^{(n)}(t) + a(t)f(x[q(t)]) = 0$. *J. Differential Equations* 28 (1978), 437–451.
- [15] *M. Naito*: On strong oscillation of retarded differential equations. *Hiroshima Math. J.* vol 11 (1981), 553–560.
- [16] *Ch. G. Philos and Y. G. Sficas*: Oscillatory and asymptotic behavior of second and third order retarded differential equations. *Czechoslovak Math. J.* 32 (1982), 169–182.
- [17] *M. Růžičková and E. Špániková*: Oscillation theorems for neutral differential equations with the quasi-derivatives. *Arch. Math.* 30 (1994), 293–300.
- [18] *W. F. Trench*: Oscillation properties of perturbed disconjugate equations. *Proc. Amer. Math. Soc.* 52 (1975), 147–155.

Author's address: Jozef Džurina, Department of Mathematical Analysis, Šafárik University, Jesenná 5, 041 54 Košice, Slovakia, e-mail: dzurina@duro.upjs.sk.