

Ivan Chajda

Tolerances on q -lattices

Czechoslovak Mathematical Journal, Vol. 46 (1996), No. 1, 21–28

Persistent URL: <http://dml.cz/dmlcz/127266>

Terms of use:

© Institute of Mathematics AS CR, 1996

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

TOLERANCES ON q -LATTICES

IVAN CHAJDA, Olomouc

(Received November 3, 1992)

The concept of a q -lattice was introduced for the first time in [1] and some of its congruence properties were studied in [2] and [3]. Recall that an algebra $(A; \wedge, \vee)$ with two binary operations is a q -lattice if it satisfies the following axioms:

(associativity)	$x \vee (y \vee z) = (x \vee y) \vee z,$	$x \wedge (y \wedge z) = (x \wedge y) \wedge z,$
(commutativity)	$x \vee y = y \vee x,$	$x \wedge y = y \wedge x,$
(weak absorption)	$x \vee (x \wedge y) = x \vee x,$	$x \wedge (x \vee y) = x \wedge x,$
(weak idempotence)	$x \vee (y \vee y) = x \vee y,$	$x \wedge (y \wedge y) = x \wedge y,$
(equalization)	$x \vee x = x \wedge x.$	

If, moreover, it satisfies also distributivity:

$$x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z),$$

the q -lattice is called *distributive*.

In every q -lattice A we can distinguish two sorts of elements: *idempotents*, i.e. such $x \in A$ for which $x = x \vee x$ (and hence also $x = x \wedge x$), and *non-idempotents* (i.e. $x \neq x \vee x$). Denote by S_A the so called *skeleton of A* , i.e. S_A is the set of all idempotents of A . It is known (see e.g. [1] or [3]) that S_A is a sub- q -lattice of A which is a sublattice with respect to the *induced quasiorder Q* :

$$\langle a, b \rangle \Leftrightarrow a \vee b = b \vee b,$$

i.e. $Q \cap S_A^2$ is an order on S_A (for some details, see [1]).

The non-idempotents occur in A in the so called cells: a subset $C_x \subseteq A$ is called a *cell* (with the idempotent x) if $\text{card } C_x > 1$ and for each $a, b \in C_x$, $a \vee a = b \vee b (= x)$.

The aim of this paper is to characterize q -lattices with distributive lattices of tolerances.

By a *tolerance* on $(A; \wedge, \vee)$ we mean a reflexive and symmetric binary relation on A satisfying the substitution property with respect operations \vee and \wedge . Denote by $\text{Tol } A$ the lattice of all tolerance of $(A; \wedge, \vee)$ (for some details on $\text{Tol } A$ and the basic properties of tolerances, see the monograph [4]). In particular, denote by ω (or ι) the least (greatest) element of $\text{Tol } A$, i.e. ω is the identity relation on A and $\iota = A \times A$. If $a, b \in A$ denote by $T(a, b)$ the least tolerance on $(A; \wedge, \vee)$ containing the pair $\langle a, b \rangle$.

An algebra A is called *tolerance trivial* if every tolerance on A is a congruence, i.e. if $\text{Tol } A = \text{Con } A$ (e.g. every boolean or every relative complementary lattice is tolerance trivial, see [4]).

Proposition. *If a q -lattice $(A; \wedge, \vee)$ has at least one non-idempotent element and at least two idempotents, then it is not tolerance trivial.*

Proof. Suppose that $(A; \wedge, \vee)$ has at least one non-idempotent. Then $(A; \wedge, \vee)$ contains at least one cell C . Let S_A be the skeleton of A . Define a binary relation T on A as follows: $\langle x, y \rangle \in T$ if and only if either $x, y \in C$ or $x, y \in S_A$ or $x = y$. It is an easy exercise to show that $T \in \text{Tol } A$. Let x be the unique idempotent of C , let $y \neq x$ be an idempotent of A and z a non-idempotent of C . Then $x, y \in S_A$, i.e. $\langle x, y \rangle \in T$, $x, z \in C$, i.e. $\langle x, z \rangle \in T$ but $\langle y, z \rangle \notin T$ which proves $T \notin \text{Con } A$. \square

Lemma. *Let $(A; \wedge, \vee)$ be a q -lattice and C its cell with the unique idempotent c .*

(i) *Let $p(x_1, \dots, x_n)$ be an n -ary term which is not a projection over $(A; \wedge, \vee)$, and let $a, a_1, \dots, a_n \in A$ and $a_i \in C$ for some i . If $a = p(a_1, \dots, a_n)$ then*

$$a = p(a_1, \dots, a_{i-1}, c, a_{i+1}, \dots, a_n).$$

(ii) *If $T \in \text{Tol } A$, $b \in C$, a is an idempotent and $\langle a, b \rangle \in T$, then $\langle a, c \rangle \in T$.*

Proof. (i) If p is not a projection then p is a composition of operations \vee and \wedge . Hence, $a = p(a_1, \dots, a_n)$ is an idempotent of $(A; \wedge, \vee)$. By induction over the rank of p , suppose first $p(x_1, \dots, x_n) = x_1 \vee x_2$, i.e. $a = a_1 \vee a_2$. If $a_1 \in C$, then clearly $a_1 \vee a_2 = c \vee a_2$; similarly for $i = 2$ and dually for the operation \wedge . By induction, we obtain the first assertion.

(ii) If $\langle a, b \rangle \in T$ and $b \in C$ and c is an idempotent of C , then $b \vee b = c$ and hence $\langle a, c \rangle = \langle a \vee a, b \vee b \rangle \in T$. \square

Theorem 1. *Let $(A; \wedge, \vee)$ be a q -lattice with just one cell C , let S_A be its skeleton. If $\text{Tol } S_A$ is distributive then also $\text{Tol } A$ is distributive.*

Proof. Let $R, S, T \in \text{Tol } A$ and $x, y \in A$. Suppose $\langle x, y \rangle \in R \wedge (S \vee T)$. Then $\langle x, y \rangle \in R$ and there exists an n -ary term $p(x_1, \dots, x_n)$ such that $x = p(a_1, \dots, a_n)$, $y = p(b_1, \dots, b_n)$, where $\langle a_i, b_i \rangle \in S$ or $\langle a_i, b_i \rangle \in T$, see e.g. [4].

(1) If at least one of the elements x, y is non-idempotent, then it cannot be the result of an operation, i.e. p is a projection, therefore $p(a_1, \dots, a_n) = pr_i(a_1, \dots, a_n) = a_i$, $p(b_1, \dots, b_n) = pr_i(b_1, \dots, b_n) = b_i$, thus $\langle x, y \rangle = \langle a_i, b_i \rangle$ and hence $\langle x, y \rangle \in S$ or $\langle x, y \rangle \in T$, i.e. $\langle x, y \rangle \in R \wedge S$ or $\langle x, y \rangle \in R \wedge T$, proving $\langle x, y \rangle \in (R \wedge S) \vee (R \wedge T)$.

(2) Suppose both x, y are idempotents. Then $x, y \in S_A$. By the Lemma, we can substitute all non-idempotents among $a_1, \dots, a_n, b_1, \dots, b_n$ by a unique idempotent $c \in C$ because $(A; \wedge, \vee)$ has just one cell C .

If $\langle a_i, b_i \rangle \in S$ and b_i is a non-idempotent and a_i an idempotent, then $\langle a_i, c \rangle \in S$. Analogously for the converse case and also for T . If both a_i, b_i are non-idempotents, we have $\langle c, c \rangle \in S$ analogously for T . By the Lemma,

$$x = p(a_1^0, \dots, a_n^0), \quad y = p(b_1^0, \dots, b_n^0)$$

where

$$\begin{aligned} a_i^0 &= a_i && \text{if } a_i \text{ is an idempotent and} \\ a_i^0 &= c && \text{in the opposite case,} \\ b_i^0 &= b_i && \text{if } b_i \text{ is an idempotent and} \\ b_i^0 &= c && \text{in the opposite case.} \end{aligned}$$

By the Lemma, $\langle a_i^0, b_i^0 \rangle \in S^0$ or T^0 , where $S^0 = S \cap (S_A \times S_A)$, $T^0 = T \cap (S_A \times S_A)$ are the restrictions of S or T onto the skeleton. But $x, y \in S_A$ implies also $\langle x, y \rangle \in R^0 = R \cap (S_A \times S_A)$. Since $\text{Tol } S_A$ is distributive, we have

$$\langle x, y \rangle \in (R^0 \wedge S^0) \vee (R^0 \wedge T^0) \subseteq (R \wedge S) \vee R \wedge T.$$

Distributivity is proved in both the cases. □

Corollary. *Let $(A; \wedge, \vee)$ be a distributive q -lattice with at most one cell. Then $\text{Tol } A$ is distributive.*

Proof. By [5], for every distributive lattice L , $\text{Tol } L$ is also distributive. If $(A; \wedge, \vee)$ has no cell then $(A; \wedge, \vee)$ is a lattice and $\text{Tol } A$ is therefore distributive. If $(A; \wedge, \vee)$ has just one cell then S_A is a distributive lattice and hence $\text{Tol } S_A$ is distributive. By Theorem 1 we are done. □

Remark 1. If $(A; \wedge, \vee)$ is a q -lattice and C is its cell and S_A its skeleton, then for each $c \in C$ and each $x \in S_A$ there exists a tolerance $T \in \text{Tol } A$ given by

$$T = \omega \cup \{\langle c, x \rangle, \langle x, c \rangle\} \cup (S_A \times S_A).$$

If $\text{Tol } S_A = \{\omega_s, \iota_s\}$ only (i.e. S_A is tolerance simple, see [4]), then all tolerances on A are determined only by the pairs $\langle c, x \rangle$ as was shown before and by all tolerances on C . This is illustrated in the following

Example 1. Let A be a q -lattice with the diagram in Fig. 1.




Fig. 1

It has just one cell $\{z, c\} = C$, z is an idempotent in C . It is evident that $\text{Tol } S_A = \{\omega_s, \iota_s\}$, where $S_A = \{0, x, y, z, 1\}$. Henceforth, for every subset $B \subseteq S_A$ there exists a tolerance $T_B \in \text{Tol } A$ given by

$$T_B = \omega \cup (S_A \times S_A) \cup \{(b, c), (c, b) ; b \in B\}.$$

Since $\text{card } S_A = 5$ we have 2^5 of such subsets; for $B = \emptyset$ we have $T_0 = \omega \cup (S_A \times S_A)$, i.e. it is the congruence collapsing S_A and having two blocks, namely S_A and $\{c\}$, i.e. $T_0 = \theta(0, 1)$. Moreover, $\text{Tol } A$ also contains $\theta(z, c)$ collapsing the cell $C = \{z, c\}$ only and ω and ι , then $\text{Tol } A$ has $2^5 + 2 = 34$ elements, see Fig. 2 (I denotes the two element lattice):




Fig. 2

Example 2. Although $(A; \wedge, \vee)$ can be “nice” and distributive, its $\text{Tol } A$ is rather big in the case if $(A; \wedge, \vee)$ contains a cell. Such $\text{Tol } A$ for a q -lattice visualized in Fig. 3 is the distributive lattice (by the foregoing Corollary) in Fig. 4. All tolerances of $\text{Tol } A$ are listed in Fig. 5.




Fig. 3




Fig. 4

Theorem 2. *If a q -lattice has at least two different cells then Tol A is not modular.*

Proof. Let A have cells $C_1 \neq C_2$, let c_i be the idempotent in C_i , $i = 1, 2$ and let $a \in C_1$, $b \in C_2$ be non-idempotents. Denote by $T(\langle u_1, v_1 \rangle, \dots, \langle u_n, v_n \rangle)$ the least tolerance of Tol A containing the pairs $\langle u_1, v_1 \rangle, \dots, \langle u_n, v_n \rangle$. Now, put

$$\begin{aligned}
 T_0 &= T(\langle a, b \rangle, \langle a, c_1 \rangle), \\
 T_x &= T(\langle a, b \rangle, \langle a, c_1 \rangle, \langle b, c_1 \rangle), \\
 T_y &= T(\langle a, b \rangle, \langle a, c_1 \rangle, \langle b, c_1 \rangle, \langle b, c_2 \rangle), \\
 T_z &= T(\langle a, b \rangle, \langle a, c_1 \rangle, \langle a, c_2 \rangle), \\
 T_1 &= T(\langle a, b \rangle, \langle a, c_1 \rangle, \langle a, c_2 \rangle, \langle b, c_1 \rangle, \langle b, c_2 \rangle).
 \end{aligned}$$

Since $\langle a, b \rangle \in T_i$ for $i \in \{0, x, y, z, 1\}$ and a, b are non-idempotents, we have also $\langle c_1, c_2 \rangle = \langle a \vee a, b \vee b \rangle \in T_i$.




Fig. 5




Fig. 6

- (1) If $c_1 < c_2$, tolerances are visualized in Fig. 6:
- (2) If c_1, c_2 are non-comparable elements (of the skeleton), the situation is visualized in Fig. 7.




Fig. 7

It is routine to show that in both of the foregoing cases, tolerances T_0, T_x, T_y, T_z, T_1 form a sublattice N_5 of $\text{Tol } A$, see Fig. 8. \square




Fig. 8

Remark 2. If $(A; \wedge, \vee)$ is a q -lattice with a skeleton S_A and $\text{Tol } S_A$ is not distributive then $\text{Tol } A$ is not distributive either since $\text{Tol } S_A$ is a sublattice of $\text{Tol } A$.

Corollary. For a distributive q -lattice $(A; \wedge, \vee)$, the following conditions are equivalent:

- (i) $\text{Tol } A$ is distributive;
- (ii) $(A; \wedge, \vee)$ has at most one cell.

References

- [1] *Chajda I.*: Lattices of quasiordered sets. *Acta UP (Olomouc)* 31 (1992), 6–12.
- [2] *Chajda I.*: Subdirectly irreducible algebras of quasiordered logic. *Acta UP (Olomouc)* 32 (1993), 21–26.
- [3] *Chajda I., Kotrle M.*: Subdirectly irreducible and congruence distributive q -lattices. *Czechoslovak Math. J.* 43 (1993), 635–642.
- [4] *Chajda I.*: Algebraic Theory of Tolerance Relations. University Palacký Olomouc Press, 1991.
- [5] *Chajda I., Zelinka B.*: Minimal compatible tolerances on lattices. *Czechoslovak Math. J.* 27 (1977), 452–459.

Author's address: Katedra algebry a geometrie, Přír. fak. UP Olomouc, Tomkova 38, 77900 Olomouc, Czech Republic.