

Salah Mecheri

Commutants and derivation ranges

*Czechoslovak Mathematical Journal*, Vol. 49 (1999), No. 4, 843–847

Persistent URL: <http://dml.cz/dmlcz/127534>

## Terms of use:

© Institute of Mathematics AS CR, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## COMMUTANTS AND DERIVATION RANGES

SALAH MECHERI, Tebessa

(Received February 12, 1997)

*Abstract.* In this paper we obtain some results concerning the set  $\mathcal{M} = \cup\{\overline{R(\delta_A)} \cap \{A\}' : A \in \mathcal{L}(\mathcal{H})\}$ , where  $\overline{R(\delta_A)}$  is the closure in the norm topology of the range of the inner derivation  $\delta_A$  defined by  $\delta_A(X) = AX - XA$ . Here  $\mathcal{H}$  stands for a Hilbert space and we prove that every compact operator in  $\overline{R(\delta_A)}^w \cap \{A^*\}'$  is quasinilpotent if  $A$  is dominant, where  $\overline{R(\delta_A)}^w$  is the closure of the range of  $\delta_A$  in the weak topology.

## INTRODUCTION

Let  $\mathcal{L}(\mathcal{H})$  be the algebra of all bounded linear operators on a complex separable and infinite dimensional Hilbert space  $\mathcal{H}$ , the inner derivation induced by  $A \in \mathcal{L}(\mathcal{H})$  being the map defined by

$$\delta_A: \mathcal{L}(\mathcal{H}) \mapsto \mathcal{L}(\mathcal{H}); \quad \delta_A(X) = AX - XA \quad (A \in \mathcal{L}(\mathcal{H})).$$

The identity is not a commutator, that is,  $I \notin R(\delta_A)$  for any  $A \in \mathcal{L}(\mathcal{H})$ , where  $R(\delta_A)$  denotes the range of  $\delta_A$ . Nevertheless, J.H. Anderson in [2] proved the remarkable result that  $I \in \overline{R(\delta_A)}$  for a large class of operators, where  $\overline{R(\delta_A)}$  denotes the closure of the range of  $\delta_A$  in the norm topology. This allowed him to define a new class of operators, called

$$J_A(\mathcal{H}) = \{A \in \mathcal{L}(\mathcal{H}) : I \in \overline{R(\delta_A)}\}.$$

Let  $\mathcal{N} = \cup\{R(\delta_A) \cap \{A\}' : A \in \mathcal{L}(\mathcal{H})\}$ , where  $\{A\}'$  denotes the commutant of  $A$ . In finite dimension the set  $\mathcal{N}$  is exactly the set of nilpotent operators, in infinite dimension the theorem of Kleĭnecke-Shirokov [3] confirms that any operator in  $\mathcal{N}$  is quasinilpotent. If we now consider instead of  $\mathcal{N}$  the set

$$\mathcal{M} = \cup\{\overline{R(\delta_A)} \cap \{A\}' : A \in \mathcal{L}(\mathcal{H})\},$$

the theorem of Kleineck-Shirokov can't be used. In other words an operator in  $\mathcal{M}$  is not necessarily quasinilpotent; we can take as a counterexample the existence of an operator  $A \in \mathcal{L}(\mathcal{H})$  such that  $I \in \overline{R(\delta_A)}$ .

J.H. Anderson [1, p. 135–136] proved that  $\overline{R(\delta_A)} \cap \{A\}' = \{0\}$  if  $A$  is normal or isometric. Here we prove that any operator in  $\mathcal{M}$  is nilpotent if  $P(A)$  is normal, isometric or co-isometric for some polynomial  $P$ .

R.E. Weber [5] confirms that every compact operator in  $\overline{R(\delta_A)}^w \cap \{A\}'$  is quasinilpotent, where  $\overline{R(\delta_A)}^w$  is the weak closure of  $R(\delta_A)$ . If we now consider the set

$$\left\{ \overline{R(\delta_A)}^w \cap \{A^*\}' : A \in \mathcal{L}(\mathcal{H}) \right\},$$

we can ask: is every compact operator in  $\overline{R(\delta_A)}^w \cap \{A^*\}'$  quasinilpotent? At this moment, we have not a global answer but we can partially answer this question with the assumption that  $A$  is dominant

**Lemma 1.** *Let  $A, X \in \mathcal{L}(H)$ ,  $T \in \{A\}'$  and  $\varepsilon > 0$ . If  $\|A\| \leq 1$  and if  $\|AX - XA - T\| < \varepsilon$ , then for every  $n \in \mathbb{N}$  we have*

$$\|(A^{n+1}X - XA^{n+1}) - (n+1)A^nT\| < (n+1)\varepsilon.$$

We recall that  $\forall A \in \mathcal{L}(H)$ ,  $\forall X \in \mathcal{L}(H)$  and  $\forall T \in \{A\}'$  we have

$$A^nX - XA^n = nA^{n-1}T - \sum_{i=1}^n A^{n-i-1}(T - (AX - XA))A^i.$$

*Proof.* For  $n = 0$  evident.

For  $n = 1$  we have

$$A^2X - XA^2 = (A^2X - AXA) + (AXA - XA^2),$$

so,

$$\begin{aligned} \|(A^2X - XA^2) - 2AT\| &= \|(A^2X - AXA) - AT + (AXA - XA^2) - TA\| \\ &= \|A(AX - XA - T) + (AX - XA - T)A\| \\ &\leq 2\|A\|\|AX - XA - T\| < 2\varepsilon. \end{aligned}$$

Now suppose that for every  $n \geq 2$  and for every  $k \leq n$  we have

$$(*) \quad \|(A^kX - XA^k) - kA^{k-1}T\| < k\varepsilon.$$

Since

$$(A^{n+1}X - X(A^{n+1}) - (n+1)A^nT) = A^n(AX - XA - T) + ((A^nX - XA^n) - nA^{n-1}T)A,$$

we have

$$\|(A^{n+1}X - X(A^{n+1}) - (n+1)A^nT)\| < \varepsilon + n\varepsilon = (n+1)\varepsilon.$$

□

**Theorem 2.** *Let  $A \in \mathcal{L}(\mathcal{H})$  and suppose that*

$$\overline{R(\delta_{P(A)})} \cap \{P(A)\}' = \{0\}$$

*for some polynomial  $P$ , then every operator in  $\overline{R(\delta_A)} \cap \{A\}'$  is nilpotent.*

*Proof.* Let  $P$  be a polynomial of degree  $n$  and let  $P^{(k)}$  be the  $k$ 'th derivative of  $P$ . If

$$T \in \overline{R(\delta_A)} \cap \{A\}',$$

then there exists a sequence  $(X_n)$  in  $\mathcal{L}(\mathcal{H})$  such that

$$AX_n - X_nA \rightarrow T;$$

since  $T \in \{A\}'$  then

$$P^{(k)}(A)X_n - X_nP^{(k)}(A) \rightarrow P^{(k+1)}(A)T.$$

So

$$P(A)X_n - X_nP(A) \rightarrow P^{(1)}(A)T,$$

which shows that

$$P^{(1)}(A)T \in \overline{R(\delta_{P(A)})} \cap \{P(A)\}',$$

that is,  $P^{(1)}(A)T = 0$ . Also we have

$$P^{(1)}(A)X_n - X_nP^{(1)}(A) \rightarrow P^{(2)}(A)T,$$

which gives

$$0 = TP^{(1)}(A)X_nT - TX_nP^{(1)}(A)T \rightarrow P^{(2)}(A)T^3,$$

that is,  $P^{(2)}(A)T^3 = 0$ . By repeating the same argument it follows that  $T^k = 0$  for a given integer number  $k$ , so  $T$  is nilpotent. In particular, every normal operator in  $\overline{R(\delta_A)} \cap \{A\}'$  vanishes. □

**Corollary 3.** Let  $A \in \mathcal{L}(\mathcal{H})$ . If  $P(A)$  is normal, isometric or co-isometric ( $AA^* = I$  or  $A^*A = I$ ) for some polynomial  $P$ , then  $\overline{R(\delta_A)} \cap \{A\}'$  is nilpotent.

**Proof.** In [1, p. 136–137] Anderson showed that

$$\overline{R(\delta_{P(A)})} \cap \{P(A)\}' = \{0\}.$$

□

**Definition 4.** An operator  $A \in \mathcal{L}(\mathcal{H})$  is called *dominant* if, for all complex  $\lambda$ ,  $\text{range}(A - \lambda) \subseteq \text{range}(A - \lambda)^*$ , or equivalently, if there is a real number  $M_\lambda \geq 1$  such that

$$\|(A - \lambda)^* f\| \leq M_\lambda \|(A - \lambda)f\|$$

for all  $f$  in  $\mathcal{H}$ . If there is a constant  $M$  such that  $M_\lambda \leq M$  for all  $\lambda$ ,  $A$  is called *M-hyponormal*, and if  $M = 1$ ,  $A$  is *hyponormal* (see [4]).

**Theorem 5** [5]. Let  $A \in \mathcal{L}(\mathcal{H})$ , then every compact operator in  $\overline{R(\delta_A)}^w \cap \{A\}'$  is quasinilpotent.

**Theorem 6.** If  $B \in \overline{R(\delta_A)}^w \cap \{A\}'$  and  $f(B)$  is compact, where  $f$  is an analytic function on an open set containing  $\sigma(A)$ , then

$$\sigma(B) \subset \{z: zf(z) = 0\}.$$

**Proof.** If  $B \in \overline{R(\delta_A)}^w \cap \{A\}'$ , then

$$AX_\alpha - X_\alpha A \xrightarrow{w} B;$$

since  $f(B) \in \{A\}'$  we have

$$AX_\alpha f(B) - X_\alpha A f(B) \xrightarrow{w} Bf(B),$$

hence

$$AX_\alpha f(B) - X_\alpha f(B)A \xrightarrow{w} Bf(B),$$

that is,

$$Bf(B) \in \overline{R(\delta_A)}^w \cap \{A\}'.$$

Since  $Bf(B)$  is compact, then  $\sigma(Bf(B)) = g(\sigma(B)) = 0$  by Theorem 5, where  $g(z) = zf(z)$ . In particular, if  $P(B)$  is compact for some polynomial  $P$ , then

$$\sigma(B) \subset \{z: zP(z) = 0\}.$$

□

**Theorem 7.** *Let  $A$  or  $A^*$  be a dominant operator.*

*If  $B \in \overline{R(\delta_A)}^w \cap \{A^*\}'$ , then*

$$\{\lambda \in \sigma_p(B^*): \dim \ker(B^* - \bar{\lambda}) < \infty\} \subset \{0\}$$

or,

$$\{\lambda \in \sigma_p(B): \dim \ker(B - \lambda) < \infty\} \subset \{0\},$$

where  $\sigma_p(A)$  is the point spectrum of  $A$ .

*Proof.* Suppose that  $A$  is dominant and  $B \in \overline{R(\delta_A)}^w \cap \{A^*\}'$ , then

$$B^* \in \overline{R(\delta_{A^*})}^w \cap \{A\}'.$$

Let  $\lambda \in \sigma_p(B^*)$  be such that  $E = \ker(B^* - \lambda)$  is finite dimensional.

The subspace  $E$  is invariant under  $B^*$  and  $A$ . It is easy to verify that  $A|_E$  is dominant, hence  $A|_E$  is normal and so  $E$  reduces  $A$  (see [4]).

Let  $H = E \oplus E^\perp$ , then we can write

$$A = \begin{pmatrix} C & 0 \\ 0 & * \end{pmatrix}, \quad B^* = \begin{pmatrix} \lambda & * \\ 0 & * \end{pmatrix}.$$

Since  $B^* \in \overline{R(\delta_{A^*})}^w$ , then  $\lambda I_E \in R(\delta_{C^*})$ , and this necessarily implies  $\lambda = 0$ . □

By the same arguments as in the above proof we achieve the proof of the present theorem.

**Corollary 8.** *If  $A$  or  $A^*$  is a dominant operator, then every compact operator in  $\overline{R(\delta_A)}^w \cap \{A^*\}'$  is quasinilpotent.*

*Proof.* Suppose that  $B \in \overline{R(\delta_A)}^w \cap \{A^*\}'$  with  $B$  compact and  $\lambda \in \sigma(B) \setminus \{0\}$ , then  $\lambda \in \sigma_p(B)$  with  $\dim \ker(B - \lambda) < \infty$  and  $\bar{\lambda} \in \sigma_p(B^*)$  with  $\dim \ker(B^* - \bar{\lambda}) < \infty$ . It follows from Theorem 7 that  $B$  is quasinilpotent. □

### References

- [1] *J.H.Anderson*: On normal derivations. Proc. Amer. Math. Soc. 38 (1973), 135–140..
- [2] *J.H.Anderson*: Derivation ranges and the identity. Bull. Amer. Math. Soc. 79 (1973), 705–708..
- [3] *D.C.Kleïnecke*: On operator commutators. Proc. Amer. Math. Soc. 8 (1957), 535–536..
- [4] *J.G.Stampfli, B.L.Wadhwa*: On dominant operators. Monatsh. Math. 84 (1977), 143–153.
- [5] *R.E.Weber*: Derivations and the trace class operators. Proc. Amer. Math. Soc. 73 (1979), 79–82.

*Author's address:* Université de Tebessa, Département des sciences exactes, 12000 Tebessa, Algeria.