

Jiří Močkoř

Construction of po -groups with quasi-divisors theory

Czechoslovak Mathematical Journal, Vol. 50 (2000), No. 1, 197–207

Persistent URL: <http://dml.cz/dmlcz/127561>

Terms of use:

© Institute of Mathematics AS CR, 2000

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

CONSTRUCTION OF po -GROUPS WITH QUASI-DIVISORS THEORY

JIŘÍ MOČKOŘ, Ostrava

(Received September 30, 1997)

Abstract. A method is presented making it possible to construct po -groups with a strong theory of quasi-divisors of finite character and with some prescribed properties as subgroups of restricted Hahn groups $H(\Delta, \mathbb{Z})$, where Δ are finitely atomic root systems. Some examples of these constructions are presented.

MSC 2000: 13F05, 06F15

Keywords: quasi-divisor theory, divisor class group

1. INTRODUCTION

In the algebraic number theory the notion of a *theory of divisors* was introduced by Borevic and Shafarevic [4] as a map h from the group of divisibility G of an integral domain A into a free abelian group $\mathbb{Z}^{(P)}$ (considered as an l -group with pointwise ordering) satisfying some conditions. It is wellknown (see [4]) that a divisibility group of a domain A admits a theory of divisors if and only if A is a Krull domain.

L. Skula [21] introduced a notion of a theory of divisors for a partly ordered group (po -group) (or, equivalently, for a semigroup with a cancellation law) as a very natural generalization of a theory of divisors for rings, and he derived an extensive theory of these po -groups.

A step towards a further generalization was done by K.E. Aubert in [2], where for the first time the notion of a quasi-divisors theory was introduced. Recall that a directed po -group (G, \cdot) has a *theory of quasi-divisors* if there exists an l -group (Γ, \cdot) and a map $h: G \longrightarrow \Gamma$ such that

- (i) h is an order isomorphism from G into Γ ;
- (ii) $(\forall \alpha \in \Gamma_+)(\exists g_1, \dots, g_n \in G_+) \alpha = h(g_1) \wedge \dots \wedge h(g_n)$.

Supported in part by the Grant Agency of Czech Republic (201/97/0433).

The principal tool for the investigation of these properties in po -groups seems to be the notion of an r -ideal. We recall here that by an r -system of ideals in a directed po -group G we mean a map $X \mapsto X_r$ (X_r is called an r -ideal) from the set of all lower bounded subsets X of G into the power set of G which satisfies the following conditions:

- (1) $X \subseteq X_r$;
- (2) $X \subseteq Y_r \implies X_r \subseteq Y_r$;
- (3) $\{a\}_r = a \cdot G^+ = (a)$ for all $a \in G$;
- (4) $a \cdot X_r = (a \cdot X)_r$ for all $a \in G$.

One of the first characterizations of po -groups with a theory of quasi-divisors was done by P. Jaffard [11]. He proved that a directed po -group G has a theory of quasi-divisors if and only if the semigroup $(\mathcal{I}_t^{(f)}, \times)$ of finitely generated t -ideals is a group, i.e. if and only if G is a t -Prüfer group. (For a comprehensive description see e.g. [2].)

In [14] we introduced a stronger version of po -groups with a theory of quasi-divisors. Recall that a theory of quasi-divisors $h: G \longrightarrow \Gamma$ is called a *strong theory of quasi-divisors*, if

$$(\forall \alpha, \beta \in \Gamma_+)(\exists \gamma \in \Gamma_+) \alpha \cdot \gamma \in h(G), \beta \wedge \gamma = 1.$$

It may be proved that any strong theory of quasi-divisors is a theory of quasi-divisors as well.

Moreover, in a classical divisor theory of po -groups an important role is played by a divisor class group. This notion was introduced by L. Skula [21] as a natural generalization of a class group known from the theory of Krull domains. This notion can be defined naturally for any o -isomorphism $h: G \longrightarrow \Gamma$ of a po -group G into another po -group Γ . Such a definition was introduced in [15] and let us recall that a *divisor class group* \mathcal{C}_h of h is then the abstract group $\Gamma/h(G)$. The canonical map $\Gamma \longrightarrow \mathcal{C}_h$ is then denoted by φ_h .

It was again L. Skula who showed that \mathcal{C}_h and φ_h have great importance when deciding whether or not h is a theory of divisors. We proved (see [19], [16]) that the divisor class group is of the same importance also for po -groups with a theory of quasi-divisors as it is for groups with the classical divisors theory. Namely, we proved that by using some properties of \mathcal{C}_h it is possible to characterize po -groups G with the strong theory of quasi-divisors of a finite character (see [16; Theorem 3.3], [19; Theorem 2.1]).

In this paper we present a general method which enables us to construct examples of po -groups with a strong theory of quasi-divisors of a finite character with some prescribed properties. Using this method we present several examples of po -groups with a quasi-divisors theory with some prescribed properties.

2. EXAMPLES GENERATING

In [16]; Theorem 3.3, we proved the following theorem characterizing po -groups with a theory of quasi-divisors with a finitely atomic value group Γ . Recall that an l -group Γ is *finitely atomic*, if for any element $\alpha \in \Gamma$, $\alpha > 1$, the set of all atoms $\sigma \in \Gamma_+$ such that $\sigma \leq \alpha$ is nonempty and finite. A trivial example of a finitely atomic l -group is the group $\mathbb{Z}^{(P)}$.

Theorem 1 ([16]; 3.3). *Let h be an o -isomorphism from a directed po -group G into an l -group Γ , let \mathcal{C}_h be a divisor class group of h and let $\varphi: \Gamma \longrightarrow \mathcal{C}_h$ be a canonical map. Let us consider the following statements:*

- (1) *h is a strong theory of quasi-divisors.*
- (2) *If $\alpha_1, \dots, \alpha_n$ are elements of Γ such that $\alpha_i > 1$ for all i , then $\varphi(\Gamma_+ \setminus \{\alpha_1, \dots, \alpha_n\}_t) = \mathcal{C}_h$, where $\{\alpha_1, \dots, \alpha_n\}_t = \{\alpha \in \Gamma: \exists 1 \leq i \leq n, \alpha \geq \alpha_i\}$.*
- (3) *If $\alpha_1, \dots, \alpha_n$ are atoms in Γ_+ , then $\varphi(\Gamma_+ \setminus \{\alpha_1, \dots, \alpha_n\}_t) = \mathcal{C}_h$.*

Then (1) \implies (2) \implies (3). If Γ is finitely atomic, then all the statements are equivalent.

A method for constructing examples of po -groups with a strong theory of quasi-divisors of a finite character that we will present here is based on an application of Theorem 1 for a special l -group, the restricted Hahn group $H(\Delta, \mathbb{Z})$.

Recall that if Δ is a *root system* (i.e. (Δ, \leq) is a partly ordered set for which $\{\alpha \in \Delta: \alpha \geq \gamma\}$ is totally ordered for any $\gamma \in \Delta$), then the *restricted Hahn group* $H(\Delta, \mathbb{Z})$ on Δ is the group $\mathbb{Z}^{(\Delta)}$ ordered in the following way:

$$a \in H(\Delta, \mathbb{Z}), a \geq 0 \Leftrightarrow a_\alpha > 0 \text{ for all } \alpha \in \text{ms}(a),$$

where $\text{ms}(a)$ is the *maximal support* of a , i.e. the set of all maximal elements in $\text{supp}(a) = \{\alpha \in \Delta: a_\alpha \neq 0\}$. Then $H(\Delta, \mathbb{Z})$ is an l -group (see e.g. [1]).

Now, let Δ_0 be the set of all minimal elements of Δ . We say that Δ is *atomic* if for any element $\alpha \in \Delta$ there exists $\beta \in \Delta_0$ such that $\alpha \geq \beta$. Moreover, we say that Δ is *finitely atomic* if for any $\alpha \in \Delta$, the set $\{\sigma \in \Delta_0: \sigma \leq \alpha\}$ is nonempty and finite. Finally, let $\alpha \in \Delta$. Then by a^α we denote the element of $H(\Delta, \mathbb{Z})$ such that

$$a_\beta^\alpha = \begin{cases} 1 & \text{if } \beta = \alpha, \\ 0 & \text{otherwise.} \end{cases}$$

In the following lemma we recall some properties of $H(\Delta, \mathbb{Z})$ which can be of interest for our examples of groups with a strong theory of quasi-divisors.

Lemma 2 ([16]; 3.4). *Let Δ be a root system.*

- (1) *Let $\alpha \in \Delta_0$, $b \in H(\Delta, \mathbb{Z})_+$. Then $b \geq a^\alpha$ if and only if there exists $\beta \in ms(b)$ such that $\beta \geq \alpha$.*
- (2) *If Δ is atomic, then $a \in H(\Delta, \mathbb{Z})$ is an atom if and only if $a = a^\alpha$ for some $\alpha \in \Delta_0$.*
- (3) *If Δ is finitely atomic, then $H(\Delta, \mathbb{Z})$ is finitely atomic.*

Our examples of groups with a strong theory of quasi-divisors will be based on the application of Theorem 1 and Lemma 2 to a group $H(\Delta, \mathbb{Z})$. Hence, we will investigate homomorphisms φ of $H(\Delta, \mathbb{Z})$ into an abelian group \mathcal{C} . We will be interested in homomorphisms $\bar{\varphi}$ which are determined by maps $\varphi: \Delta \rightarrow \mathcal{C}$ in the following way.

We say that a homomorphism $\psi: H(\Delta, \mathbb{Z}) \rightarrow \mathcal{C}$ is determined by a map $\varphi: \Delta \rightarrow \mathcal{C}$ if for any $a \in H(\Delta, \mathbb{Z})$ we have

$$\psi(a) = \sum_{\alpha \in \Delta} \varphi(\alpha) a_\alpha.$$

In this case the following diagram commutes:

$$\begin{array}{ccc} \Delta & \xrightarrow{\varphi} & \mathcal{C} \\ \text{nat} \downarrow & & \parallel \\ H(\Delta, \mathbb{Z}) & \xrightarrow{\psi} & \mathcal{C} \end{array}$$

where $\text{nat}(\alpha) = a^\alpha$. The homomorphism ψ will be then denoted by $\bar{\varphi}$.

Corollary (Examples generating method). *Let (Δ, \leq) be a finitely atomic root system, \mathcal{C} an abelian group and let $\varphi: \Delta \rightarrow \mathcal{C}$ be a map such that for any finite set $\{\alpha_1, \dots, \alpha_n\}$ of atoms in Δ the set $\varphi(\Delta \setminus \{\alpha_1, \dots, \alpha_n\}_t)$ is a semigroup generator of \mathcal{C} , where $\{\alpha_1, \dots, \alpha_n\}_t = \{\alpha \in \Delta: \exists i, \alpha \geq \alpha_i\}$. Then in the diagram*

$$\text{Ker } \bar{\varphi} \xrightarrow{h} H(\Delta, \mathbb{Z}) \xrightarrow{\bar{\varphi}} \mathcal{C},$$

the inclusion map h is a strong theory of quasi-divisors of a finite character and \mathcal{C} is a divisor class group of h .

P r o o f. Since Δ is finitely atomic, according to Lemma 2, the l -group $H(\Delta, \mathbb{Z})$ is finitely atomic. Let $\{a_1, \dots, a_n\}$ be a finite set of atoms in $H(\Delta, \mathbb{Z})$. Then according to Lemma 2, for any $i, 1 \leq i \leq n$, there exists an atom $\alpha_i \in \Delta$ such that $a_i = a^{\alpha_i}$. Let $\mathbf{a} \in \mathcal{C}$. Since $\varphi(\Delta \setminus \{\alpha_1, \dots, \alpha_n\}_t)$ is a semigroup generator of \mathcal{C} , there exist $\beta_1, \dots, \beta_m \in \Delta \setminus \{\alpha_1, \dots, \alpha_n\}_t$ and natural numbers k_1, \dots, k_m such that $\mathbf{a} =$

$\sum_{j=1}^m \varphi(\beta_j)k_j$. If $a^{\beta_j}k_j \geq a_i = a^{\alpha_i}$ for some i , then we have $\beta_j \geq \alpha_i$ according to Lemma 2, a contradiction. Hence $b = \sum_{j=1}^m a^{\beta_j}k_j \in H(\Delta, \mathbb{Z})_+ \setminus \{a_1, \dots, a_n\}_t$ and

$$\bar{\varphi}(b) = \sum_{\alpha \in \Delta} \varphi(\alpha) \cdot b_\alpha = \sum_{j=1}^m \varphi(\beta_j)k_j = \mathbf{a}.$$

We will show that $\text{Ker } \bar{\varphi} = G$ is a directed po -group. Let $a \in G$. We put

$$b(\alpha) = \begin{cases} a(\alpha) & \text{if } \alpha \in \text{ms}(a), a(\alpha) > 0, \\ 0 & \text{if } \alpha \in \text{ms}(a), a(\alpha) < 0, \\ |a(\alpha)| & \text{if } \alpha \in \text{supp}(a) \setminus \text{ms}(a), \\ 0 & \text{otherwise.} \end{cases}$$

Then $b \geq a, 0$ in $H(\Delta, \mathbb{Z})$, since $(b - a)(\alpha) > 0$ for all $\alpha \in \text{ms}(b - a)$. Let $\mathbf{a} = \sum_{\alpha \in \text{supp}(b)} b(\alpha) \cdot \varphi(\alpha)$ and let

$$\Phi^b = \{\alpha \in \Delta : \alpha \text{ is an atom in } \Delta, \alpha \leq \beta \text{ for some } \beta \in \text{supp}(b)\}.$$

Then Φ^b is a finite set in Δ and $\varphi(\Delta \setminus \Phi_t^b)$ is a semigroup generator of \mathcal{C} . Hence, there exist $\beta_1, \dots, \beta_n \in \Delta \setminus \Phi_t^b$ and positive numbers $c_1, \dots, c_n \in \mathbb{Z}$ such that $-\mathbf{a} = \varphi(\beta_1)c_1 + \dots + \varphi(\beta_n)c_n$. Then $\beta_i \notin \text{supp}(b)$ for any i and we put

$$c(\alpha) = \begin{cases} b(\alpha) & \text{if } \alpha \in \text{supp}(b), \\ c_i & \text{if } \alpha = \beta_i, i = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

Then $c > 0$ in $H(\Delta, \mathbb{Z})$ and $c \geq b \geq a$. Moreover,

$$\sum_{\alpha \in \Delta} c(\alpha)\varphi(\alpha) = \sum_{\alpha \in \text{supp}(b)} b(\alpha)\varphi(\alpha) + \sum_{i=1}^n c_i\varphi(\beta_i) = 0.$$

Hence, $c \in G$ and since G is a convex subgroup, we obtain that $b \in G$ and G is directed.

Now, according to [17]; 2.2, Γ satisfies Conrad's (F)-condition and it follows that h is of finite character. The result then follows from Theorem 1. \square

Corollary (Skula, L. [21]). *Let G be a p_0 -group and let $h: G \longrightarrow \mathbb{Z}^{(P)}$ be an o -isomorphism into. Then the following conditions are equivalent.*

- (1) h is a strong theory of divisors.
- (2) For $p_1, \dots, p_n \in P$ ($n \geq 1$), the set $\varphi_h(P \setminus \{p_1, \dots, p_n\})$ is a semigroup generator of a divisor class group $\mathbb{Z}^{(P)}/h(G)$.

We investigate first some additional properties of the inclusion $h: G = \text{Ker } \bar{\varphi} \longrightarrow \Gamma = H(\Delta, \mathbb{Z})$. In what follows we will always assume that Δ is a finitely atomic root system.

Let Δ_0 be the set of all atoms in Δ . Then Δ_0 is the maximal disjoint set in Δ and it follows that $\{\text{nat}(\alpha): \alpha \in \Delta_0\}$ is a base of Γ . Moreover, the set of polars of $\text{nat}(\alpha), \alpha \in \Delta_0$, then defines an l -realization of Γ , i.e. if we put

$$\Delta_\alpha^+ = (\text{nat}(\alpha)')_+ = \{g \in \Gamma_+: g \wedge \text{nat}(\alpha) = 0\}$$

for $\alpha \in \Delta_0$ then since Γ satisfies the Conrad (F)-condition (see [17]; 2,2), the set

$$W = \{w_\alpha: \Gamma \xrightarrow{w_\alpha} \Gamma/\Delta_\alpha \text{ is a canonical } l\text{-homomorphism, } \alpha \in \Delta_0\}$$

is a defining family of a finite character of Γ (see [5]; p. 3.29). Moreover, for any $\alpha \in \Delta_0$ we have

$$\begin{aligned} \Delta_\alpha^+ &= \{g \in \Gamma_+: \text{supp}(g) \cap (\alpha)_t = \emptyset\} \\ &= \{g \in \Gamma_+: g(\beta) = 0 \text{ for all } \beta \in \Delta, \beta \geq \alpha\} \end{aligned}$$

according to [16]; 3.4.

In the following proposition we present some properties of this defining family W .

Proposition 3. *Let W be a defining family of Γ introduced above.*

- (1) W is an independent defining family if and only if for all $\alpha, \beta \in \Delta_0, \alpha \neq \beta, (\alpha)_t \cap (\beta)_t = \emptyset$ holds.
- (2) If $\alpha \in \Delta_0$ is such that $\alpha < \beta$ for some $\beta \in \Delta$, then Γ/Δ_α is not isomorphic to \mathbb{Z} .

Proof. (1). Let W be independent and let us assume that there are $\alpha \neq \beta$ in W such that $\gamma \geq \alpha, \beta$ for some γ . Let $g = \text{nat}(\gamma) \in \Gamma$. Then $g \notin \Delta_\alpha \cup \Delta_\beta$. If $g \in [\Delta_\alpha, \Delta_\beta]$, then there exist $a \in \Delta_\alpha, b \in \Delta_\beta$ such that $a + b \geq g$. But in this case $a(\gamma) = b(\gamma) = 0$ and it follows that $\gamma \in \text{ms}((a + b) - g)$. Hence, $a + b \not\geq g$, a

contradiction. Conversely, let $(\alpha)_t \cap (\beta)_t = \emptyset$ for all $\alpha \neq \beta \in \Delta_0$. Let $g \in \Gamma_+$ and $\alpha \neq \beta \in \Delta_0$. We put

$$a(\gamma) = \begin{cases} 0 & \gamma \geq \alpha, \\ g(\gamma) & \gamma \not\geq \alpha, \end{cases}$$

$$b(\gamma) = \begin{cases} 0 & \gamma \geq \beta, \\ g(\gamma) & \gamma \not\geq \beta. \end{cases}$$

Then $a \in \Delta_\alpha, b \in \Delta_\beta$ and it follows from $(\alpha)_t \cap (\beta)_t = \emptyset$ that $\gamma \in \text{ms}((a+b) - g)$ iff $\gamma \in \text{ms}(g)$. Hence, $a+b > g$ and $[\Delta_\alpha, \Delta_\beta] = G$.

(2) Let $a, b, c_n; n = 1, 2, \dots$ be defined such that

$$a(\gamma) = 2 \text{ if } \gamma = \beta,$$

$$b(\gamma) = 1 \text{ if } \gamma = \beta,$$

$$c_n(\gamma) = 2 \text{ if } \gamma = \beta,$$

$$c_n(\gamma) = -n, \text{ if } \gamma = \alpha,$$

$$a(\gamma) = b(\gamma) = c_n(\gamma) = 0 \text{ otherwise.}$$

Then

$$a + \Delta_\alpha > c_1 + \Delta_\alpha > c_2 + \Delta_\alpha > \dots > b + \Delta_\alpha > \Delta_\alpha.$$

Hence, Γ/Δ_α cannot be order isomorphic to \mathbb{Z} . □

Example 1. For any infinite cardinal number \mathbf{a} there exists a po -group G with a strong theory of quasi-divisors of a finite character such that G_+ has at least \mathbf{a} maximal prime t -ideals.

In fact, let Δ be a set with cardinality \mathbf{a} . Let Δ be considered to be an antichain. Let $\Delta = \Delta_1 \cup \Delta_2$ be a partition such that any Δ_i is infinite and let us define a map $\varphi: \Delta \rightarrow \mathbf{Z}$ such that

$$\varphi(\delta) = \begin{cases} 1 & \text{if } \delta \in \Delta_1, \\ -1 & \text{if } \delta \in \Delta_2. \end{cases}$$

Hence, according to the first Corollary, $G = \ker(\bar{\varphi}) \xrightarrow{h} \Gamma = H(\Delta, \mathbb{Z})$ is a po -group with a strong theory of quasi-divisors of a finite character. Moreover, since Δ is a set of all atoms in (Δ, \leq) , then for any $\delta \in \Delta$, $\Delta_\delta = \{g \in \Gamma: g_\beta = 0, \forall \beta \geq \delta\}$ is a minimal prime l -ideal in Γ . Hence, according to [14]; Prop. 2.4, $P_\delta = \Gamma_+ \setminus \Delta_\delta$ is a maximal prime t -ideal in Γ . Since an embedding h is a (t, t) -morphism, it follows that $Q_\delta = h^{-1}(P_\delta)$ is a prime t -ideal in G as well. Moreover, Q_δ is maximal. In fact, if Q is a prime t -ideal in G such that $Q_\delta \subset Q$ then since G is a t -Prüfer po -group (see [17];

Theorem 2.1, for example), a canonical map $w_Q: G \longrightarrow G/H$ is a t -valuation, where H is the quotient group of a semigroup $G_+ \setminus Q$. Let \hat{w}_Q be an extension of w_Q onto a t -valuation of Γ . The existence of this extension follows from the universal property of Γ (see [2], e.g.). Then $\Gamma_+ \setminus \ker(\hat{w}_Q)$ is a prime t -ideal strictly containing P_δ , a contradiction. Hence, $\{Q_\delta: \delta \in \Delta\}$ is a set of maximal prime t -ideals of G .

Our next two examples will concern G -dense l -ideals of Γ . Recall that an l -ideal Δ of Γ is called G -dense (with respect to an o -isomorphism h from G into Γ) if for any $\alpha \in \Delta$ there exists $g \in G$ such that $\alpha \leq h(g) \in \Delta$. In [18]; 3.3, it was proved that there exists a bijection between the set of o -ideals of a (t -closed) po -group G and the set of G -dense l -ideals of its Lorenzen l -group $\Lambda_t(G)$. In the first example we show that even in the case that an inclusion $G \longrightarrow \Lambda_t(G)$ is a strong theory of quasi-divisors of a finite character, in $\Lambda_t(G)$ there exist (in general) l -ideals which are not G -dense.

Example 2. There exists a po -group G with a strong theory of quasi-divisors of a finite character $h: G \longrightarrow \Gamma$ such that in Γ there exists an l -ideal which is not G -dense.

In fact, let A be an infinite finitely atomic root system as in Fig. 1. We define a map $\varphi: A \longrightarrow \mathbb{Z}$ such that

$$\varphi(\alpha_i) = (-1)^i; \quad \varphi(\alpha_{ij}) = 0.$$




Fig. 1

Then for any finite set F of atoms from A , $\varphi(A \setminus F_t)$ is a semigroup generator of \mathbb{Z} and according to Corollary, $G = \text{Ker } \bar{\varphi} \xrightarrow{h=\text{id}} \Gamma = H(A, \mathbb{Z})$ is a strong theory of quasi-divisors of a finite character with \mathbb{Z} as a divisor class group.

We set

$$\Delta = \bigcap_{i \text{ odd}} \Delta_{\alpha_i} = \{g \in \Gamma: g(\beta) = 0, \text{ for all } \beta \geq \alpha_i, i \text{ odd}\}$$

where Δ_{α_i} is a polar of $\text{nat}(\alpha_i)$. Then Δ is an l -ideal of Γ . Let $a = \text{nat}(\alpha_2) \in \Gamma$. Then $a \in \Delta$. Now, if Δ is G -dense, there exists $g \in G$ such that $g \geq a$, $g \in \Delta$. Hence, $g(\alpha_i) = 0 = g(\alpha_{i,i+1})$ for all i odd. Since $g \in G$ we then have $0 = \sum_i g(\alpha_i)(-1)^i = \sum_{i \text{ even}} g(\alpha_i)$. If $g(\alpha_2) < 1$, then $\alpha_2 \in \text{ms}(g - a)$ and $(g - a)(\alpha_2) < 0$, a contradiction. Hence, $g(\alpha_2) \geq 1$ and there exists another i such that $g(\alpha_i) < 0$. But then $i \in \text{ms}(g)$ and $g \not\geq 0$, a contradiction. Therefore, Δ is not G -dense.

In [18]; 3.5, it was proved that any intersection of finitely many prime l -ideals of Γ is G -dense. In Example 2 it was shown that there exists an intersection of infinitely many prime l -ideals from a base of Γ which is not G -dense. In the next example we show that this is not a typical case.

Example 3. There exists a po -group G with a strong theory of quasi-divisors of a finite character $h: G \rightarrow \Gamma$ such that any intersection of infinitely many prime l -ideals from a base of Γ is G -dense.

In fact, let A be an infinite finitely atomic root system as in Fig. 2. We define a map $\varphi: A \rightarrow \mathbb{Z}$ such that

$$\varphi(\alpha_i) = \varphi(\bar{\alpha}_i) = (-1)^i; \quad i = 1, 2, \dots$$

Then for any finite set F of atoms from A , $\varphi(A \setminus F_t)$ is a semigroup generator of \mathbb{Z} and according to Corollary, $G = \text{Ker } \bar{\varphi} \xrightarrow{h=\text{id}} \Gamma = H(A, \mathbb{Z})$ is a strong theory of quasi-divisors of a finite character.




Fig. 2

Then $\{\Delta_{\alpha_i} : i = 1, 2, \dots\}$ is a base of Γ , where $\Delta_{\alpha} = \{g \in \Gamma : g(\beta) = 0, \beta \in A, \beta \geq \alpha\}$. Let $B \subseteq \{\alpha_1, \dots\}$ be an arbitrary infinite set and let $\Delta = \bigcap_{\beta \in B} \Delta_{\beta}$. Let $a \in \Delta_+$ be arbitrary, i.e. $a(\alpha) = a(\bar{\alpha}) = 0$ for any $\alpha \in B$. We define an element $g \in \Gamma$ such

that

$$g(\bar{\alpha}) = \begin{cases} 0 & \text{if } \alpha \in B, \\ a(\bar{\alpha}) + 1 & \text{if } a(\bar{\alpha}) \neq 0, \\ 1 & \text{if } a(\bar{\alpha}) = 0, a(\alpha) \neq 0, \\ 0 & \text{if } a(\bar{\alpha}) = a(\alpha) = 0, \end{cases}$$

$$g(\alpha) = -g(\bar{\alpha}).$$

Then $\sum_i (g(\alpha_i) + g(\bar{\alpha}_i))(-1)^i = 0$ and $g \in G$. Moreover, $g \in \Delta$ and $g \geq a$. Hence, Δ is G -dense.

References

- [1] *Anderson, M., Feil, T.*: Lattice-ordered Groups. D. Reidl Publ. Co., Dordrecht, Tokyo, 1988.
- [2] *Aubert, K.E.*: Divisors of finite character. *Annali di matem. pura ed appl.* 33 (1983), 327–361.
- [3] *Aubert, K.E.*: Localizations dans les systèmes d'idéaux. *C.R.Acad. Sci. Paris* 272 (1971), 465–468.
- [4] *Borevich, Z. I. and Shafarevich, I. R.*: Number Theory. Academic Press, New York, 1966.
- [5] *Conrad, P.*: Lattice Ordered Groups. Tulane University, 1970.
- [6] *Chouinard, L. G.*: Krull semigroups and divisor class group. *Canad. J. Math.* 33 (1981), 1459–1468.
- [7] *Geroldinger, A., Močkoř, J.*: Quasi-divisor theories and generalizations of Krull domains. *J. Pure Appl. Algebra* 102 (1995), 289–311.
- [8] *Gilmer, R.*: Multiplicative Ideal Theory. M. Dekker, Inc., New York, 1972.
- [9] *Griffin, M.*: Rings of Krull type. *J. Reine Angew. Math.* 229 (1968), 1–27.
- [10] *Griffin, M.*: Some results on v -multiplication rings. *Canad. J. Math.* 19 (1967), 710–722.
- [11] *Jaffard, P.*: Les systèmes d'idéaux. Dunod, Paris, 1960.
- [12] *Močkoř, J.*: Groups of Divisibility. D. Reidl Publ. Co., Dordrecht, 1983.
- [13] *Močkoř, J., Alajbegovic, J.*: Approximation Theorems in Commutative Algebra. Kluwer Academic publ., Dordrecht, 1992.
- [14] *Močkoř, J., Kontolatou, A.*: Groups with quasi-divisor theory. *Comm. Math. Univ. St. Pauli, Tokyo* 42 (1993), 23–36.
- [15] *Močkoř, J., Kontolatou, A.*: Divisor class groups of ordered subgroups. *Acta Math. Inform. Univ. Ostraviensis* 1 (1993).
- [16] *Močkoř, J., Kontolatou, A.*: Quasi-divisors theory of partly ordered groups. *Grazer Math. Ber.* 318 (1992), 81–98.
- [17] *Močkoř, J.*: t -Valuation and theory of quasi-divisors. To appear in *J. Pure Appl. Algebra*.
- [18] *Močkoř, J., Kontolatou, A.*: Some remarks on Lorezen r -group of partly ordered group. *Czechoslovak Math. J.* 46(121) (1996), 537–552.
- [19] *Močkoř, J.*: Divisor class group and the theory of quasi-divisors. To appear.
- [20] *Ohm, J.*: Semi-valuations and groups of divisibility. *Canad. J. Math.* 21 (1969), 576–591.
- [21] *Skula, L.*: Divisorentheorie einer Halbgruppe. *Math. Z.* 114 (1970), 113–120.

[22] *Skula, L.*: On c -semigroups. *Acta Arith.* *31* (1976), 247–257.

Author's address: Department of Mathematics, University of Ostrava, CZ-702 00 Ostrava, e-mail: `Jiri.Mockor@osu.cz`.