

Chong-Yun Chao; Shaocen Han  
On the toughness of cycle permutation graphs

*Czechoslovak Mathematical Journal*, Vol. 51 (2001), No. 2, 239–260

Persistent URL: <http://dml.cz/dmlcz/127645>

## Terms of use:

© Institute of Mathematics AS CR, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## ON THE TOUGHNESS OF CYCLE PERMUTATION GRAPHS

CHONG-YUN CHAO, SHAO-CEN HAN, Pittsburgh

(Received January 30, 1998)

*Abstract.* Motivated by the conjectures in [11], we introduce the maximal chains of a cycle permutation graph, and we use the properties of maximal chains to establish the upper bounds for the toughness of cycle permutation graphs. Our results confirm two conjectures in [11].

*Keywords:* cycle permutation graph, toughness, maximal chain

*MSC 2000:* 05C58

## 1. INTRODUCTION

Chartrand and Wilson, in [5], introduced a series of properties for the Petersen graph. They used a conjecture of Tutte to explain why so much attention had been paid to the Petersen graph and its various generalizations. (That is, every known bridgeless 3-regular graph whose edges cannot be colored with three colors contains a subgraph isomorphic to the Petersen graph. Tutte conjectured that this is always the case (see [12]).

Two classes of generalization of Petersen graphs are generalized Petersen graphs and cycle permutation graphs. Let  $n$  and  $k$  be integers with  $n \geq 5$  and  $k \geq 1$ . A generalized Petersen graph,  $G(n, k)$ , is the graph with vertex set  $V(G(n, k)) = \{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n\}$ , and edge set  $E(G(n, k)) = \{[x_i, x_{i+1}], [y_i, y_{i+k}], [x_i, y_i], i = 1, 2, \dots, n$  where the subscripts are taken modulo  $n\}$ . The subgraph induced from  $x_1, x_2, \dots, x_n$  is denoted by  $C_x$  and the subgraph induced from  $y_1, y_2, \dots, y_n$  is denoted by  $C_y$ . When  $n = 5$  and  $k = 2$ ,  $G(5, 2)$  is the Petersen graph.

Let  $\alpha$  be a permutation of the symmetric group,  $S_n$ , acting on the set  $\{1, 2, \dots, n\}$ . A cycle permutation graph  $P_n(\alpha)$  is the graph with  $2n$  vertices,  $V(P_n(\alpha)) = V_1 \cup V_2$  where  $V_i = \{v_{i1}, v_{i2}, \dots, v_{in}\}$  for  $i = 1, 2$ ,  $V_1 \cap V_2 = \emptyset$ , and  $E(P_n(\alpha)) = E_1 \cup E_2 \cup E_{12}$

where  $E_i = \{[v_{ij}, v_{i(j+1)}]\}$  for  $j = 1, 2, \dots, n\}$  for  $i = 1, 2$  and  $E_{1,2} = \{[v_1 v_{2\alpha(t)}]; t = 1, 2, \dots, n\}$ . (See [6], [7].) When  $n = 5$  and  $\alpha = (1)(2453)$ ,  $P_5(\alpha)$  is the Petersen graph.

The toughness  $t(G)$  of a graph  $G$  was defined by Chvátal [6]. If  $G$  is not a complete graph,

$$t(G) = \min_S \left\{ \frac{|S|}{\omega(G-S)} \right\}$$

where  $S$  is taken over all disconnecting subsets of the vertex set of  $G$ ,  $|S|$  is the cardinality of  $S$  and  $\omega(G-S)$  is the number of components in the subgraph induced from  $G-S$ .

Recently, the toughness of graphs has received a lot of attention. Much work has been done concerning the toughness, which is considered to be more sensitive to the structure of the graph than the connectivity of the graph (see [1] and [2]).

In [6], Chvátal first considered the toughness of the cross product of two complete graphs. Guichard, Piazza, and Stueckle, in [10], proved that for  $\alpha \in S_{m+n}$  and  $m \leq n$ , the toughness of a cycle permutation graph is given by

$$t(P_\alpha(K_{m,n})) = \begin{cases} \frac{2m}{n+m-q} & \text{if } q < \frac{n^2+m^2}{n+3m}, \\ \frac{n+m}{n+q} & \text{if } q \geq \frac{n^2+m^2}{n+3m} \end{cases}$$

where  $K_{m,n}$  is the complete bipartite graph of  $mn$  vertices.

Some results and conjectures were given by Piazza, Ringeisen, and Stueckle in [11]; the authors proved that the toughness of  $G(n, k)$  is more than  $n/(n-1)$ , if  $n$  is an positive odd integer with  $n$  and  $k$  being relatively prime, and  $k \notin \{1, n-1\}$ . An upper bound for the toughness of cycle permutation graphs was obtained as follows:

$$t(P_n(\alpha)) \leq \frac{(k+2)}{(k+1)}, \quad \text{if } \alpha(i) = i \text{ for all } 1 \leq i \leq k \leq n-2.$$

Based on the set of permutations which generate all nonisomorphic cycle permutation graphs of  $C_n$ ,  $n \leq 8$  in [11], the authors found that for all  $\sigma \in S_n$ , the toughness of  $P_3(\sigma)$  is equal to  $3/2$  and the toughness of  $P_n(\sigma)$  is less than or equal to  $4/3$  for  $4 \leq n \leq 8$ . Three conjectures were stated as follows.

**Conjecture 1.** For  $n \geq 4$  and  $\alpha \in S_n$ ,  $t(P_n(\alpha)) \leq 4/3$ .

If this upper bound cannot be obtained, perhaps, the following looser upper bound can be obtained.

**Conjecture 2.** For  $n \geq 4$  and  $\alpha \in S_n$ ,  $t(P_n(\alpha)) < 3/2$ .

Since  $G(5, 2)$  and  $G(9, 2)$  have their toughness equal to  $4/3$ , could such a class be the generalized Petersen graphs when  $n \equiv 1 \pmod{4}$  and  $k = 2$ ?

**Conjecture 3.** *If  $n \geq 5$  and  $n \equiv 1 \pmod{4}$ , then  $t(G(n, 2)) = 4/3$ .*

In [3] and [8], the authors proved that the upper bound for toughness of generalized Petersen graph is  $4/3$ . Here, we shall study the structure of cycle permutation graphs with some maximal chains and establish the upper bound for the toughness of cycle permutation graphs. Some of these results confirm Conjectures 1 and 2. Throughout this paper, all integers and subscripts are taken modulo  $n$ .

## 2. MAXIMAL CHAINS

Let  $P_n(\alpha)$  be a cycle permutation graph consisting of two  $n$ -cycles  $C_n$  and  $C'_n$  with a connecting set of edges, i.e.,  $V(P_n(\alpha)) = V(C_n) \cup V(C'_n)$  where  $V(C_n) = \{1, 2, \dots, n\}$ ,  $V(C'_n) = \{y_1, y_2, \dots, y_n\}$  such that  $V(C_n) \cap V(C'_n) = \emptyset$ , and  $E(P_n(\alpha)) = E(C_n) \cup E(C'_n) \cup E_{1,2}$  where  $E(C_n) = \{[i, i+1] \text{ for } i = 1, 2, \dots, n\}$ ,  $E(C'_n) = \{[y_i, y_{i+1}] \text{ for } i = 1, 2, \dots, n\}$  and  $E_{1,2} = \{[i, y_{\alpha(i)}] \text{ for } i = 1, 2, \dots, n\}$ .

Let  $B$  be a nonempty (proper) subset of  $V(C_n)$ . On  $C'_n$ , a chain of edges  $[y_i, y_{i+1}][y_{i+1}, y_{i+2}] \dots [y_{t-1}, y_t]$  is said to be related to  $B$ , if  $\alpha^{-1}(i), \alpha^{-1}(i+1), \dots, \alpha^{-1}(t)$  belong to  $B$ . For simplicity, this chain of edges on  $C'_n$  will be written as  $y_i y_{i+1} \dots y_t$ . A chain of edges,  $y_i y_{i+1} \dots y_t$  is said to be maximal, if  $\alpha^{-1}(i-1) \notin B$  and  $\alpha^{-1}(t+1) \notin B$ . A maximal chain related to  $B$  will be denoted by  $M(B)$ . Similarly, replacing  $B$  by  $C_n - B$ , we may define chains related to  $C_n - B$ , and maximal chains related to  $C_n - B$ . Also, a maximal chain related to  $C_n - B$  will be denoted by  $M(C_n - B)$ . Two chains on  $C'_n$ ,  $y_i y_{i+1} \dots y_t$  and  $y_j y_{j+1} \dots y_s$  are said to be related, denoted by  $y_i y_{i+1} \dots y_t \sim y_j y_{j+1} \dots y_s$  or  $y_j y_{j+1} \dots y_s \sim y_i y_{i+1} \dots y_t$ , if  $[y_t, y_j] \in E(C'_n)$  or  $[y_s, y_i] \in E(C'_n)$ .

For a nonempty independent subset  $B$  of  $V(C_n)$ ,  $C'_n$  is partitioned into disjoint maximal chains  $M_1(B), M_2(B), \dots, M_k(B)$  related to  $B$ , and disjoint maximal chains  $M_1(C_n - B), M_2(C_n - B), \dots, M_k(C_n - B)$  related to  $C_n - B$ . This partition of maximal chains will be denoted by  $p(n, \alpha, B)$ . We note that since  $C'_n$  is a cycle, the number of maximal chains related to  $B$  is equal to the number of maximal chains related to  $(C_n - B)$ .

**Example 1.** Let  $n = 12$ , and

$$\alpha = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 \\ 3 & 2 & 4 & 5 & 7 & 9 & 6 & 12 & 1 & 8 & 10 & 11 \end{pmatrix}.$$

In  $P_{12}(\alpha)$ , let  $B = \{1, 3, 5, 7, 10\}$ . Then we have  $\alpha^{-1}(3) = 1$ ,  $\alpha^{-1}(4) = 3$ ,  $\alpha^{-1}(7) = 5$ ,  $\alpha^{-1}(6) = 7$ ,  $\alpha^{-1}(8) = 10$ ,  $M_1(B) = y_3 y_4$ ,  $M_2(B) = y_6 y_7 y_8$ ,  $M_1(C_{12} -$

$B) = y_5$ ,  $M_2(C_{12} - B) = y_9y_{10}y_{11}y_{12}y_1y_2$ , and  $p(12, \alpha, B)$  is  $M_1(B) \smile (C_{12} - B) \smile M_2(B) \smile M_2(C_{12} - B) \smile M_1(B)$ . Also,  $y_6y_7$  is a chain related to  $B$ , but not a maximal chain related to  $B$ . Similarly,  $y_{12}y_1y_2$  is a chain related to  $(C_{12} - B)$ , but not a maximal chain related to  $(C_{12} - B)$ .

Each of  $M_1(B)$  and  $M_2(C_{12} - B)$  is said to have an even cardinality (i.e., each contains an even number of vertices), and each of  $M_2(B)$  and  $M_1(C_{12} - B)$  has an odd cardinality.

**Theorem 1.** *Let  $n$  be an integer  $\geq 4$ ,  $\alpha \in S_n$ ,  $P_n(\alpha)$  be a cycle permutation graph with  $2n$  vertices,  $B$  be a nonempty independent subset of vertices of  $C_n$  in  $P_n(\alpha)$ , and  $p(n, \alpha, B)$  be the partition of maximal chains related to  $B$ . Then*

$$(1) \quad t(P_n(\alpha)) \leq \frac{2|B| + n + e_1(B) - e_2(B)}{3|B| + e_1(B)}$$

where  $|B|$  is the cardinality of  $B$ ,  $e_1(B)$  is the number of maximal chains related to  $B$  with odd cardinality, and  $e_2(B)$  is the number of maximal chains related to  $C_n - B$  with odd cardinality.

*Proof.* For some positive integer  $q$ ,  $p(n, \alpha, B)$  is  $M_1(B) \smile M_1(C_n - B) \smile M_2(B) \smile M_2(C_n - B) \smile \dots \smile M_q(B) \smile M_q(C_n - B) \smile M_1(B)$ .  $\square$

We construct a disconnecting subset  $S$  of  $P_n(\alpha)$  as follows:

$$(2) \quad S = B \cup B_y \quad \text{with } B_y = \bigcup_{j=1}^q [K_j(B) \cup K_j(C_n - B)]$$

where for  $M_j(B) = y_{t_j+1}y_{t_j+2} \dots y_{t_j+m_j}$ ,  $1 \leq j \leq q$ ,

$$K_j(B) = \begin{cases} \varnothing & \text{if } m_j = 1 \text{ or } 2, \\ \{y_{t_j+2}, y_{t_j+4}, \dots, y_{t_j+m_j-1}\} & \text{if } m_j \text{ is odd } > 1, \\ \{y_{t_j+2}, y_{t_j+4}, \dots, y_{t_j+m_j-2}\} & \text{if } m_j \text{ is even } > 2. \end{cases}$$

For  $M_j(C_n - B) = y_{s_j+1}y_{s_j+2} \dots y_{s_j+m'_j}$ ,  $1 \leq j \leq q$ ,

$$K_j(C_n - B) = \begin{cases} \{y_{s_j+1}, y_{s_j+3}, \dots, y_{s_j+m'_j-2}, y_{s_j+m'_j}\} & \text{if } m'_j \text{ is odd,} \\ \{y_{s_j+1}, y_{s_j+3}, \dots, y_{s_j+m'_j-1}, y_{s_j+m'_j}\} & \text{if } m'_j \text{ is even.} \end{cases}$$

Thus, we have, for  $j = 1, 2, \dots, q$ ,

$$|M_j(B)| = m_j, \quad |K_j(B)| = \left\lceil \frac{m_j - 1}{2} \right\rceil,$$

and

$$|M_j(C_n - B)| = m'_j, \quad |K_j(C_n - B)| = \left\lceil \frac{m'_j + 2}{2} \right\rceil$$

where  $[x]$  is the largest integer  $\leq x$ .

For  $1 \leq j \leq q$ , the components of the induced graph  $M_j(B) - B_y$  are:

$$\{y_{t_j+1}\}, \{y_{t_j+3}\}, \dots, \{y_{t_j+2k+1}\}, \dots, \{y_{t_j+m_j}\} \quad \text{if } m_j \text{ is odd,}$$

and

$$\{y_{t_j+1}\}, \{y_{t_j+3}\}, \dots, \{y_{t_j+2k+1}\}, \dots, \{y_{t_j+m_j-1}, y_{t_j+m_j}\} \quad \text{if } m_j \text{ is even.}$$

Since each vertex  $v \in V(M_j(C_n - B) - B_y)$  is incident with an edge  $[v, i]$  in  $E(P_n(\alpha))$  for some  $i \in C_n - S$ , the number of components,  $\omega(M_j(B) - B_y)$ , of the induced subgraph  $M_j(B) - B_y$  is equal to  $\left\lceil \frac{m_j+1}{2} \right\rceil$  for  $j = 1, 2, \dots, q$ , and

$$\begin{aligned} (2) \quad \omega(P_n(\alpha) - 5) &= \omega(C_n - B) + \sum_{j=1}^q \omega(M_j(B) - B_y) \\ &= |B| + \sum_{j=1}^q \left\lceil \frac{m_j + 1}{2} \right\rceil \\ &= |B| + \frac{|B|}{2} + \frac{e_1(B)}{2} \end{aligned}$$

where  $|B| = \sum_{j=1}^q m_j$  is used.

By using

$$\left\lceil \frac{m_j - 1}{2} \right\rceil = \begin{cases} \frac{m_j - 1}{2} & \text{if } m_j \text{ is odd,} \\ \frac{m_j + 2}{2} & \text{if } m_j \text{ is even,} \end{cases}$$

$$\left\lceil \frac{m'_j + 2}{2} \right\rceil = \begin{cases} \frac{m'_j + 1}{2} & \text{if } m'_j \text{ is odd,} \\ \frac{m_j + 2}{2} & \text{if } m'_j \text{ is even,} \end{cases}$$

and  $n = \sum_{j=1}^q m_j + \sum_{j=1}^q m'_j$ , we have

$$\begin{aligned}
 (3) \quad |S| &= |B| + |B_y| \\
 &= |B| + \sum_{j=1}^q \left[ \frac{m_j - 1}{2} \right] + \sum_{j=1}^q \left[ \frac{m'_j + 2}{2} \right] \\
 &= |B| + \frac{1}{2} \left( \left( \sum_{j=1}^q m_j \right) - e_1(B) - 2(q - e_1(B)) \right) \\
 &\quad + \frac{1}{2} \left( \left( \sum_{j=1}^q m'_j \right) + e_2(B) + 2(q - e_2(B)) \right) \\
 &= |B| + \frac{n}{2} + \frac{e_1(B) - e_2(B)}{2}.
 \end{aligned}$$

By using (2) and (3), we have

$$t(P_n(\alpha)) \leq \frac{|S|}{\omega(P_n(\alpha) - S)} = \frac{2|B| + n + e_1(B) - e_2(B)}{3|B| + e_1(B)}.$$

**Example 2.** Let  $n, \alpha, P_n(\alpha)$  and  $B$  be the same as in our Example 1. We have

$$\begin{aligned}
 M_1(B) &= y_3y_4, & M_2(B) &= y_6y_7y_8, & M_1(C_{12} - B) &= y_5, \\
 M_2(C_{12} - B) &= y_9y_{10}y_{11}y_{12}y_1y_2.
 \end{aligned}$$

Thus,

$$K_1(B) = \emptyset, \quad K_2(B) = \{y_7\}, \quad K_1(C_{12} - B) = \{y_5\}, \quad K_2(C_{12} - B) = \{y_9, y_{11}, y_1, y_2\}.$$

$S = B \cup B_y = \{1, 3, 5, 7, 10\} \cup \{y_7\} \cup \{y_5\} \cup \{y_9, y_{11}, y_1, y_2\}$ , and the components in  $P_{12}(\alpha) - S$  are:  $\langle 2 \rangle, \langle 4 \rangle, \langle 6 \rangle, \langle 8, 9, y_{12} \rangle, \langle 11, 12, y_{10} \rangle, \langle y_3, y_4 \rangle, \langle y_6 \rangle$  and  $\langle y_8 \rangle$ . (We note that  $[y_{12}, 8]$  and  $[y_{10}, 11]$  do belong to  $E(P_{12}(\alpha))$ ). Thus,  $|S| = 11$ ,  $\omega(P_{12}(\alpha) - S) = 8$ , and  $t(P_{12}(\alpha)) \leq \frac{|S|}{\omega(P_{12}(\alpha) - S)} = \frac{11}{8}$ .

Using our (1), with  $|B| = 5$ ,  $n = 12$ ,  $e_1(B) = 1$  and  $e_2(B) = 1$ , we have

$$t(P_{12}(\alpha)) \leq \frac{2(5) + 12 + 1 - 1}{3(5) + 1} = \frac{22}{16} = \frac{11}{8}.$$

**Corollary 1.1.** Let  $n$  be an integer  $\geq 4$  and  $n \neq 4k + 3$ ,  $\alpha \in S_n$  and  $P_n(\alpha)$  be a cycle permutation graph with  $2n$  vertices. Then

$$t(P_n(\alpha)) \leq \frac{4}{3}.$$

Proof. Let

$$B = \begin{cases} \{1, 3, \dots, n-1\} & \text{if } n \text{ is even,} \\ \{1, 3, \dots, n-2\} & \text{if } n \text{ is odd.} \end{cases}$$

Then  $|B| = \lfloor n/2 \rfloor$ .

For  $n = 4k$ ,  $|B| = 2k$ , and by using (1) in Theorem 1, we have

$$t(P_k(\alpha)) \leq \frac{2(2k) + 4k + e_1(B) - e_2(B)}{3(2k) + e_1(B)} \leq \frac{8k + e_1(B)}{6k + e_1(B)} \leq \frac{4}{3}.$$

For  $n = 4k + 1$  and  $|B| = 2k$ . We claim that  $e_1(B) \geq 1$ . Let  $p(4k + 1, \alpha, B) = \left(\bigcup_{j=1}^q M_j(B)\right) \cup \left(\bigcup_{j=1}^q M_j(C_{4k+1} - B)\right)$  for some positive integer  $q$ . Since  $\sum_{j=1}^q |M_j(B)| = |B| = 2k$  and  $\sum_{j=1}^q |M_j(B)| + \sum_{j=1}^q |M_j(C_{4k+1} - B)| = 4k + 1$ ,  $\sum_{j=1}^q |M_j(C_{4k+1} - B)| = 2k + 1$ .

Consequently,  $e_2(B) \neq 0$ , i.e.,  $e_2(B) \geq 1$ .

By using (1) in Theorem 1 with  $e_2(B) \geq 1$ , we have

$$t(P_{4k+1}(\alpha)) \leq \frac{2(2k) + 4k + 1 + e_1(B) - e_2(B)}{3(2k) + e_1(B)} = \frac{8k + 1 + e_1(B) - e_2(B)}{6k + e_1(B)} \leq \frac{4}{3}$$

where  $3 \leq e_1(B) + 3e_2(B)$  is used.

For  $n = 4k + 2$ ,  $|B| = 2k + 1$ . By using (1) in Theorem 1, we have

$$t(P_{4k+2}(\alpha)) \leq \frac{2(2k + 1) + 4k + 2 + e_1(B) - e_2(B)}{3(2k + 1) + e_1(B)} \leq \frac{8k + 4 + e_1(B)}{6k + 3 + e_1(B)} \leq \frac{4}{3}.$$

□

**Corollary 1.2.** Let  $n$  be an integer  $\geq 4$ ,  $\alpha \in S_n$  and  $P_n(\alpha)$  be a cycle permutation graph with  $2n$  vertices. Then

$$t(P_n(\alpha)) < \frac{3}{2}.$$

Proof. For  $n = 4k, 4k + 1$  and  $4k + 2$ , by Corollary 1.1, we have  $t(P_n(\alpha)) \leq \frac{4}{3} < \frac{3}{2}$ . For  $n = 4k + 3$ , let  $B = \{1, 3, \dots, 4k + 1\}$ . Then  $|B| = 2k + 1$ . By using (1) in Theorem 1, we have

$$\begin{aligned} t(P_{4k+3}(\alpha)) &\leq \frac{2(2k + 1) + (4k + 3) + e_1(B) - e_2(B)}{3(2k + 1) + e_1(B)} \\ &\leq \frac{8k + 5 + e_1(B)}{6k + 3 + e_1(B)} < \frac{3}{2} \end{aligned}$$

where  $k \geq 1$  is used. □

Our Corollary 1.2 confirms the conjecture 2 in [11].



### 3. CYCLE PERMUTATION GRAPHS AND GENERALIZED PETERSEN GRAPHS

We shall show that, for  $n = 4k + 3$  with  $k \geq 1$ , a certain  $P_n(\alpha)$  is isomorphic to a generalized Petersen graph. We shall also show that the toughness of this generalized Petersen graph is  $\leq \frac{4}{3}$ , and we use the results to prove  $t(P_n(\alpha)) \leq \frac{4}{3}$ .

For  $n = 4k + 3$  and  $|B| = 2k + 1$ , by using (1) in Theorem 1, we have

$$\begin{aligned} t(P_{4k+3}(\alpha)) &\leq \frac{2(2k+1) + (4k+3) + e_1(B) - e_2(B)}{3(2k+1) + e_1(B)} \\ &= \frac{8k+5 + e_1(B) - e_2(B)}{6k+3 + e_1(B)}. \end{aligned}$$

In order to have  $t(P_{4k+3}(\alpha)) \leq \frac{4}{3}$ , we need

$$(4) \quad 3 \leq e_1(B) + 3e_2(B).$$

If  $e_2(B_i) \geq 1$ , then (4) holds. If  $e_2(B) = 0$  and  $E_1(B_i) \geq 3$ , then (4) holds. Since  $n = 4k + 3$  and  $|B| = 2k + 1$  are odd integers, we cannot have the case of  $e_2(B) = 0$  and  $e_1(B_i) = 2$ . The only case which we need to consider is  $e_2(B) = 0$  and  $e_1(B_i) = 1$ .

Let  $n = 4k + 3$  with  $k \geq 1$ , and  $B_i = \{1 + i, 3 + i, \dots, (4k + 1) + i\}$  for  $i = 0, 1, \dots, 4k + 2$ . (The integers are taken modulo  $n$ .) Then each  $B_i$  is an independent set of vertices in  $C_n$ . Each of  $p(n, \alpha, B_i)$ ,  $e_1(B_i)$  and  $e_2(B_i)$  are defined in the same way as  $p(n, \alpha, B_0) = p(n, \alpha, B)$ ,  $e_1(B_0) = e_1(B)$  and  $e_2(B_0) = e_2(B)$  respectively.

**Example 3.** We give two cycle permutation graphs. One has  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for some  $i$ . The other one has  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all integers  $i$ .

Let  $P_{11}(\beta)$  be the cycle permutation graph with

$$\beta = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 \\ 10 & 7 & 5 & 9 & 3 & 8 & 4 & 2 & 11 & 1 & 6 \end{pmatrix}$$

and  $B = B_0 = \{1, 3, 5, 7, 9\}$ . Then  $p(11, \beta, B_0)$  is  $M_1(B_0) = y_3y_4y_5 \smile M_1(C_{11} - B_0) = y_6y_7y_8y_9 \smile M_2(B_0) = y_{10}y_{11} \smile M_2(C_{11} - B_0) = y_1y_2 \smile M_1(B_0)$ . Thus,  $e_1(B_0) = 1$  and  $e_2(B_0) = 0$ . For  $B_3 = \{4, 6, 8, 10, 1\}$ ,  $p(11, \beta, B_3)$  is  $M_1(B_3) = y_1y_2 \smile M_1(C_{11} - B_3) = y_3y_4y_5y_6y_7 \smile M_2(B_3) = y_8y_9y_{10} \smile M_2(C_{11} - B_3) = y_{11} \smile M_1(B_3)$ . Thus,  $e_1(B_3) = 1$  and  $e_2(B_3) = 2$ .

Let  $P_{11}(\alpha)$  be the cycle permutation graph with

$$\alpha = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 \\ 1 & 3 & 5 & 7 & 9 & 11 & 2 & 4 & 6 & 8 & 10 \end{pmatrix}$$

with  $B = B_0 = \{1, 3, 5, 7, 9\}$ . Then  $p(11, \alpha, B_0)$  is

$$\begin{aligned} M_1(B_0) &= y_1 y_2 \smile M_1(C_{11} - B_0) = y_3 y_4 \smile M_2(B_0) = y_5 y_6 \\ &\smile M_2(C_{11} - B_0) = y_7 y_8 \smile M_3(B_0) = y_9 \smile M_3(C_{11} - B_0) = y_{10} y_{11} \smile M_1(B_0). \end{aligned}$$

Thus,  $e_1(B_0) = 1$  and  $e_2(B_0) = 0$ .

For  $B_i = \{1 + i, 3 + i, 5 + i, 7 + i, 9 + i\}$ ,  $i = 1, 2, \dots, 10$ ,  $p(11, \alpha, B_i)$  is

$$\begin{aligned} M_1(B_i) &= y_{1+2i} y_{2+2i} \smile M_1(C_{11} - B_i) = y_{3+2i} y_{4+2i} \smile M_2(B_i) \\ &= y_{5+2i} y_{6+2i} \smile M_2(C_{11} - B_i) = y_{7+2i} y_{8+2i} \smile M_3(B_i) = y_{9+2i} \smile M_3(C_{11} - B_i) \\ &= y_{10+2i} y_{11+2i} \smile M_1(B_i). \end{aligned}$$

Thus,  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$ , i.e.,  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all integers  $i$ .

**Theorem 2.** *Let  $P_n(\alpha)$  be a cycle permutation graph with  $2n$  vertices where  $n = 4k + 3$  and  $k \geq 1$ , and  $B_i$  be the same as above with  $e_1(B_i) \geq 3$  or  $e_2(B_i) \geq 1$  for some integer  $i$ . Then*

$$t(P_n(\alpha)) \leq \frac{4}{3}.$$

*Proof.* Replacing  $B$  in Theorem 1 by  $B_i$ , we have the inequality (1). By using the inequality (4), we have  $t(P_n(\alpha)) \leq \frac{4}{3}$ .  $\square$

**Theorem 3.** *Let  $P_n(\alpha)$  be a cycle permutation graph with  $2n$  vertices with  $n = 4k + 3$  and  $k \geq 1$ , and  $B_i$  be the same as above for  $i = 0, 1, \dots, 4k + 2$ . If  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all  $i$ , then  $P_n(\alpha)$  is isomorphic to  $G(n, 2r)$  for some positive integer  $r$  such that  $r$  divides  $2(k + 1)$ .*

In order to prove our Theorem 3, we need the following lemmas.

**Lemma 3.1.** *Let  $P_n(\alpha)$  be a cycle permutation graph with  $2n$  vertices where  $n = 4k + 3$  and  $k \geq 1$ ,  $B_i$  be the same as above with  $e_1(B_i) = 1$ , and  $e_2(B_i) = 0$  for all integers  $i$ , and for some integer  $t$  and some integer  $l$   $M_1(B_t) = y_{l+1} y_{l+2} \dots y_{l+m}$  where  $m$  is a positive odd integer. Then*

$$\alpha\{t - 1, t\} = \{l, l + m + 1\}$$

where

$$\alpha\{t - 1, t\} = \{\alpha(t - 1), \alpha(t)\}.$$

*Proof.* We claim that if  $M_1(B_0) = y_{l+1}y_{l+2} \dots y_{l+m}$ , then  $\alpha\{n-1, n\} = \{l, l+m+1\}$ . Suppose the contrary, i.e.,  $\alpha\{n-1, n\} \neq \{l, l+m+1\}$ . Then there are two cases to be considered:

*Case 1.*  $\alpha^{-1}(l+m+1) \notin \{n-1, n\}$ .

(i)  $\alpha^{-1}(l) \notin \{n-1, n\}$ .

Let  $B_0 = \{1, 3, \dots, 4k+1\}$ ,  $M_1(B_0) = y_{l+1}y_{l+2} \dots y_{l+m}$  with  $m$  being a positive odd integer, and  $B_1 = \{2, 4, \dots, 4k+2\}$ . Then  $C_n - B_1$  consists of all odd integers in  $\{1, 2, \dots, n\}$  and  $\{\alpha^{-1}(i); y_i \in M_1(B_0)\} \subseteq C_n - B_1$ , i.e.,  $M_1(B_0)$  is a chain related to  $C_n - B_1$ . Since  $\alpha^{-1}(l)$  and  $\alpha^{-1}(l+m+1) \notin \{n-1, n\}$ , neither the vertex  $y_l$  nor the vertex  $y_{l+m+1}$  is in the chain  $M_1(B_0)$ . Thus,  $M_1(B_0)$  is a maximal chain related to  $C_n - B_1$ . Since  $|M_1(B_0)| = m$  is an odd integer,  $e_2(B_1) \geq 1$  which is a contradiction to  $e_2(B_i) = 0$  for all integers  $i$ . Hence,  $\alpha^{-1}(l) \in \{n-1, n\}$ .

(ii)  $\alpha^{-1}(l) = n-1$ .

Similar to the proof of (i) in the Case 1. Since  $n-1 \notin C_n - B_1$ ,  $M_1(B_0)$  is a maximal chain related to  $C_n - B_1$ , and  $e_2(B_1) \geq 1$  which is a contradiction.

(iii)  $\alpha^{-1}(l) = n$ .

Similar to the proof of (i) in the Case 1. Replacing  $B_1$  by  $B_{-1} = B_{4k+2}$ ,  $M_1(B_0)$  is a maximal chain related to  $C_n - B_{-1}$ , and  $e_2(B_{-1}) \geq 1$  which is a contradiction. Consequently, we have  $\alpha^{-1}\{l, l+m+1\} \in \{n-1, n\}$ .

*Case 2.*  $\alpha^{-1}(l) \notin \{n-1, n\}$  and  $\alpha^{-1}(l+m+1) \in \{n-1, n\}$ .

(i)  $\alpha^{-1}(l+m+1) = n-1$ .

Similar to the proof of (ii) in the Case 1,  $e_2(B_1) \geq 1$  which is a contradiction.

(ii)  $\alpha^{-1}(l+m+1) = n$ .

Similar to the proof of (iii) in the Case 1,  $e_2(B_{-1}) \geq 1$  which is a contradiction. Hence, we have  $\alpha^{-1}\{l, l+m+1\} = \{n-1, n\}$ , i.e.,  $\alpha\{n-1, n\} = \{l, l+m+1\}$ .

We claim that if, for some nonzero  $t$ ,

$$M_1(B_t) = y_{l+1}y_{l+2} \dots y_{l+m},$$

then  $\alpha\{t-1, t\} = \{l, l+m+1\}$ . Relabeling  $k$  by  $t+k$  for all  $k \in V(C_n)$ , we obtain a new cycle permutation graph  $P_n(\beta)$  for a permutation  $\beta \in S_n$  such that  $\beta(i) = \alpha(t+i)$  for  $i = 1, 2, \dots, n$ .

Obviously,  $P_n(\beta) \cong P_n(\alpha)$  and, in  $P_n(\beta)$ ,  $e_1(B_i) = 1$  and  $e_2(\beta_i) = 0$  for all integers  $i$ . Since  $M_1(B_0) = y_{l+1}y_{l+2} \dots y_{l+m}$  in  $P_n(\beta)$ ,  $\beta\{n-1, n\} = \{l, l+m+1\}$ . Thus,

$$\alpha\{(n-1) + t, n+t\} = \alpha\{t-1, t\} = \{l, l+m+1\},$$

i.e.,

$$\alpha\{t-1, t\} = \{l, l+m+1\}.$$

□

**Lemma 3.2.** *Let  $n = 4k + 3$  with  $k \geq 1$ ,  $P_n(\alpha)$  be a cycle permutation graph with  $B_0 = \{1, 3, \dots, 4k + 1\}$  and  $p(n, \alpha, B_0)$  given by*

$$M_1 \smile M'_1 \smile M_2 \smile M'_2 \smile \dots \smile M_r \smile M'_r \smile M_1$$

where

$$(5) \quad M_j = M_j(B_0) = v_{j,1}v_{j,2} \dots v_{j,m_j}$$

and

$$(6) \quad M'_j = M_j(C_n - B_0) = w_{j,1}w_{j,2} \dots w_{j,m'_j}$$

for  $j = 1, 2, \dots, r$ . If  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all integers  $i$ , and  $|M_i| = m$  is a positive odd integer, then

$$(7) \quad n \smile w_{1,1} \quad \text{and} \quad (n-1) \smile w_r, m'_r$$

or

$$(8) \quad n \smile w_r, m'_r \quad \text{and} \quad (n-1) \smile w_{1,1}$$

where  $w_{1,1}$  and  $w_r, m'_r$  are the initial vertex of  $M'_1$  and the terminal vertex of  $M'_r$ , respectively, and  $q \smile w$  means  $[q, w] \in E(P_n(\alpha))$  where  $q \in C_n$  and  $w \in C'_n$ .

*Proof.* This follows from Lemma 3.1. □

We shall repeatedly use (5), (6), (7) and (8) in the following

**Lemma 3.3.** *Let  $n = 4k + 3$  and  $k \geq 1$ ,  $P_n(\alpha)$  be a cycle permutation graph with  $2n$  vertices, and  $M_j$  and  $M'_j$  for  $j = 1, 2, \dots, r$  be the same as in (5) and (6). If  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all integers  $i$ ,  $|M_1| = m_1$  is an odd positive integer, and  $n \smile w_{1,1}$  and  $(n-1) \smile w_r, m'_r$ , then we have*

- (1)  $(2dr - 1) \smile v_{1,d}$  and  $2dr \smile w_{1,d+1}$  for  $1 \leq d \leq \min_{1 \leq j \leq r} \{|M_j|, |M'_j|\}$ , and
- (2)  $(2sr + 2t - 1) \smile v_{t+1, s+1}$ , and  $(2sr + 2t) \smile w_{t+1, s+1}$  for  $0 \leq s \leq m$  and  $1 \leq t \leq r - 1$ .

Since the proof of Lemma 3.3. is very lengthy, we shall first consider the following examples which demonstrate Lemma 3.3.

**Example 4.** Let  $n = 4(3) + 3 = 15$ ,  $P_n(\alpha)$  be the cycle permutation graph with

$$\alpha = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 \\ 1 & 5 & 9 & 13 & 2 & 6 & 10 & 14 & 3 & 7 & 11 & 15 & 4 & 8 & 12 \end{pmatrix}$$

and  $B_0 = \{1, 3, 5, 7, 9, 11, 13\}$ . Then we have  $M_1 = y_9y_{10}y_{11} \smile M'_1 = y_{12}y_{13}y_{14}y_{15} \smile M_2 = y_1y_2y_3y_4 \smile M'_2 = y_5y_6y_7y_8 \smile M_1$ . Thus,  $e_1(B_0) = 1$  and  $e_2(B_0) = 0$ . In fact, for  $B_i = \{1+i, 3+i, 5+i, 7+i, 9+i, 11+i, 13+i\}$ , we have  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all integers  $i$ . We see that, in this case,  $k = 3$ ,  $r = 2$ ,  $\min_{1 \leq j \leq r} \{|M_j|, |M'_j|\} = 3$ , and  $(15 = n) \smile (w_{1,1} = y_{12} = y_{\alpha(15)})$ , and  $(14 = n - 1) \smile (w_{2,m'_2} = y_8 = y_{\alpha(14)})$ .

$$(1) \quad (2dr - 1) \smile v_{1,d} \text{ and } 2dr \smile w_{1,d+1} \text{ for } d = 1, 2, 3.$$

That is,

$$\begin{aligned} ((2(1)(2) - 1) = 3) &\smile (v_{1,1} = y_9 = y_{\alpha(3)}), \\ ((2(2)(2) - 1) = 7) &\smile (v_{1,2} = y_{10} = y_{\alpha(7)}), \\ ((2(3)(2) - 1) = 11) &\smile (v_{1,3} = y_{11} = y_{\alpha(11)}), \\ ((2(1)(2)) = 4) &\smile (w_{1,2} = y_{13} = y_{\alpha(4)}), \\ ((2(2)(2)) = 8) &\smile (w_{1,3} = y_{14} = y_{\alpha(8)}), \\ ((2(3)(2)) = 12) &\smile (w_{1,4} = y_{15} = y_{\alpha(12)}). \end{aligned}$$

$$(2) \quad (2sr + 2t - 1) \smile v_{t+1,s+1} \text{ and } (2sr + 2t) \smile w_{t+1,s+1} \\ \text{for } s = 0, 1, 2, 3 \text{ and } t = 1.$$

$$\begin{aligned} ((2(0)(2) + 2(1) - 1) = 1) &\smile (v_{2,1} = y_1 = y_{\alpha(1)}), \\ ((2(1)(2) + 2(1) - 1) = 5) &\smile (v_{2,2} = y_2 = y_{\alpha(5)}), \\ ((2(2)(2) + 2(1) - 1) = 9) &\smile (v_{2,3} = y_3 = y_{\alpha(9)}), \\ ((2(3)(2) + 2(1) - 1) = 13) &\smile (v_{2,4} = y_4 = y_{\alpha(13)}), \\ ((2(0)(2) + 2(1)) = 2) &\smile (w_{2,1} = y_5 = y_{\alpha(2)}), \\ ((2(1)(2) + 2(1)) = 6) &\smile (w_{2,2} = y_6 = y_{\alpha(6)}), \\ ((2(2)(2) + 2(1)) = 10) &\smile (w_{2,3} = y_7 = y_{\alpha(10)}), \\ ((2(3)(2) + 2(1)) = 14) &\smile (w_{2,4} = y_8 = y_{\alpha(14)}). \end{aligned}$$

We recall that the generalized Petersen graph  $G(15, 4)$  has

$$V(G(15, 4)) = \{1', 2', \dots, 15', y'_1, y'_2 \dots y'_{15}\}$$

and  $E(G(15, 4)) = \{[i', i' + 1], [y'_i, y'_{i+4}], [i', y'_i]\}$  for  $i = 1, 2, \dots, 15\}$ . Then  $P_{15}(\alpha) \simeq G(15, 4)$  where the isomorphic map  $\theta: V(P_{15}(\alpha)) \rightarrow V(G(15, 4))$  is defined by  $\theta(i) = i'$  and  $\theta(y_i) = y'_{\alpha^{-1}(i)}$  for  $i = 1, 2, \dots, 15$ .

**Example 5.** The following two cycle permutation graphs are isomorphic. But one has the property (7) and the other has the property (8).

Let  $P_7(\alpha)$  be the cycle permutation graph with 14 vertices,

$$\alpha = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 4 & 7 & 3 & 6 & 2 & 5 \end{pmatrix},$$

and  $B_i = \{1+i, 3+i, 5+i\}$  for  $i = 0, 1, \dots, 6$ . Then  $M_1 = M_1(B_0) = y_6 y_7 y_1 \smile M'_1 = M_1(C_7 - B_0) = y_2 y_3 y_4 y_5 \smile M_1$ , and  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for  $i = 0, 1, \dots, 6$ .  $(n = 7) \smile (w_{1,4} = y_5 = y_{\alpha(7)})$  and  $((n - 1) = 6) \smile (w_{1,1} = y_2 = y_{\alpha(6)})$  which is the case of (8).

Let  $P_7(\beta)$  be the cycle permutation graph with 14 vertices,

$$\beta = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 5 & 2 & 6 & 3 & 7 & 4 \end{pmatrix},$$

and  $B_i = \{1+i, 3+i, 5+i\}$  for  $i = 0, 1, \dots, 6$ . Then  $M_1 = M_1(B_0) = y_1 y_2 y_3 \smile M'_1 = M_1(C_7 - B_0) = y_4 y_5 y_6 y_7 \smile M_1$ , and  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for  $i = 0, 1, \dots, 6$ .  $(n = 7) \smile (w_{1,1} = y_4 = y_{\beta(7)})$  and  $((n - 1) = 6) \smile (w_{1,4} = y_7 = y_{\beta(6)})$  which is the case of (7):

$$P_7(\alpha) \simeq G(7, 3) \simeq G(7, 4) \simeq P_7(\beta).$$

We also note that  $r = 1$  for both  $P_7(\alpha)$  and  $P_7(\beta)$ . Also, in  $P_7(\alpha)$ , if we use  $B_3 = \{4, 6, 1\}$  instead of  $B_0 = \{1, 3, 5\}$ , then we have  $M_1 = y_1 y_2 y_3$  and  $M'_1 = y_4 y_5 y_6 y_7$ .

*Proof.* The proof of Lemma 3.2 goes as follows.

(1) We show that  $(2dr - 1) \smile v_{1,d}$  and  $2dr \smile w_{1,d+1}$  for  $1 \leq d \leq m = \min_{1 \leq j \leq r} \{|M_j|, |M'_j|\}$ .

There are two cases to be considered:

*Case 1.*  $r = 1$ .

Let  $B_0 = \{1, 3, \dots, 4k + 1\}$ , and let the maximal chains be

$$M_1(B_0) = M_1 = v_{1,1} \dots v_{1,m_1},$$

and

$$M_1(C_n - B_0) = M'_1 = w_{1,1} w_{1,2} \dots w_{1,m'_1}$$

where  $m_1$  is a positive odd integer, and

$$n \smile w_{1,1} \quad \text{and} \quad (n - 1) \smile (w_{r,m'_r} = w_{1,m'_1}).$$

Since  $r = 1$  and  $n = 4k + 3$ ,  $m_1 = 2k + 1$  and  $m'_1 = 2k + 2$ , i.e.,  $m_1 = m = \min\{|M_1|, |M'_1|\}$ . Let  $B_2 = \{3, 5, \dots, 4k + 1, 4k + 3\}$ . Then

$$\{\alpha^{-1}(i); y_i \in M'_1 - w_{1,1}\} \subseteq C_n - B_2.$$

We claim that  $1 \smile v_{1,1}$ . If not, then  $M'_1 - w_{1,1}$  is a maximal chain related to  $C_n - B_2$ , and  $|M'_1 - w_{1,1}|$  is an odd integer. Thus,  $e_2(B_2) \geq 1$  which is a contradiction to  $e_2(B_i) = 0$  for all integers  $i$ . Hence,  $1 \smile v_{1,1}$ . Since  $M_1 + w_{1,1} - v_{1,1}$  is a maximal chain related to  $B_2$  and  $M'_1 - w_{1,1} + v_{1,1}$  is a maximal chain related to  $C_n - B_2$ , by Lemma 3.1,  $1 \smile v_{1,1}$  (the terminal vertex of  $M'_1 - w_{1,1} + v_{1,1}$ ), and  $2 \smile w_{1,2}$  (the initial vertex of  $M'_1 - w_{1,1} + v_{1,1}$ ).

Repeatedly applying the same argument to  $B_{2dr} = B_{2d}$  for  $1 \leq d \leq m (= 2k + 1)$ , we obtain a maximal chain related to  $B_{2d}$ :

$$M_1 + w_{1,1}w_{1,2} \dots w_{1,d} - v_{1,1}v_{1,2} \dots v_{1,d},$$

a maximal chain related to  $C_n - B_{2d}$ ,  $M'_1 - w_{1,1}w_{1,2} \dots w_{1,d} + v_{1,1}v_{1,2} \dots v_{1,d}$ , and  $(2d - 1) \smile v_{1,d}$  and  $2d \smile w_{1,d+1}$  for  $d = 1, 2, \dots, (2k + 1)$ . (We already know that  $n \smile w_{1,1}$ .)

*Case 2.*  $r > 1$ .

(i) We show that  $2t - 1 \smile v_{t+1,1}$  and  $2t \smile w_{t+1,1}$  for  $t = 1, 2, \dots, r - 1$ .

Let  $B_0 = \{1, 3, \dots, 4k + 1\}$ , let the maximal chains be, for  $j = 2, \dots, r$ ,

$$M_j = M_j(B_0) = v_{j,1}v_{j,2}, \dots, v_{j,m_j},$$

and

$$M'_j = M_j(C_n - B_0) = w_{j,1}w_{j,2} \dots w_{j,m'_j}$$

where  $m_1$  is a positive odd integer, and

$$n \smile w_{1,1} \quad \text{and} \quad (n - 1) \smile w_{r,m'_r}.$$

Similar to the proof of Case 1, let  $B_2 = \{3, 5, \dots, 4k + 1, 4k + 3\}$ . Then

$$\{\alpha^{-1}(i); y_i \in M'_1 - w_{1,1}\} \subseteq C_n - B_2.$$

We claim that  $1 \smile v_{2,1}$ . If not, then  $M'_1 - W_{1,1}$  is a maximal chain related to  $C_n - B_2$ , and  $|M'_1 - w_{1,1}|$  is an odd integer. Thus,  $e_2(B_2) \geq 1$  which is a contradiction to  $e_2(B_i) = 0$  for all integers  $i$ . Hence,  $1 \smile v_{2,1}$ . Consequently,

$$M_1 + w_{1,1}, M_2 - v_{2,1}, M_3, \dots, M_r$$

are maximal chains related to  $B_2$ , and

$$M'_1 - w_{1,1} + v_{2,1}, M'_2, M'_3, \dots, M'_r$$

are maximal chains related to  $C_n - B_2$ . Since  $|M_2 - v_{2,1}| = m_2 - 1$  is odd and the others are even, by Lemma 2.1, we have  $1 \smile v_{2,1}$  (the terminal vertex of  $M'_1 - w_{1,1} + v_{2,1}$ ), and  $2 \smile w_{2,1}$  (the initial vertex of  $M'_2$ ).

Repeatedly applying the same argument to  $B_{2t}$  for  $1 \leq t \leq r - 1$ , we obtain the maximal chains related to  $B_{2t}$ :

$$M_1 + w_{1,1}, M_2 - v_{2,1} + w_{2,1}, M_3 - v_{3,1} + w_{3,1}, \dots, M_{t+1} - v_{t+1,1}, M_{t+2} \dots, M_r,$$

the maximal chains related to  $C_n - B_{2t}$ :

$$M'_1 - w_{1,1} + v_{2,1}, M'_2 - w_{2,1} + v_{3,1}, M'_3 - w_{3,1} + v_{4,1}, \dots, M'_t - w_{t,1} + v_{t+1,1}, \\ M'_{t+1}, M'_{t+2}, \dots, M'_r,$$

and  $(2t - 1) \smile v_{t+1,1}$  and  $2t \smile v_{t+1,1}$  and  $2t \smile w_{t+1,1}$  for  $t = 1, 2, \dots, r - 1$ .

(ii) We show that  $(2dr - 1) \smile v_{1,d}$  and  $2dr \smile w_{1,d+1}$  for  $d = 1, 2, \dots, m$  (where  $m = \min_{1 \leq j \leq r} \{|M_j|, |M'_j|\}$ ). For  $t = r - 1$ , by the Case 2 (i), we obtain the maximal chains related to  $B_{2(r-1)}$ :

$$M_1 + w_{1,1}, M_2 - v_{2,1} + w_{2,1}, M_3 - v_{3,1} + w_{3,1}, \dots, M_r - v_{r,1},$$

the maximal chains related to  $C_n - B_{2(r-1)}$ :

$$M'_1 - w_{1,1} + v_{2,1}, M'_2 - w_{2,1} + v_{3,1}, M'_3 - w_{3,1} + v_{4,1}, \dots, M'_{r-1} - w_{r-1,1} + v_{r,1}, M'_r$$

and  $(2r - 3) \smile v_{r,1}$  and  $(2r - 2) \smile w_{r,1}$ .

Let  $B_{2r} = \{1 + 2r, 3 + 2r, \dots, 4k + 1 + 2r\}$ . Since  $4k + 1 + 2r$  is congruent to  $2r - 2$  modulo  $n$ ,  $\{\alpha^{-1}(i); y_i \in M'_r - w_{r,1}\} \subseteq C_n - B_{2r}$ .

We claim that  $(2r - 1) \smile v_{1,1}$ . If not, then  $M'_r - w_{r,1}$  is a maximal chain related to  $C_n - B_{2r}$ , and  $|M'_r - w_{r,1}| = m'_r - 1$  is an odd integer. Thus,  $e_2(B_{2r}) \geq 1$  which is a contradiction to  $e_2(B_i) = 0$  for all integers  $i$ . Hence,  $(2r - 1) \smile v_{1,1}$ .

The maximal chains related to  $B_{2r}$  are:

$$M_1 - v_{1,1} + w_{1,1}, M_2 - v_{2,1} + w_{2,1}, M_3 - v_{3,1} + w_{3,1}, \dots, M_r - v_{r,1} + w_{r,1},$$

and the maximal chains related to  $C_n - B_{2r}$  are:

$$M'_1 - w_{1,1} + v_{2,1}, M'_2 - w_{2,1} + v_{3,1}, M'_3 - w_{3,1} + v_{4,1}, \dots, M'_r - w_{r,1} + v_{1,1}.$$



Since  $|M_1 - v_{1,1} + w_{1,1}|$  is an odd integer, by Lemma 3.1,  $(2r - 1) \smile v_{1,1}$  (the terminal vertex of  $M'_r - w_{1,1} + v_{1,1}$ ), and  $2r \smile w_{2,1}$  (the initial vertex of  $M'_1 - w_{1,1} + v_{2,1}$ ).

Repeatedly applying the same argument to  $B_{2dr}$  for  $d = 1, 2, \dots, m$ , we obtain the maximal chains related to  $B_{2dr}$ :

$$\begin{aligned} M_1 - v_{1,1}v_{1,2} \dots v_{1,d} + w_{1,1}w_{1,2} \dots w_{1,d} \\ M_2 - v_{2,1}v_{2,2} \dots v_{2,d} + w_{2,1}w_{2,2} \dots w_{2,d} \\ \vdots \\ M_r - v_{r,1}v_{r,2} \dots v_{r,d} + w_{r,1}w_{r,2} \dots w_{r,d}, \end{aligned}$$

the maximal chains related to  $C_n - B_{2dr}$ :

$$\begin{aligned} M'_1 - w_{1,1}w_{1,2} \dots w_{1,d} + v_{2,1}v_{2,2} \dots v_{2,d}, \\ M'_2 - w_{2,1}w_{2,2} \dots w_{2,d} + v_{3,1}v_{3,2} \dots v_{3,d}, \\ \vdots \\ M'_r - w_{r,1}w_{r,2} \dots w_{r,d} + v_{1,1}v_{1,2} \dots v_{1,d}, \end{aligned}$$

and  $(2dr - 1) \smile v_{1,d}$  and  $2dr \smile w_{1,d+1}$  for  $d = 1, 2, \dots, m$ . Thus, the proof of (1) is completed.

(2) Repeatedly applying the same argument as in the proof of Case 2 (ii) to  $B_{2kr}$  for  $k = 1, 2, \dots, m - 1$ , then applying the same argument as in the proof of Case 2 (i) to  $B_{2kr+2t}$  for  $t = 1, 2, \dots, r - 1$ , we obtain the maximal chains related to  $B_{2kr+2t}$ :

$$\begin{aligned} M_1 - v_{1,1}v_{1,2} \dots v_{1,k} + w_{1,1}, w_{1,2} \dots w_{1,k}w_{1,k+1}, \\ M_2 - v_{2,1}v_{2,2} \dots v_{2,k}v_{2,k+1} + w_{2,1}w_{2,2} \dots w_{2,k}w_{2,k+1}, \\ \vdots \\ M_t - v_{t,1}v_{t,2} \dots v_{t,k}v_{t,k+1} + w_{t,1}w_{t,2} \dots w_{t,k}w_{t,k+1}, \\ M_{t+1} - v_{t+1,1}v_{t+1,2} \dots v_{t+1,k}v_{t+1,k+1} + w_{t+1,1}w_{t+1,2} \dots w_{t+1,k}, \\ M_{t+2} - v_{t+2,1}v_{t+2,2} \dots v_{t+2,k} + w_{t+2,1}w_{t+2,2} \dots w_{t+2,k}, \\ \vdots \\ M_r - v_{r,1}v_{r,2} \dots v_{r,k} + w_{r,1}w_{r,2} \dots w_{r,k}, \end{aligned}$$

the maximal chains related to  $C_n - B_{2kr+2t}$ :

$$\begin{aligned}
 &M'_1 - w_{1,1}w_{1,2} \dots w_{1,k}w_{1,k+1} + v_{2,1}v_{2,2} \dots v_{2,k}v_{2,k+1}, \\
 &M'_2 - w_{2,1}w_{2,2} \dots w_{2,k}w_{2,k+1} + v_{3,1}v_{3,2} \dots v_{3,k+1}, \\
 &\quad \vdots \\
 &M'_t - w_{t,1}w_{t,2} \dots w_{t,k}w_{t,k+1} + v_{t+1,1}v_{t+1,2} \dots v_{t+1,k}v_{t+1,k+1}, \\
 &M'_{t+1} - w_{t+1,1}w_{t+1,2} \dots w_{t+1,k} + v_{t+2,1}v_{t+2,2} \dots v_{t+2,k}, \\
 &\quad \vdots \\
 &M'_r - w_{r,1}w_{r,2} \dots w_{r,k} + v_{1,1}v_{1,2} \dots v_{1,k},
 \end{aligned}$$

and

$$(2kr + 2t - 1) \smile v_{t+1,k+1} \quad \text{and} \quad (2kr + 2t) \smile w_{t+1,k+1}$$

for  $k = 1, 2, \dots, m - 1$  and  $t = 1, 2, \dots, r - 1$ . □

**Lemma 3.4.** *Let  $n = 4k + 3$  with  $k \geq 1$ ,  $P_n(\alpha)$ ,  $B_i$ ,  $M_j$  and  $M'_j$  for  $j = 1, 2, \dots, r$  be the same as in Lemma 3.3. If  $|M_1| = m$  is an odd positive integer,  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all integers  $i$ , and  $n \smile w_{1,1}$  and  $(n - 1) \smile w_{r,m'_r}$ , then  $|M_j| = m + 1$  for  $j = 2, 3, \dots, r$ , and  $|M'_j| = m + 1$  for  $j = 1, 2, \dots, r$  where  $r(m + 1) = 2(k + 1)$ .*

*Proof.* Let  $|M_1| = m = \min_{1 \leq j \leq r} \{|M_j|, |M'_j|\}$ . We claim that  $|M_j| = m + 1$  for  $j = 2, 3, \dots, r$  and  $|M'_j| = m + 1$  for  $j = 1, 2, \dots, r$ . First, we show that  $|M'_1| > m$ . Suppose  $|M'_1| = m$ . Then by Lemma 3.3, Case 2 (ii),  $2mr \smile w_{1,m+1}$ . Since  $|M'_1| = m$ ,  $w_{1,m+1} = v_{2,1}$ . By Lemma 3.3, Case 2 (i),  $1 \smile v_{2,1}$ . Since  $P_n(\alpha)$  is a cycle permutation graph, each vertex in  $C_n$  is incident with exactly one vertex in  $C'_n$ . Thus,  $2mr \equiv 1 \pmod{n}$ , i.e.,  $2mr - 1 = n = 4k + 3$  which is a contradiction to  $2mr \leq 4k + 3$ . Hence,  $|M'_1| > m$ .

Suppose  $|M_t| = m$  for some  $t = 2, 3, \dots, r$ . Then

$$M_t(B_{2mr}) = M_t - v_{t,1}v_{t,2} \dots v_{t,m} + w_{t,1}w_{t,2} \dots w_{t,m},$$

i.e.,  $M_t(B_{2mr}) = w_{t,1}w_{t,2} \dots w_{t,m}$  is a maximal chain related to  $B_{2mr}$ . Repeatedly using the procedure in Lemma 3.3, Case 2 (i) with  $B_{2mr+2(t-1)}$ , we have  $(2mr + 2(t - 1) - 1) \smile v_{t,m+1}$ . Since  $|M_t| = m$ ,  $v_{t,m+1} = w_{t,1}$ . By Lemma 3.3, Case 2 (i),  $w_{t,1} \smile 2t - 2$ . Thus,  $2mr + 2t - 3 \equiv 2t - 2 \pmod{n}$ ,  $2mr - 1 = n = 4k + 3$  which is a contradiction to  $2mr \leq 4k + 3$ . Hence,  $|M_t| > m$  for  $t = 2, 3, \dots, r$ . By using a similar reasoning, we have  $|M'_t| > m$  for  $t = 1, 2, \dots, r$ .

We want to show that

$$|M_j| = m + 1 \quad \text{for } j = 2, 3, \dots, r,$$

and

$$|M'_j| = m + 1 \quad \text{for } j = 1, 2, \dots, r.$$

We know that  $|M_j| \geq m + 1$  for  $j = 2, 3, \dots, r$ , and  $|M'_j| \geq m + 1$  for  $j = 1, 2, \dots, r$ . By using the procedure in Lemma 3.3, Case 2 (i) with  $B_{2mr+2(r-1)}$  repeatedly, we have  $(2mr + 2(r - 1) - 1) \sim w_{(r-1)+1, m+1}$  and  $(2mr + 2(r - 1)) \sim w_{(r-1)+1, m+1} = w_{r, m+1}$ . If  $w_{r, m+1}$  is the terminal vertex of  $M'_r$ , then  $2mr + 2(r - 1) \equiv 4k + 2 \pmod{n}$ , i.e.,  $2r(m + 1) = 4k + 4$ . Hence,  $|M_j| = m + 1$  for  $j = 2, 3, \dots, r$ ,  $|M'_j| = m + 1$  for  $j = 1, 2, \dots, r$ , and  $r$  divides  $2(k + 1)$ . If  $w_{r, m+1}$  is not the terminal vertex of  $M'_r$ , then, by using the procedure in Lemma 3.3, Case 2 (ii) with  $B_{2mr+2r}$ , we have  $(2(m+1)r-1) \sim v_{1, m+1}$ . Since  $v_{1, m+1} = w_{1, 1}$  and  $4k+3 \sim w_{1, 1}$ ,  $2(m+1)r-1 \equiv 4k+3 \pmod{n}$ , i.e.,  $2r(m+1) = 4k+4$ . Hence,  $|M_j| = m+1$  for  $j = 2, 3, \dots, r$ ,  $|M'_j| = m+1$  for  $j = 1, 2, \dots, r$  and  $r$  divides  $2(k + 1)$ .

*Proof of Theorem 3* goes as follows. We want to show that, for  $n = 4k + 3$  with  $k \geq 1$ ,  $P_n(\alpha)$  is isomorphic to  $G(n, 2r)$  for some  $r$  which divides  $2(k + 1)$ . Let  $V(G(n, 2r)) = \{1', 2', \dots, (4k + 3)', y'_1, y'_2, \dots, y'_{4k+3}\}$  and  $E(G(n, 2r)) = \{[i', (i+1)'], [y'_i, y'_{i+2r}], [i', y'_i] \text{ for } i = 1, 2, \dots, 4k + 3\}$ . Also, let  $V(P_n(\alpha)) = \{1, 2, \dots, 4k + 3, y_1, y_2, \dots, y_{4k+3}\}$  and  $E(P_n(\alpha)) = \{[i, i + 1], [y_i, y_{i+1}][i, y_{\alpha(i)}] \text{ for } i = 1, 2, \dots, 4k + 3\}$ . In  $P_n(\alpha)$ , let  $B_i = \{1 + i, 3 + i, \dots, (4k + 1) + i\}$ , with  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for  $i = 0, 1, \dots, 4k + 2$ , the maximal chains  $M_j = M_j(B_0) = v_{j, 1}v_{j, 2} \dots v_{j, m_j}$  and  $M'_j = M_j(C_n - B_0) = w_{j, 1}w_{j, 2} \dots w_{j, m'_j}$ , for  $j = 1, 2, \dots, r$ , where  $r$  divides  $2(k + 1)$ .

*Case 1.*  $n \sim w_{1, 1}$  and  $(n - 1) \sim w_{r, m'_r}$ . We define a map  $\theta: V(P_n(\alpha)) \rightarrow V(G(n, 2r))$  by  $\theta(i) = i'$  and  $\theta(y_i) = y'_{\alpha^{-1}(i)}$  for  $i = 1, 2, \dots, 4k + 3$ . Then  $\theta$  is a well defined map between the vertices of these two cubic graphs. We show that  $\theta$  preserves the edges: Since  $[i, j] \in E(P_n(\alpha))$  if and only if  $j = i + 1$  and  $[\theta(i), \theta(j)] = [i', j'] \in G(n, 2r)$  if and only if  $j' = i' + 1$ ,  $\theta[i, j] = [\theta(i), \theta(j)]$  for  $i, j = 1, 2, \dots, 4k + 3$  and  $i \neq j$ . Since  $[i, y_j] \in E(P_n(\alpha))$  if and only if  $j = \alpha(i)$  and  $[\theta(i), \theta(y_j)] = [i', y'_{\alpha^{-1}(j)}] \in E(G(n, 2r))$  if and only if  $j = \alpha(i)$ ,  $\theta[i, y_j] = [\theta(i), \theta(y_j)]$  for  $i, j = 1, 2, \dots, 4k + 3$ . We know that  $[y_i, y_j] \in E(P_n(\alpha))$  if and only if  $j = i + 1$ , and  $[\theta(y_i), \theta(y_j)] = [y'_{\alpha^{-1}(i)}, y'_{\alpha^{-1}(j)}]$ . Say,  $\alpha^{-1}(i) = q$  for some  $q$  such that  $1 \leq q \leq 4k + 3$ . Then  $\alpha(q) = i$ . By (1) and (2) of lemma 3.3, we know that  $[y_i, y_{i+1}] \in E(P_n(\alpha))$  if and only if  $i + 1 = \alpha(q) + (2r)$ . By Lemma 3.4, this holds for all  $i = 1, 2, \dots, 4l + 3$ . Since  $2r(m + 1) = 4k + 4 \equiv 1 \pmod{n}$ ,  $[\theta(y_i), \theta(y_{i+1})] = [y'_{\alpha^{-1}(i)}, y'_{\alpha^{-1}(i+1)}] = [y'_q, y'_{q+2r}] \in E(G(n, 2r))$ . Thus,  $\theta[y_i, y_j] = [\theta(y_i), \theta(y_j)]$  for all  $i, j = 1, 2, \dots, 4k + 3$  and  $i \neq j$ . Hence,  $P_n(\alpha) \simeq G(n, 2r)$ .

*Case 2.*  $n \sim w_{r, M'_r}$  and  $(n-1) \sim w_{1,1}$ . Relabeling the vertices on  $C'_n$  by  $v_{1, m_1} \rightarrow z_1, v_{1, m_1-1} \rightarrow z_2, \dots, v_{1,1} \rightarrow z_{m_1}, w_{r, m'_r} \rightarrow z_{m_1+1}, z_{r, m'_r-1} \rightarrow z_{m_1+2}, \dots, w_{1,1} \rightarrow z_n$ , we obtain a new cycle permutation graph  $P_n(\beta)$  such that  $P_n(\beta)$  has a cycle with vertex set  $\{z_1, z_2, \dots, z_n\}$  and  $n \sim z_{m_1+1}$  (the initial vertex of  $M_1(C'_n - B_0)$ ) and  $(n-1) \sim w_{r, m'_r}$  (the end vertex related to  $M_r(C'_n - B_0)$ ). Clearly, by Case 1 above, we have  $P_n(\beta) = P_n(\alpha) \simeq G(n, 2r)$ .  $\square$

We know that  $G(n, t)$  is isomorphic to a cycle permutation graph  $P_n(\alpha)$  for some  $\alpha \in S_n$  if and only if  $t$  and  $n$  are relatively prime. The following example shows that the converse of Theorem 3 does not hold.

**Example 6.** Consider  $G(11, 4)$  where

$$V(G(11, 4)) = \{1, 2, \dots, 11, y_1, y_2, \dots, y_{11}\},$$

and  $E(G(11, 4)) = \{[i, i+1], [i, y_i], [y_i, y_{i+4}] \text{ for } i = 1, 2, \dots, 11\}$ , i.e., the outer cycle of  $G(11, 4)$ ,  $C_x$ , is  $1 - 2 - \dots - 11 - 1$ , and the inner cycle of  $G(11, 4)$ ,  $C_y$ , is  $y_1 - y_5 - y_9 - y_2 - y_6 - y_{10} - y_3 - y_7 - y_{11} - y_4 - y_8 - y_1$ . Let

$$\gamma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 \\ 1 & 5 & 9 & 2 & 6 & 10 & 3 & 7 & 11 & 4 & 8 \end{pmatrix}$$

and

$$\alpha = \gamma^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 \\ 1 & 4 & 7 & 10 & 2 & 5 & 8 & 11 & 3 & 6 & 9 \end{pmatrix}.$$

Clearly,  $G(11, 4) \simeq P_{11}(\alpha)$ . Let  $B_0 = \{1, 3, 5, 7, 9\}$ . Then the maximal chains in  $G(11, 4)$  are:

$$M_1 = y_3 y_7 \sim M'_1 = y_{11} y_4 y_8 \sim M_2 = y_1 y_5 y_9 \sim M'_2 = y_2 y_6 y_{10} \sim M_1.$$

Thus,  $r = 2$  and  $m = 2$ . But  $e_1(B_0) = 1$  and  $e_2(B_0) = 2$ .

**Theorem 4.** Let  $n = 4k + 3$  with  $k \geq 1$  and  $r$  divide  $2(k + 1)$ . Then,  $t(G(n, 2r)) \leq 4/3$ .

*Proof.* Let  $V(G(n, 2r)) = \{1, 2, \dots, 4k+3, y_1, y_2, \dots, y_{4k+3}\}$  and  $E(G(n, 2r)) = \{[i, i+1], [i, y_i], [y_i, y_{i+2r}] \text{ for } i = 1, 2, \dots, 4k+3\}$ .

*Case 1.*  $r = 2(k + 1)$ ,  $G(n, 2r) = G(n, 1)$ . Take the disconnecting set  $S = \{1, 3, 5, \dots, 4k+1, y_2, y_4, y_6, \dots, y_{4k+2}\}$ . Then,  $\omega(G(n, 1) - S)$ , the set of components induced from  $G(n, 1) - S$  is  $\{\{2\}, \{4\}, \{6\}, \dots, \{4k\}, \{y_3\}, \{y_5\}, \{y_7\}, \dots, \{y_{4k+1}\}, \{4k+2, 4k+3, y_{4k+3}, y_1\}\}$ . Thus,  $|S| = (2k+1) + (2k+1) = 4k+2$ ,  $\omega(G(n, 1) - S) = 2k + 2k + 1 = 4k + 1$  and

$$t(G(n, 1)) \leq \frac{4k+2}{4k+1} < \frac{4}{3}.$$

*Case 2.*  $r = 1$ .  $G(n, 2r) = G(n, 2)$ . It was proved in [3] and [8] that  $t(G(n, 2)) \leq \frac{4}{3}$ .

*Case 3.*  $r = k + 1$ .  $G(n, 2r) = G(n, 2k + 2)$ . Let  $B_0 = \{1, 3, 5, \dots, 4k + 1\}$ . Then the partition of maximal chains is:

$$\begin{aligned} & y_{2k+1} \smile y_{4k+3}y_{2k+2} \smile y_1y_{2k+3} \smile y_2y_{2k+4} \\ & \smile y_3y_{2k+5} \smile y_4y_{2k+6} \smile \dots \smile y_{2k-1}y_{4k+1} \smile y_{2k}y_{4k+2}, \end{aligned}$$

and the maximal chains are:

$$\begin{aligned} M_1(B_0) &= y_{2k+1}, & M_1(C_n - B_0) &= y_{4k+3}y_{2k+2}, \\ M_2(B_0) &= y_1y_{2k+3}, & M_2(C_n - B_0) &= y_2y_{2k+4}, \\ & \vdots \\ M_{k+1}(B_0) &= y_{2k-1}y_{4k-1}, & M_{k+1}(C_n - B_0) &= y_{2k}y_{4k+2}. \end{aligned}$$

Let  $B_y = \{y_{2k+2}, y_2, y_{2k+4}, y_4, y_{2k+6}, \dots, y_{2t}, y_{2k+2t+2}, \dots, y_{2k-2}, y_{4k}, y_{2k}\}$  and  $S = B_0 \cup B_y$ . Then the components of  $G(n, 2k + 2) - S$  are the following sets:

$$\begin{aligned} & \{2\}, \{4\}, \dots, \{2t\}, \dots, \{4k\}, \\ & \{y_1, y_{2k+3}\}, \{y_3, y_{2k+5}\}, \dots, \{y_{2t-1}, y_{2k+2t+1}\}, \dots, \\ & \{y_{2k-1}, y_{4k+1}\}, \{y_{4k+3}, 4k + 3, 4k + 2, y_{4k+2}, y_{2k+1}\}. \end{aligned}$$

Thus,

$$\begin{aligned} |S| &= (2k + 1) + 2(k - 1) + 2 = 4k + 1, \\ \omega(G(n, 2k + 2) - S) &= 2k + k + 1 = 3k + 1, \\ \text{and } t(G(n, 2k + 2)) &\leq \frac{|S|}{\omega(G(n, 2k + 2) - S)} = \frac{4k + 1}{3k + 1} < \frac{4}{3}. \end{aligned}$$

*Case 4.* Let  $1 < r < k + 1$  and  $r(m + 1) = 2k + 2$ . Let  $B_0 = \{1, 3, 5, \dots, 4k + 1\}$ . Then the partition of maximal chains is:

$$\begin{aligned} & y_{2r-1}y_{4r-1} \dots y_{2mr-1} \smile y_{4k+3}y_{2r}y_{4r} \dots y_{2mr} \\ & \smile y_1y_{1+2r}y_{1+4r} \dots y_{1+2mr} \smile y_2y_{2+2r}y_{2+4r} \dots y_{2+2mr} \\ & \smile y_{1+2(3-2)}y_{1+2(3-2)+2r}y_{1+2(3-2)+4r} \dots y_{1+2(3-2)+2mr} \\ & \smile y_{2+2(3-2)}y_{2+2(3-2)+2r}y_{2+2(3-2)+4r} \dots y_{2+2(3-2)+2mr} \\ & \vdots \\ & \smile y_{1+2(r-2)}y_{1+2(r-2)+2r}y_{1+2(r-2)+4r} \dots y_{1+2(r-2)+2mr} \\ & \smile y_{2+2(r-2)}y_{2+2(r-2)+2r}y_{2+2(r-2)+4r} \dots y_{2+2(r-2)+2mr}. \end{aligned}$$

Since  $1 < r < k + 1$  and  $r(m + 1) = 2k + 2$ ,  $m + 1 \geq 3$  and  $m \geq 2$ . If  $m = 2$ , then

$$|M_1(C_n - B_0)| = |y_{4k+3}y_{2r}y_{4r} \dots y_{2mr}| = m + 1$$

is a positive odd integer. Thus,  $e_2(B_0) \geq 1$  and by (1) in Theorem 1 or by (4),  $t(G(n, 2r)) \leq 4/3$ , since  $G(n, 2r)$  is isomorphic to a cycle permutation graph. If  $m \geq 3$ , then we define an independent set of vertices in  $C_n$  by

$$B = (B_0 \cup \{4r\}) \setminus \{4r - 1, 4r + 1\}.$$

Then, since  $m \geq 3$  and  $4r \neq 2mr$ ,  $y_{4r-1}, y_{6r}, \dots, y_{2mr}$  and  $y_{4r+1}$  are three maximal chains, related to  $C_n - B$ , of odd cardinalities. Thus,  $e_2(B) \geq 3$  and by (1) in Theorem 1 or by (4),  $t(G(n, 2r)) \leq \frac{4}{3}$  since  $G(n, 2r)$  is isomorphic to a cycle permutation graph.  $\square$

**Corollary 4.1.** For  $n \geq 4$  and  $\alpha \in S_n$ ,

$$t(P_n(\alpha)) \leq \frac{4}{3}.$$

*P r o o f.* If  $n \neq 4k + 3$ , then, by Corollary 1.1,  $t(P_n(\alpha)) \leq \frac{4}{3}$  for every  $\alpha \in S_n$ . By Theorem 2, if  $e_1(B_i) \geq 3$  or  $e_2(B_i) \geq 1$ , then  $t(P_{4k+3}(\alpha)) \leq \frac{4}{3}$  for every  $\alpha \in S_n$  and any positive integer  $k$ . We know that the case of  $n = 4k + 3$  with  $k \geq 1$ ,  $e_1(B_i) = 2$  and  $e_2(B_i) = 0$  does not exist, because this means that there are exactly 2 maximal chains with odd cardinalities and the rest are of even cardinalities. Then the total number of vertices in  $C'_n$  is even which contradicts  $n = 4k + 3$ . Thus, the remaining case which we have to consider is  $n = 4k + 3$  with  $k \geq 1$ ,  $e_1(B_i) = 1$  and  $e_2(B_i) = 0$  for all integers  $i$ . By Theorem 3,  $P_{4k+3}(\alpha)$ , for every  $\alpha \in S_n$ , is isomorphic to  $G(4k + 3, 2r)$  for some positive integer  $r$  which divides  $2k + 2$ . By Theorem 4,  $t(G(4k + 3, 2r)) \leq \frac{4}{3}$ . Hence,  $t(P_n(\alpha)) \leq \frac{4}{3}$  for every integer  $n \geq 4$  and every  $\alpha \in S_n$ .  $\square$

Our Corollary 4.1 confirms the conjecture 1 in [11].

#### References

- [1] *D. Bauer, E. Schmeichel, and H. J. Veldman*: Some recent results on long cycles in tough graphs. Graph Theory, Combinatorics, and Applications—Proceedings of the Sixth Quadrennial International Conference on the Theory and Applications of Graphs (Y. Alavi, G. Chartrand, O.R. Oellermann, and A. J. Schwenk, eds.). John Wiley & Sons, New York, 1991, pp. 113–121.
- [2] *D. Bauer, E. Schmeichel, and H. J. Veldman*: Cycles in tough graphs—updating the last four years. Graph Theory, Combinatorics, and Applications—Proceedings of the Seventh Quadrennial International Conference on the Theory and Applications of Graphs (Y. Alavi and A. J. Schwenk, eds.). Wiley & Sons, New York, 1995, pp. 19–34.

- [3] *C. Y. Chao and S. C. Han*: A note on the toughness of generalized Petersen graphs. *J. Math. Res. Exposition* 12 (1992), 183–186.
- [4] *C. Y. Chao and S. C. Han*: On the classification and toughness of generalized permutation star-graphs. *Czechoslovak Math. J.* 47 (1997), 431–452.
- [5] *G. Chartrand and R. J. Wilson*: The Petersen graph. *Graphs and Applications* (F. Harary and J. S. Maybee, eds.). John Wiley & Sons, 1982, pp. 69–99.
- [6] *V. Chvátal*: Tough graphs and hamiltonian circuits. *Math. Slovaca* 28 (1979), 215–228.
- [7] *W. Döfler*: On mapping graphs and permutation graphs. *Math. Slovaca* 28 (1979), 277–288.
- [8] *K. Ferland*: On the toughness of some generalized Petersen graphs. *Ars Combin.* (1993), 65–88.
- [9] *W. Goddard*: The toughness of cubic graphs. *Graphs Combin.* 12 (1996), 17–22.
- [10] *D. Guichard, B. Piazza, and S. Stueckle*: On the vulnerability of permutation graphs of complete and complete bipartite graphs. *Ars Combin.* 31 (1991), 149–157.
- [11] *B. Piazza, R. Ringeisen, and S. Stueckle*: On the vulnerability of cycle permutation graphs. *Ars Combin.* 29 (1990), 289–296.
- [12] *W. T. Tutte*: On the algebraic theory of graph colorings. *J. Combin. Theory* 1 (1960), 15–50.

*Authors' address:* Chong-Yun Chao, Shaocen Han, Department of Mathematics, University of Pittsburgh, Pittsburgh, PA 15260, USA, e-mail: [cyc@pitt.edu](mailto:cyc@pitt.edu).