

Alexander Lomtadze; P. Torres

On a two-point boundary value problem for second order singular equations

*Czechoslovak Mathematical Journal*, Vol. 53 (2003), No. 1, 19–43

Persistent URL: <http://dml.cz/dmlcz/127778>

## Terms of use:

© Institute of Mathematics AS CR, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A TWO-POINT BOUNDARY VALUE PROBLEM  
FOR SECOND ORDER SINGULAR EQUATIONS

A. LOMTATIDZE, Brno, and P. TORRES, Granada

(Received November 3, 1999)

*Abstract.* The problem on the existence of a positive in the interval  $]a, b[$  solution of the boundary value problem

$$u'' = f(t, u) + g(t, u)u'; \quad u(a+) = 0, \quad u(b-) = 0$$

is considered, where the functions  $f$  and  $g: ]a, b[ \times ]0, +\infty[ \rightarrow \mathbb{R}$  satisfy the local Carathéodory conditions. The possibility for the functions  $f$  and  $g$  to have singularities in the first argument (for  $t = a$  and  $t = b$ ) and in the phase variable (for  $u = 0$ ) is not excluded. Sufficient and, in some cases, necessary and sufficient conditions for the solvability of that problem are established.

*Keywords:* second order singular equation, two-point boundary value problem, solvability

*MSC 2000:* 34B10, 34B18

## 1. STATEMENT OF THE PROBLEM AND FORMULATION OF MAIN RESULTS

The following notation is used throughout the paper.

$\mathbb{R}$  is the set of real numbers,  $\mathbb{R}_+ = [0, +\infty[$ .

$L(]a, b[; D)$ , where  $D \subset \mathbb{R}$ , is the set of functions  $p: ]a, b[ \rightarrow D$  which are Lebesgue integrable on the segment  $[a, b]$ .

$L_{\text{loc}}(]a, b[; D)$ , where  $D \subset \mathbb{R}$ , is the set of functions  $u: ]a, b[ \rightarrow D$  which are Lebesgue integrable on each segment contained in  $]a, b[$ .

$C([a, b]; D)$ , where  $D \subset \mathbb{R}$ , is the set of continuous functions  $u: [a, b] \rightarrow D$ .

---

Supported by the Grant No. 201/99/0295 of the Grant Agency of the Czech Republic.

$AC'([a, b]; D)$ , where  $D \subset \mathbb{R}$ , is the set of functions  $u: [a, b] \rightarrow D$  which are absolutely continuous together with their first derivatives on  $[a, b]$ .

$AC'_{\text{loc}}(I; D)$ , where  $I \subseteq ]a, b[$ ,  $D \subset \mathbb{R}$ , is the set of functions  $u: I \rightarrow D$  such that  $u \in AC'(I_0; D)$  for every segment  $I_0 \subset I$ .

$\text{Car}(]a, b[ \times D; \mathbb{R})$ , where  $D \subset \mathbb{R}$ , is the Carathéodory class, i.e., the set of functions  $f: ]a, b[ \times D \rightarrow \mathbb{R}$  such that  $f(t, \cdot): D \rightarrow \mathbb{R}$  is continuous for almost all  $t \in ]a, b[$ ,  $f(\cdot, x): ]a, b[ \rightarrow \mathbb{R}$  is measurable for all  $x \in D$ , and

$$\sup\{|f(\cdot, x)|: x \in D_0\} \in L(]a, b[; \mathbb{R}_+)$$

for any compact  $D_0 \subset D$ .

$\text{Car}_{\text{loc}}(]a, b[ \times D; \mathbb{R})$ , where  $D \subset \mathbb{R}$ , is the set of functions  $f: ]a, b[ \times D \rightarrow \mathbb{R}$  whose restrictions to  $[a + \varepsilon, b - \varepsilon] \times D$  belong to  $\text{Car}([a + \varepsilon, b - \varepsilon] \times D; \mathbb{R})$  for any  $\varepsilon \in ]0, \frac{1}{2}(b - a)[$ .

$$[p]_- = \frac{1}{2}(|p| - p).$$

$u(s+)$  and  $u(s-)$  are one-sided limits of the function  $u$  at the point  $s$  from the right and from the left, respectively.

Consider the boundary value problem

$$(1.1) \quad u'' = f(t, u) + g(t, u)u',$$

$$(1.2) \quad u(a+) = 0, \quad u(b-) = 0,$$

where  $f, g \in \text{Car}_{\text{loc}}(]a, b[ \times ]0, +\infty[; \mathbb{R})$ . Under a solution of problem (1.1), (1.2) we understand a function  $u \in AC'_{\text{loc}}(]a, b[; ]0, +\infty[)$  satisfying equation (1.1) almost everywhere in  $]a, b[$  and boundary conditions (1.2).

The aim of the present paper is to investigate the problem of solvability of problem (1.1), (1.2) provided the functions  $f$  and  $g$  possess singularities both in the independent (for  $t = a$  and  $t = b$ ) and in the phase (for  $u = 0$ ) variable. Singular problems of such a type arise frequently in applications (cf., for example, [1], [3], [4], [6]–[8], [22]–[24]). The first essential step in their investigation was made by S. Taliaferro in his work [25] in which he established a necessary and sufficient condition for the solvability of problem (1.1), (1.2) with  $g(t, x) \equiv 0$  and  $f(t, x) = -h(t)/x^\lambda$ , where  $\lambda > 0$  and  $h \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$ . Problem (1.1), (1.2) has been more often considered in the case when the function  $g$  does not depend on the second argument, and  $f(t, x) \leq 0$  for  $a < t < b$ ,  $x > 0$  (cf., for example, [1]–[18], [21]–[27] and references therein). In that case, equation (1.1) is in its turn easily reduced to a two-term equation of the type

$$(1.1_0) \quad u'' = f(t, u)$$

with nonpositive right-hand side. The restriction on the sign of the function  $f$  was overcome for the first time in [19], where criteria for the solvability of problem (1.1<sub>0</sub>), (1.2) were established under the assumption that

$$(1.3) \quad \int_a^b (s-a)(b-s)f_r^*(s) ds < +\infty \quad \text{for } r > 1,$$

where  $f_r^*(\cdot) = \max\{|f(\cdot, x)|: \frac{1}{r} \leq x \leq r\}$  for  $r > 1$  (see also later works [12]–[14]). In [17], the equation of a more general type  $u'' = H(t, u, u')$  was considered, but again under the assumption that the function  $H$  is nonpositive. So, in spite of a large number of publications, the question of the solvability of problem (1.1), (1.2) has not yet been studied thoroughly enough. Below we will give new sufficient, and in some cases, necessary and sufficient conditions for the solvability of problem (1.1), (1.2). Moreover, as it has been noted above, the possibility for both the functions  $f$  and  $g$  to have singularities in the first argument and in the phase variable, is not excluded. Note also that Theorem 1.2 below enables one to establish criteria for the solvability of problem (1.1), (1.2) even in the case when condition (1.3) is not fulfilled.

Before we proceed to formulating the main results, we introduce the following definition.

**Definition 1.1.** The continuous function  $\sigma: ]a, b[ \rightarrow ]0, +\infty[$  is said to be a lower (upper) function of equation (1.1) if  $\sigma \in AC'_{\text{loc}}(]a, b[ \setminus \{t_1, t_2, \dots, t_n\}; ]0, +\infty[)$ , where  $a < t_1 < t_2 < \dots < t_n < b$ , there exist finite limits  $\sigma(a+)$ ,  $\sigma(b-)$ ,  $\sigma'(t_i+)$ ,  $\sigma'(t_i-)$ ,  $i = \overline{1, n}$ ,

$$\sigma'(t_i-) < \sigma'(t_i+) \quad (\sigma'(t_i-) > \sigma'(t_i+)), \quad i = \overline{1, n},$$

and almost everywhere in  $]a, b[$  the inequality

$$\sigma''(t) \geq f(t, \sigma(t)) + g(t, \sigma(t))\sigma'(t) \quad (\sigma''(t) \leq f(t, \sigma(t)) + g(t, \sigma(t))\sigma'(t))$$

is fulfilled.

Definition 1.1 is a particular case of the definition of lower and upper functions introduced in [15] (see also [17] and [19]).

**Theorem 1.1.** *Let  $\sigma_1$  and  $\sigma_2$  be respectively lower and upper functions of equation (1.1) and let*

$$(1.4) \quad \begin{aligned} \sigma_1(t) &\leq \sigma_2(t) \quad \text{for } a < t < b, \\ \sigma_1(a+) &= 0, \quad \sigma_1(b-) = 0, \quad \sigma_2(a+) \neq 0, \quad \sigma_2(b-) \neq 0. \end{aligned}$$

Assume, moreover, that for every  $0 < \eta < \min\{\sigma_2(t) : a \leq t \leq b\}$  there exist  $\gamma \in ]a, b[$  and functions  $p_\eta, q_\eta \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$  such that

$$(1.5) \quad \int_a^b q_\eta(s) \, ds < +\infty, \quad \int_a^b (s-a)(b-s)p_\eta(s) \, ds < +\infty$$

and

$$\begin{aligned} |f(t, x)| &\leq p_\eta(t), \quad g(t, x) \operatorname{sgn}(\gamma - t) \geq -q_\eta(t) \\ &\text{for } a < t < b, \quad \sigma_{1\eta}(t) \leq x \leq \sigma_2(t), \end{aligned}$$

where

$$(1.6) \quad \sigma_{1\eta}(\cdot) = \max\{\eta, \sigma_1(\cdot)\}.$$

Then the problem (1.1), (1.2) has at least one solution  $u$  such that

$$(1.7) \quad \sigma_1(t) \leq u(t) \leq \sigma_2(t) \quad \text{for } a < t < b.$$

**Remark 1.1.** Theorem 1.1 covers the case when both the functions  $f$  and  $g$  have nonintegrable singularities with respect to the independent variable. Note that singularities of the function  $g$  may be “sufficiently large”. As an example, consider the problem

$$(1.8) \quad \begin{aligned} u'' &= \frac{\lambda}{t(1-t)} - \frac{\lambda}{t(1-t)u^3} + \left(1 + \frac{1-2t}{t^2(1-t)^2}\right) \frac{u'}{u^2}, \\ u(0+) &= 0, \quad u(1-) = 0, \end{aligned}$$

where  $\lambda > 0$ . We can easily see that  $\sigma_2(t) \equiv 1$  is an upper function of the equation, and  $\sigma_1(t) = \varepsilon t(1-t)$  for  $0 \leq t \leq 1$ , where  $0 < \varepsilon < \frac{\lambda}{2+\lambda}$  is a lower function. Putting now  $\gamma = \frac{1}{2}$ , by Theorem 1.1 problem (1.8) has at least one solution.

**Corollary 1.1.** Let a function  $f$  be nondecreasing in the second argument and let there exist  $r > 0$  such that

$$(1.9) \quad f(t, r) \leq 0 \quad \text{for } a < t < b,$$

with the strict inequality on the subset of  $]a, b[$  of a positive measure. Let, moreover, on the set  $]a, b[ \times ]0, +\infty[$  the inequality

$$(1.10) \quad |g(t, x)| \leq q^*(t)$$

hold, where  $q^* \in L(]a, b[; \mathbb{R}_+)$ . Then the condition

$$(1.11) \quad \int_a^b (s-a)(b-s)|f(s, x)| \, ds < +\infty \quad \text{for } 0 < x \leq r$$

is necessary and sufficient for the solvability of problem (1.1), (1.2).

**Remark 1.2.** In the case when the function  $g$  does not depend on the second argument, problem (1.1), (1.2) under the conditions of Corollary 1.1 is uniquely solvable. Note also that for  $g(t, x) \equiv 0$  the above corollary implies Theorem 1.2 in [19] (see also [17], Theorem 4.3<sub>1</sub>).

**Corollary 1.2.** *Let a function  $f$  be nondecreasing in the second argument and there exist  $r > 0$  such that conditions (1.9) are fulfilled. Let, moreover, there exist  $c \in ]a, b[$  such that the mapping  $(t, x) \mapsto g(t, x) \operatorname{sgn}(c - t)$  is nondecreasing in the second argument, and*

$$\int_a^b |g(s, x)| \, ds < +\infty \quad \text{for } x > 0.$$

Then condition (1.11) is necessary and sufficient for the solvability of problem (1.1), (1.2).

As an example, consider the equation

$$(1.12) \quad u'' = h(t) \left( \delta u^\alpha - \frac{1}{u^\lambda} \right) + \frac{g(t)}{u^\mu} u' - \varphi(t),$$

where  $\delta \geq 0$ ,  $\alpha > 0$ ,  $\lambda > 0$ ,  $\mu > 0$ ,  $h, \varphi \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$ ,  $h(t) \not\equiv 0$ ,  $c \in ]a, b[$ , and

$$g(t) \operatorname{sgn}(c - t) \leq 0 \quad \text{for } a < t < b, \quad \int_a^b |g(s)| \, ds < +\infty.$$

Then by Corollary 1.2, for the solvability of the problem (1.12), (1.2) it is necessary and sufficient to have

$$(1.13) \quad \int_a^b (s - a)(b - s)h(s) \, ds < +\infty \quad \text{and} \quad \int_a^b (s - a)(b - s)\varphi(s) \, ds < +\infty.$$

Consider now the equation

$$(1.14) \quad u'' = \frac{h(t)}{u},$$

where  $h \in L_{\text{loc}}(]a, b[; \mathbb{R})$  can, in general, change its sign.

**Corollary 1.3.** *Let the function  $h$  admit the representation*

$$h(t) = p(t) - q(t), \quad p(t) \geq 0, \quad q(t) \geq 0 \quad \text{for } a < t < b,$$

where

$$\int_a^b \frac{p(s)}{(s-a)(b-s)} ds < +\infty, \quad \int_a^b (s-a)(b-s)q(s) ds < +\infty$$

and

$$(1.15) \quad (b-t) \int_a^t (s-a)q(s) ds + (t-a) \int_t^b (b-s)q(s) ds \\ > (t-a)(b-t)\sqrt{b-a} \sqrt{\int_a^b (s-a)(b-s)q(s) ds} \\ \times \exp\left[\frac{2}{b-a} \int_a^b \frac{p(s)}{(s-a)(b-s)} ds\right] \quad \text{for } a < t < b.$$

Then problem (1.14), (1.2) has at least one solution.

According to Corollary 1.3, for example the problem

$$u'' = \left( \frac{kt(1-t)}{\sqrt{|2t-1|}} - \lambda \right) \frac{1}{u}; \quad u(0+) = 0, \quad u(1-) = 0,$$

where  $k > 0$  and  $\lambda > \frac{2}{3}e^{8k}$ , has at least one solution.

**Theorem 1.2.** Let  $\sigma_1$  and  $\sigma_2$  be respectively lower and upper functions of equation (1.1) satisfying conditions (1.4). Assume, moreover, that for every  $0 < \eta < \min\{\sigma_2(t) : a \leq t \leq b\}$  there exist functions  $p_\eta, q_\eta \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$  such that conditions (1.5) and

$$f(t, x) \geq -p_\eta(t), \quad |g(t, x)| \leq q_\eta(t) \quad \text{for } a < t < b, \quad \sigma_{1\eta}(t) \leq x \leq \sigma_2(t)$$

are fulfilled, where  $\sigma_{1\eta}$  is the function defined by (1.6). Then problem (1.1), (1.2) has at least one solution.

**Remark 1.3.** Theorem 1.2 covers also the case when condition (1.3) is not fulfilled for the function  $f$ . Indeed, consider the problem

$$(1.16) \quad u'' = u - \frac{1}{u^2} + \frac{\lambda(1-2t)}{u^2} u' + \frac{\alpha}{t^2(1-t)^2}; \quad u(0+) = 0, \quad u(1-) = 0,$$

where  $\alpha > 0$  and  $\lambda \in \mathbb{R}$ . It is easily seen that  $\sigma_1(t) = \varepsilon t(1-t)$ , where  $0 < \varepsilon < \frac{1}{1+\alpha+|\lambda|}$ , is a lower function and  $\sigma_2(t) \equiv 1$  is an upper function. Consequently, by Theorem 1.2, problem (1.16) is solvable. It should also be noted that in the case  $\lambda \leq 0$  and  $\alpha < 0$ , problem (1.16) by Corollary 1.2 has no solution.

From Theorem 1.2, for the equation

$$(1.17) \quad u'' = h(t)u^\lambda - \frac{p(t)}{u^\mu} + \varphi(t),$$

where  $\lambda > 0$  and  $\mu > 0$ , we obtain the following

**Corollary 1.4.** *Let  $r > 0$ ,  $n > 0$ ,  $p_0 > 0$ ,  $p, h, \varphi \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$ , and*

$$\int_a^b (s-a)(b-s)p(s) ds < +\infty,$$

$$p_0 \leq p(t), \quad (\varphi(t) + h(t))[(t-a)(b-t)]^n \leq r \quad \text{for } a < t < b.$$

*Then problem (1.17), (1.2) is uniquely solvable.*

As is readily seen from this corollary, the functions  $h$  and  $\varphi$  need not satisfy conditions (1.13).

## 2. SOME AUXILIARY PROPOSITIONS

In this section, lemmas on a priori estimates and a lemma on the solvability of problem (1.1), (1.2) will be established in the case when  $f, g \in \text{Car}_{\text{loc}}(]a, b[ \times \mathbb{R}; \mathbb{R})$ . Everywhere in what follows, functions  $h_1, h_2 \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$  will be assumed to satisfy the conditions

$$(2.1) \quad \int_a^b (s-a)(b-s)h_1(s) ds < +\infty, \quad \int_a^b h_2(s) ds < +\infty.$$

**Lemma 2.1.** *Let  $r_0 > 0$ , and let  $h_1, h_2 \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$  satisfy conditions (2.1). Then there exist  $c_0 > 0$  and functions  $H_1 \in C([a, \frac{a+b}{2}]; \mathbb{R}_+)$ ,  $H_2 \in C([\frac{a+b}{2}, b]; \mathbb{R}_+)$  satisfying the conditions  $H_1(a) = 0$ ,  $H_2(b) = 0$  and such that for any  $a_1 \in ]a, \frac{a+b}{2}[$ ,  $b_1 \in ]\frac{a+b}{2}, b[$  and  $u \in AC'([a_1, b_1]; \mathbb{R})$  satisfying the inequalities*

$$(2.2) \quad \begin{aligned} u''(t) &\geq -h_1(t) - h_2(t)|u'(t)| && \text{for } a_1 < t < b_1, \\ |u(t)| &\leq r_0 && \text{for } a_1 < t < b_1, \end{aligned}$$

*the following estimates hold:*

$$(2.3) \quad (t - a_1)(b_1 - t)|u'(t)| \leq c_0 \quad \text{for } a_1 < t < b_1,$$

$$(2.4) \quad \begin{aligned} u(t) &\leq u(a_1) + H_1(t) && \text{for } a_1 \leq t \leq \frac{a+b}{2}, \\ u(t) &\leq u(b_1) + H_2(t) && \text{for } \frac{a+b}{2} \leq t \leq b_1. \end{aligned}$$



*P r o o f.* Let a function  $u \in AC'([a_1, b_1]; \mathbb{R})$  satisfy the conditions of the lemma. Suppose

$$\psi_0(t) = -h_2(t) \operatorname{sgn} u'(t), \quad \psi_1(t) = u''(t) + h_2(t)|u'(t)| \quad \text{for } a_1 < t < b_1.$$

Clearly,  $u$  is a solution of the equation

$$(2.5) \quad u'' = \psi_0(t)u' + \psi_1(t)$$

and

$$(2.6) \quad \psi_1(t) \geq -h_1(t) \quad \text{for } a_1 < t < b_1.$$

Let  $t_0 \in ]a_1, b_1[$  be an arbitrary point such that  $u'(t_0) \neq 0$ . Then either

$$(2.7) \quad u'(t_0) > 0$$

or

$$(2.8) \quad u'(t_0) < 0.$$

Suppose that (2.7) ((2.8)) is satisfied. Put

$$\begin{aligned} \mu(t) &= \int_t^{b_1} \exp \left[ \int_t^s \psi_0(\xi) d\xi \right] ds \quad \text{for } t_0 < t < b_1 \\ \left( \mu(t) &= \int_{a_1}^t \exp \left[ \int_t^s \psi_0(\xi) d\xi \right] ds \quad \text{for } a_1 < t < t_0 \right). \end{aligned}$$

Multiplying both sides of (2.5) by  $\mu$  and integrating from  $t_0$  to  $b_1$  (from  $a_1$  to  $t_0$ ), we obtain

$$\begin{aligned} -\mu(t_0)u'(t_0) - u(b_1) + u(t_0) &= \int_{t_0}^{b_1} \mu(s)\psi_1(s) ds \\ \left( \mu(t_0)u'(t_0) + u(t_0) - u(a_1) &= \int_{a_1}^{t_0} \mu(s)\psi_1(s) ds \right). \end{aligned}$$

Hence by (2.2), (2.6) and (2.7) ((2.8)) we get

$$(2.9) \quad \begin{aligned} \mu(t_0)|u'(t_0)| &\leq 2r_0 + \int_{t_0}^{b_1} h_1(s)\mu(s) ds \\ \left( \mu(t_0)|u'(t_0)| &\leq 2r_0 + \int_{a_1}^{t_0} h_1(s)\mu(s) ds \right). \end{aligned}$$

We can easily check that

$$\begin{aligned} h_0^{-1}(b_1 - t) &\leq \mu(t) \leq h_0(b_1 - t) & \text{for } t_0 < t < t_1 \\ (h_0^{-1}(t - a_1) &\leq \mu(t) \leq h_0(t - a_1) & \text{for } a_1 < t < t_0), \end{aligned}$$

where

$$(2.10) \quad h_0 = \exp \left[ \int_a^b h_2(s) \, ds \right].$$

Taking this into account, from (2.9) we find that inequality (2.3), where

$$(2.11) \quad c_0 = \left[ 2r_0(b - a) + \int_a^b (s - a)(b - s)h_1(s) \, ds \right] \exp \left[ 2 \int_a^b h_2(s) \, ds \right],$$

is fulfilled. □

Let us now show that estimates (2.4) are satisfied, where

$$(2.12) \quad \begin{aligned} H_1(t) &= \left[ \frac{4r_0}{b - a}(t - a) + \int_a^t (s - a)h_1(s) \, ds \right. \\ &\quad \left. + (t - a) \int_t^{\frac{a+b}{2}} h_1(s) \, ds \right] h_0^2 \quad \text{for } a < t \leq \frac{a+b}{2}, \\ H_2(t) &= \left[ \frac{4r_0}{b - a}(b - t) + \int_t^b (b - s)h_1(s) \, ds \right. \\ &\quad \left. + (b - t) \int_{\frac{a+b}{2}}^t h_1(s) \, ds \right] h_0^2 \quad \text{for } \frac{a+b}{2} \leq t < b. \end{aligned}$$

Here the number  $h_0$  is defined by (2.10).

Let us denote by  $w_1$  a solution of the boundary value problem

$$w'' = \psi_0(t)w' - h_1(t) \quad w(a_1) = 0, \quad w\left(\frac{a+b}{2}\right) = 2r_0$$

and show that

$$(2.13) \quad \tilde{w}(t) = u(t) - u(a_1) - w_1(t) \leq 0 \quad \text{for } a_1 < t_0 < \frac{a+b}{2}.$$

Assume the contrary. Let (2.13) be violated. Then there exist  $t_* \in [a_1, \frac{a+b}{2}[$  and  $t^* \in ]t_*, \frac{a+b}{2}]$  such that

$$(2.14) \quad \tilde{w}(t) > 0 \quad \text{for } t_* < t < t^*, \quad \tilde{w}(t_*) = 0 \quad \text{and} \quad \tilde{w}(t^*) = 0.$$

Taking into consideration (2.5) and (2.6), we obtain

$$\tilde{w}''(t) \geq \psi_0(t)\tilde{w}'(t) \quad \text{for } t_* < t < t^*.$$

However, this contradicts condition (2.14). Thus (2.13) is valid.

We can directly verify that

$$w_1(t) \leq H_1(t) \quad \text{for } a_1 \leq t \leq \frac{a+b}{2},$$

where  $H_1$  is the function defined by (2.12). Consequently, the first of inequalities (2.4) is valid.

Analogously one can prove that

$$u(t) \leq u(b_1) + w_2(t), \quad w_2(t) \leq H_2(t) \quad \text{for } \frac{a+b}{2} \leq t \leq b_1,$$

where  $w_2$  is a solution of the boundary value problem

$$w'' = \psi_0(t)w' - h_1(t); \quad w\left(\frac{a+b}{2}\right) = 2r_0, \quad w(b_1) = 0.$$

□

**Lemma 2.2.** *Let  $r_0 > 0$ ,  $\alpha \in ]a, \frac{a+b}{2}[$ ,  $\beta \in ]\frac{a+b}{2}, b[$ ,  $\gamma \in ]\alpha, \beta[$ , and let  $h_1, h_2 \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$  satisfy conditions (2.1). Then there exists a function  $\varphi \in L(]a, b[; \mathbb{R}_+)$  such that  $\varphi$  is bounded in  $]a, b[$ , and for any  $a_1 \in ]a, \alpha[$ ,  $b_1 \in ]\beta, b[$ , and a function  $u \in AC'([a_1, b_1]; \mathbb{R})$  satisfying condition (2.2) and inequalities*

$$(2.15) \quad u''(t) \operatorname{sgn}((\gamma - t)u'(t)) \geq -h_1(t) - h_2(t)|u'(t)| \quad \text{for } a_1 < t < b_1,$$

$$(2.16) \quad u''(t) \geq -h_1(t) - h_2(t)|u'(t)| \quad \text{for } \alpha < t < \beta,$$

the estimate

$$(2.17) \quad |u'(t)| \leq \varphi(t) \quad \text{for } a_1 < t < b_1$$

holds.

*Proof.* Let  $u \in AC'([a_1, b_1]; \mathbb{R})$  satisfy the conditions of the lemma. By virtue of (2.16), (2.2) and Lemma 2.1, the estimate

$$(2.18) \quad |u'(\gamma)| \leq \frac{c_0}{(\gamma - \alpha)(\beta - \gamma)}$$

is fulfilled, where  $c_0$  is the number given by (2.11). Let us show that

$$(2.19) \quad |u'(t)| \leq \varphi_0(t) \quad \text{for } a_1 < t < \gamma,$$

where

$$(2.20) \quad \varphi_0(t) = \left( \frac{c_0}{(\gamma - \alpha)(\beta - \gamma)} + \int_t^\gamma h_1(s) ds \right) \exp \left[ \int_a^b h_2(s) ds \right] \quad \text{for } a < t < \gamma.$$

Assume the contrary. Let (2.19) be violated. Then by (2.18) and (2.20), there exist  $t_* \in ]a_1, \gamma[$  and  $t^* \in ]t_*, \gamma[$  such that

$$(2.21) \quad \begin{aligned} |u'(t)| &> \varphi_0(t) \quad \text{for } t_* \leq t < t^*, \\ |u'(t^*)| &= \frac{c_0}{(\gamma - \alpha)(\beta - \gamma)}. \end{aligned}$$

For the sake of definiteness we assume that  $u'(t) > 0$  for  $t_* < t < t^*$ . Then by virtue of the lemma on the differential inequality, from (2.15) we find

$$|u'(t_*)| \leq \left( |u'(t^*)| + \int_{t_*}^{t^*} \exp \left[ \int_{t_*}^s h_2(\xi) d\xi \right] h_1(s) ds \right) \leq \varphi_0(t_*).$$

However, the last inequality contradicts condition (2.21). Consequently, estimate (2.19) is valid.

Analogously we can show that

$$(2.22) \quad |u'(t)| \leq \varphi_1(t) \quad \text{for } \gamma \leq t < b_1,$$

where

$$\varphi_1(t) = \left( \frac{c_0}{(\gamma - \alpha)(\beta - \gamma)} + \int_\gamma^t h_1(s) ds \right) \exp \left[ \int_a^b h_2(s) ds \right] \quad \text{for } \gamma < t < b.$$

Suppose now

$$\varphi(t) = \begin{cases} \varphi_0(t) & \text{for } a < t \leq \gamma \\ \varphi_1(t) & \text{for } \gamma < t < b \end{cases}.$$

It is evident from (2.19) and (2.22) that estimate (2.17) is satisfied. It is not also difficult to see that the function  $\varphi$  is bounded in  $]a, b[$ , and  $\varphi \in L(]a, b[; \mathbb{R}_+)$ .  $\square$

**Lemma 2.3.** Let  $f, g \in \text{Car}_{\text{loc}}(]a, b[ \times \mathbb{R}; \mathbb{R})$ , and let  $\sigma_1$  and  $\sigma_2$  be lower and upper functions of equation (1.1) satisfying conditions (1.4). Moreover, let there exist functions  $p_0, q_0 \in L_{\text{loc}}(]a, b[; \mathbb{R}_+)$  such that

$$\int_a^b (s-a)(b-s)p_0(s) ds < +\infty, \quad \int_a^b q_0(s) ds < +\infty$$

and

$$(2.23) \quad f(t, x) \geq -p_0(t), \quad |g(t, x)| \leq q_0(t) \text{ for } a < t < b, \quad \sigma_1(t) \leq x \leq \sigma_2(t),$$

$$(2.24) \quad (|f(t, x)| \leq p_0(t), \quad g(t, x) \operatorname{sgn}(c-t) \geq -q_0(t) \text{ for } a < t < b, \quad \sigma_1(t) \leq x \leq \sigma_2(t)).$$

Then problem (1.1), (1.2) has at least one solution  $u$  satisfying (1.7).

**P r o o f.** Let conditions (2.23) be fulfilled. Choose sequences  $(t_{ik})_{k=1}^{+\infty}$ ,  $(s_{ik})_{k=1}^{+\infty}$ , and  $(c_{ik})_{k=1}^{+\infty}$  ( $i = 1, 2$ ) such that

$$a < t_{1k+1} < t_{1k} < \frac{a+b}{2} < t_{2k} < t_{2k+1} < b, \quad k = 1, 2, 3, \dots,$$

$$(2.25) \quad s_{1k+1} \in ]t_{1k+1}, t_{1k}[, \quad s_{2k+1} \in ]t_{2k}, t_{2k+1}[, \quad t_{11} < s_{11} < \frac{a+b}{2} < s_{21} < t_{21},$$

$$(2.26) \quad c_{1k} = \sigma_1(t_{1k}), \quad c_{2k} = \sigma_1(t_{2k}), \quad k = 1, 2, 3, \dots,$$

$$(2.26) \quad \lim_{k \rightarrow +\infty} t_{1k} = a, \quad \lim_{k \rightarrow +\infty} t_{2k} = b.$$

Let  $r_0 = \max\{|\sigma_1(t)| + |\sigma_2(t)| : a \leq t \leq b\}$  and let  $c_0$  be the number defined by (2.11), where  $h_1(t) = p_0(t)$ ,  $h_2(t) = q_0(t)$  for  $a < t < b$ ,

$$(2.27) \quad \tilde{\varphi}(t) = \begin{cases} \frac{c_0}{(t-t_{1k+1})(t_{2k+1}-t)} & \text{for } t \in ]s_{1k+1}, s_{1k}[ \cup ]s_{2k}, s_{2k+1}[, \\ \frac{c_0}{(t-t_{11})(t_{21}-t)} & \text{for } t \in ]s_{11}, s_{21}[, \end{cases}$$

$$(2.28) \quad \varrho(t) = \tilde{\varphi}(t) + |\sigma_1'(t)| + |\sigma_2'(t)| + 1 \text{ for } a < t < b,$$

$$F(t, x, y) = f(t, x) + g(t, x)y \text{ for } a < t < b, \quad x, y \in \mathbb{R},$$

$$(2.29) \quad \tilde{F}(t, x, y) = \begin{cases} F(t, x, y) & \text{for } |y| \leq \varrho(t), \\ \left(2 - \frac{|y|}{\varrho(t)}\right) F(t, x, y) & \text{for } \varrho(t) < |y| \leq 2\varrho(t), \\ 0 & \text{for } |y| \geq 2\varrho(t), \end{cases}$$

$$\begin{aligned}
q(t) &= |g(t, \sigma_1(t))| + |g(t, \sigma_2(t))| \quad \text{for } a < t < b, \\
(2.30) \quad F_0(t, x, y) &= \begin{cases} \tilde{F}(t, \sigma_1(t), y) + \frac{\sigma_1(t) - x}{1 + |\sigma_1(t) - x|} q(t) & \text{for } x \leq \sigma_1(t), \\ \tilde{F}(t, x, y) & \text{for } \sigma_1(t) < x < \sigma_2(t), \\ \tilde{F}(t, \sigma_2(t), y) + \frac{x - \sigma_2(t)}{1 + |x - \sigma_2(t)|} q(t) & \text{for } x \geq \sigma_2(t). \end{cases}
\end{aligned}$$

We can easily see that  $F_0 \in \text{Car}([t_{1k}, t_{2k}[ \times \mathbb{R}^2; \mathbb{R})$  and for any natural  $k$  there exists a function  $F_k^* \in L([t_{1k}, t_{2k}[; \mathbb{R}_+)$  such that

$$|F_0(t, x, y)| \leq F_k^*(t) \quad \text{for } t_{1k} < t < t_{2k}, \quad x, y \in \mathbb{R}.$$

Therefore the boundary value problem

$$(2.31) \quad u'' = F_0(t, u, u'),$$

$$(2.32) \quad u(t_{1k}) = c_{1k}, \quad u(t_{2k}) = c_{2k}$$

has at least one solution  $u_k$  (cf., for example, Lemma 2.1 in [17]).

We will show that

$$(2.33) \quad v(t) = \sigma_1(t) - u_k(t) \leq 0 \quad \text{for } t_{1k} \leq t \leq t_{2k}.$$

Assume on the contrary that for some  $\tilde{t} \in ]t_{ik}, t_{2k}[$  the inequality  $v(\tilde{t}) > 0$  holds. Since  $v(t_{ik}) = 0$ ,  $i = 1, 2$ , there exist  $t_* \in [t_*, t_{2k}[$  and  $t^* \in ]t_k, t_{2k}[$  such that  $v \in AC'([t_*, t^*]; \mathbb{R})$ ,

$$(2.34) \quad v(t) > v(t_*) > 0 \quad \text{for } t_* < t < t^*, \quad v(t_*) = v(t^*),$$

$$(2.35) \quad |v'(t)| < \frac{v(t)}{1 + v(t)} \quad \text{for } t_* < t < t^*.$$

Then

$$(2.36) \quad |u'_k(t)| < \varrho(t) \quad \text{for } t_* < t < t^*.$$

In view of (2.29), (2.30) and (2.34)–(2.36), the inequality

$$\begin{aligned}
v''(t) &\geq F(t, \sigma_1(t), \sigma'_1(t)) - F(t, \sigma_1(t), u'_k(t)) + q(t) \frac{v(t)}{1 + v(t)} \\
&= g(t, \sigma_1(t))v'(t) + q(t) \frac{v(t)}{1 + v(t)} \geq 0
\end{aligned}$$

is satisfied almost everywhere in  $]t_*, t^*[$ . Since  $v \in AC'([t_*, t^*]; \mathbb{R})$ , the above inequality contradicts conditions (2.34). Consequently, inequality (2.33) holds. Analogously we can see that  $u_k(t) \leq \sigma_2(t)$  for  $t_{1k} \leq t \leq t_{2k}$ . Thus

$$(2.37) \quad \sigma_1(t) \leq u_k(t) \leq \sigma_2(t) \quad \text{for } t_{1k} \leq t \leq t_{2k}.$$

Taking this into account, by (2.30) we have

$$(2.38) \quad u_k''(t) = \widetilde{F}(t, u_k(t), u_k'(t)) \quad \text{for } t_{1k} \leq t \leq t_{2k},$$

whence in view of condition (2.23) we conclude that the function  $u_k$  satisfies the conditions of Lemma 2.1 for  $a_1 = t_{1k}$  and  $b_1 = t_{2k}$ . Hence,

$$(2.39) \quad (t - t_{1k})(t_{2k} - t)|u_k'(t)| \leq c_0 \quad \text{for } t_{1k} < t < t_{2k}$$

and

$$(2.40) \quad \begin{aligned} u_k(t) &\leq c_{1k} + H_1(t) \quad \text{for } t_{1k} < t < \frac{a+b}{2}, \\ u_k(t) &\leq c_{2k} + H_2(t) \quad \text{for } \frac{a+b}{2} \leq t \leq t_{2k}, \end{aligned}$$

where  $H_1$  and  $H_2$  are the functions appearing in Lemma 2.1. Due to (2.27)–(2.29) and (2.39), from (2.38) we find

$$(2.41) \quad u_k''(t) = F(t, u_k(t), u_k'(t)) \quad \text{for } s_{1k} < t < s_{2k}.$$

Now (2.37), (2.39) and (2.41) imply uniform boundedness and equicontinuity of sequences  $(u_k)_{k=1}^{+\infty}$  and  $(u_k')_{k=1}^{+\infty}$  in  $]a, b[$  (i.e., on each compact contained in  $]a, b[$ ). Therefore without loss of generality we assume that

$$\lim_{k \rightarrow +\infty} u_k(t) = u_0(t), \quad \lim_{k \rightarrow +\infty} u_k'(t) = u_0'(t)$$

uniformly in  $]a, b[$ . Clearly,  $u_0$  is a solution of equation (1.1). Moreover, from (1.4), (2.25), (2.37) and (2.40) we conclude that  $u_0$  satisfies boundary conditions (1.2) and inequalities (1.7).

In the case when (2.24) is satisfied, the lemma can be proved analogously. The only difference is that there is no need to introduce sequences  $(s_{ik})_{k=1}^{+\infty}$ ,  $i = 1, 2$ , but we must put  $\widetilde{\varphi}(t) = \varphi(t)$  for  $a < t < b$ , where  $\varphi$  is the function appearing in Lemma 2.2 for the case  $\alpha = \frac{a+\gamma}{2}$ ,  $\beta = \frac{b+\gamma}{2}$ , and  $h_1(t) = p_0(t)$ ,  $h_2(t) = q_0(t) + \widetilde{q}(t)$ . Here  $\widetilde{q}(t) = 0$  for  $t \in ]a, \alpha[ \cup ]\beta, b[$ , and  $\widetilde{q}(t) = \max\{|g(t, x)|: \sigma_* \leq x \leq \sigma^*\}$  for  $\alpha < t < \beta$ , where  $\sigma_* = \min\{\sigma_1(t): \alpha \leq t \leq \beta\}$ ,  $\sigma^* = \max\{\sigma_2(t): \alpha \leq t \leq \beta\}$ .  $\square$

Finally, for the sake of convenience we give without proof some lemmas on properties of solutions of the linear equation

$$(2.42) \quad u'' = p(t)u + q(t)u',$$

where  $p, q \in L_{\text{loc}}([a, b]; \mathbb{R})$ .

**Definition 2.1.** Equation (2.42) is said to be oscillatory on the segment  $[a, b]$  if every its nontrivial solution has at least one zero in the interval  $]a, b[$ .

**Lemma 2.4.** *Let*

$$(2.43) \quad \begin{aligned} p(t) \leq 0 \quad \text{for } a < t < b, \\ \int_a^b |q(s)| ds < +\infty, \quad \int_a^b (s-a)(b-s)|p(s)| ds < +\infty, \end{aligned}$$

and

$$\int_a^b (s-a)^2(b-s)^2|p(s)| ds \geq (b-a)^3 \exp \left[ 8 \int_a^b |q(s)| ds \right].$$

Then equation (2.42) is oscillatory.

**Lemma 2.5.** *Let (2.43) be fulfilled and*

$$\begin{aligned} \int_a^b |q(s)| ds < +\infty, \quad \int_a^b (s-a)|p(s)| ds < +\infty \quad \left( \int_a^b (b-s)|p(s)| ds < +\infty \right), \\ \int_a^b (s-a)|p(s)| ds < (b-a) \exp \left[ -3 \int_a^b |q(s)| ds \right] \\ \left( \int_a^b (b-s)|p(s)| ds < (b-a) \exp \left[ -3 \int_a^b |q(s)| ds \right] \right). \end{aligned}$$

Then equation (2.42) has a solution  $v_1$  (a solution  $v_2$ ) satisfying the conditions

$$\begin{aligned} v_1'(t) > 0 \quad \text{for } a < t \leq b, \quad v_1(a+) = 1 \\ (v_2'(t) < 0 \quad \text{for } a \leq t < b, \quad v_2(b-) = 1). \end{aligned}$$

The above two lemmas immediately follow from the results obtained in [20].

**Lemma 2.6.** *Let  $p(t) \geq 0$  for  $a < t < b$ , and  $\int_a^b (s-a)(b-s)p(s) ds < +\infty$ . Then Green's function  $G$  of the problem*

$$u'' = p(t)u; \quad u(0+) = 0, \quad u(b-) = 0$$



admits the estimates (cf., for example, [16]).

$$\begin{aligned} \frac{1}{\mu(b-a)}(t-a)(b-\tau) &\leq -G(t, \tau) \leq \frac{\mu^2}{b-a}(t-a)(b-\tau) \\ &\text{for } a \leq t < \tau \leq b, \\ \frac{1}{\mu(b-a)}(\tau-a)(b-t) &\leq -G(t, \tau) \leq \frac{\mu^2}{b-a}(\tau-a)(b-t) \\ &\text{for } a \leq \tau \leq t \leq b, \end{aligned}$$

where  $\mu = \exp[(b-a)^{-1} \int_a^b (s-a)(b-s)p(s) ds]$ .

### 3. PROOF OF MAIN RESULTS

Proof of Theorem 1.1. Let

$$\sigma_* = \min\{\sigma_2(t) : a \leq t \leq b\}, \quad \sigma^* = \max\{\sigma_2(t) : a \leq t \leq b\}, \quad \varepsilon = \frac{\sigma_*}{2}.$$

Choose  $a_0 \in ]a, \gamma[$  and  $b_0 \in ]\gamma, b[$  such that  $\sigma_1(t) < \varepsilon$  for  $t \in ]a, a_0[ \cup ]b_0, b[$ . Denote by  $v_1$  and  $v_2$  respectively solutions of the boundary value problems

$$\begin{aligned} v'' &= -q_\varepsilon(t)v' - p_\varepsilon(t); & v(a+) &= \varepsilon, & v(a_0) &= \sigma^* + 1, \\ v'' &= q_\varepsilon(t)v' - p_\varepsilon(t); & v(b_0) &= \sigma^* + 1, & v(b-) &= \varepsilon. \end{aligned}$$

Obviously, there exist  $a_1 \in ]a, a_0[$  and  $b_1 \in ]b_0, b[$  such that

$$\begin{aligned} \varepsilon &< v_1(t) < \sigma_2(t), & v_1'(t) &> 0 & \text{for } a < t < a_1, \\ v_1(a_1) &= \sigma_2(a_1), & v_1'(a_1) &\geq \sigma_2'(a_1+), \\ \varepsilon &< v_2(t) < \sigma_2(t), & v_2'(t) &< 0 & \text{for } b_1 < t < b, \\ v_2(b_1) &= \sigma_2(b_1), & v_2'(b_1) &\leq \sigma_2'(b_1-). \end{aligned}$$

Taking this and the conditions of the theorem into account, we conclude that

$$\begin{aligned} v_1''(t) &\leq -[f(t, v_1(t)) + g(t, v_1(t))v_1'(t)]_- & \text{for } a < t < a_1, \\ v_2''(t) &\leq -[f(t, v_2(t)) + g(t, v_2(t))v_2'(t)]_- & \text{for } b_1 < t < b. \end{aligned}$$

Choose sequences  $(t_{1k})_{k=1}^{+\infty}$  and  $(t_{2k})_{k=1}^{+\infty}$  such that  $t_{11} \in ]a, a_1[$ ,  $t_{21} \in ]b_1, b[$ ,  $t_{1k+1} < t_{1k}$ ,  $t_{2k} < t_{2k+1}$ ,  $k = 1, 2, \dots$ ,

$$\begin{aligned} \sigma_1(t) &< \sigma_1(t_{1k}) & \text{for } a < t < t_{1k}, & \sigma_1(t) < \sigma_1(t_{2k}) & \text{for } t_{2k} < t < b, \\ \sigma_1(t_{1k+1}) &< \sigma_1(t_{1k}), & \sigma_1'(t_{1k}) &\geq 0, & \sigma_1(t_{2k+1}) < \sigma_2(t_{2k}), & \sigma_1'(t_{2k}) \leq 0, \\ \sigma_1(t_{1k}) &= \sigma_1(t_{2k}), & \lim_{k \rightarrow +\infty} t_{1k} &= a, & \lim_{k \rightarrow +\infty} t_{2k} &= b. \end{aligned}$$

Suppose

$$\begin{aligned}
 \chi_k(x) &= \begin{cases} \sigma_1(t_{1k}) & \text{for } x < \sigma_1(t_{1k}) \\ x & \text{for } x \geq \sigma_1(t_{1k}) \end{cases} \quad k = 1, 2, \dots, \\
 F_k(t, x, y) &= \begin{cases} -[f(t, \chi_k(x)) + g(t, \chi_k(x))y]_- & \text{for } t \in ]a, t_{1k}[ \cup ]t_{2k}, b[, \\ f(t, \chi_k(x)) + g(t, \chi_k(x))y & \text{for } t \in ]t_{1k}, t_{2k}[, \end{cases} \\
 (3.1) \quad \tilde{\sigma}_{1k}(t) &= \begin{cases} \sigma_1(t_{1k}) & \text{for } a \leq t \leq t_{1k}, \\ \sigma_1(t) & \text{for } t_{1k} < t < t_{2k}, \\ \sigma_1(t_{2k}) & \text{for } t_{2k} \leq t \leq b, \end{cases} \\
 \tilde{\sigma}_2(t) &= \begin{cases} v_1(t) & \text{for } a \leq t \leq a_1, \\ \sigma_2(t) & \text{for } a_1 < t < b_1, \\ v_2(t) & \text{for } b_1 \leq t \leq b, \end{cases}
 \end{aligned}$$

and consider the boundary value problem

$$(3.2_k) \quad u'' = F_k(t, u, u'),$$

$$(3.3_k) \quad u(a+) = \sigma_1(t_{1k}), \quad u(b-) = \sigma_1(t_{2k}).$$

It is easy to see that  $F_k \in \text{Car}_{\text{loc}}(]a, b[ \times \mathbb{R}^2; \mathbb{R})$ , and  $\tilde{\sigma}_{11}$  and  $\tilde{\sigma}_2$  are respectively lower and upper functions of equation (3.2<sub>1</sub>). Therefore, by Lemma 2.3, problem (3.2<sub>1</sub>), (3.3<sub>1</sub>) has at least one solution  $u_1$  satisfying

$$\tilde{\sigma}_{11}(t) \leq u_1(t) \leq \tilde{\sigma}_2(t) \quad \text{for } a \leq t \leq b.$$

Further,  $\tilde{\sigma}_{12}$  is a lower and  $u_1$  is an upper function of equation (3.2<sub>2</sub>). Therefore, by Lemma 2.3, problem (3.2<sub>2</sub>), (3.3<sub>2</sub>) has a solution  $u_2$  satisfying the condition

$$\tilde{\sigma}_{12}(t) \leq u_2(t) \leq u_1(t) \quad \text{for } a \leq t \leq b.$$

Continuing this process, we obtain a sequence of functions  $(u_k)_{k=1}^{+\infty}$  satisfying equation (3.2<sub>k</sub>), conditions (3.3<sub>k</sub>), and

$$(3.4) \quad \tilde{\sigma}_{1k+1}(t) \leq u_{k+1}(t) \leq u_k(t) \quad \text{for } a \leq t \leq b, \quad k = 1, 2, \dots$$

Applying (3.1), (3.4) and Lemma 2.2 (in the case,  $r_0 = \sigma^*$ ,  $\alpha = a_0$ ,  $\beta = b_0$ ,  $\eta = \min\{\tilde{\sigma}_1(a_1), \tilde{\sigma}_1(b_1)\}$ ,  $h_1(t) = p_\eta(t)$ ,  $h_2(t) = q_\eta(t) + q^*(t)$ , where  $q^*(t) = \max\{|g(t, x)|: \varepsilon_0 \leq x \leq \sigma^*\}$  and  $\varepsilon_0 = \min\{\sigma_1(t): a_0 \leq t \leq b_0\}$ ), we find that sequences  $(u_k)_{k=1}^{+\infty}$

and  $(u'_k)_{k=1}^{+\infty}$  are uniformly bounded and equicontinuous in  $]a, b[$  (i.e., on each segment contained in  $]a, b[$ ). Therefore without loss of generality we can assume that

$$\lim_{k \rightarrow +\infty} u_k(t) = u_0(t), \quad \lim_{k \rightarrow +\infty} u'_k(t) = u'_0(t)$$

uniformly in  $]a, b[$ .

Clearly,  $u_0$  is a solution of equation (1.1), and

$$0 < \sigma_1(t) \leq u_0(t) \leq u_k(t) \quad \text{for } a < t < b, \quad k = 1, 2, \dots$$

This with regard to (3.3<sub>k</sub>) implies

$$\begin{aligned} 0 &\leq \lim_{k \rightarrow a+} \inf u_0(t) \leq \lim_{k \rightarrow a+} \sup u_0(t) \leq \sigma_1(t_{1k}), \quad k = 1, 2, \dots, \\ 0 &\leq \lim_{k \rightarrow b-} \inf u_0(t) \leq \lim_{k \rightarrow b-} \sup u_0(t) \leq \sigma_1(t_{2k}), \quad k = 1, 2, \dots \end{aligned}$$

Consequently,  $u_0(a+) = 0$  and  $u_0(b-) = 0$ . □

**Proof of Corollary 1.1.** First we prove the sufficiency. By Theorem 1.1, it suffices to show that there exist lower and upper functions  $\sigma_1$  and  $\sigma_2$  of equation (1.1) satisfying conditions (1.4).

Choose  $\delta \in ]0, r[$  so small that either of the equations

$$(3.5) \quad u'' = \frac{1}{\delta} f(t, r)u + q^*(t)u'$$

and

$$(3.6) \quad u'' = \frac{1}{\delta} f(t, r)u - q^*(t)u'$$

is oscillatory in the intervals  $[a, \frac{a+b}{2}]$  and  $[\frac{a+b}{2}, b]$ , respectively. For this, by Lemma 2.4, it is sufficient to assume

$$\begin{aligned} \delta &< \frac{8}{(b-a)^3} \exp \left[ -8 \int_a^b q^*(s) ds \right] \\ &\times \min \left\{ \int_a^{\frac{a+b}{2}} (s-a)^2 \left( \frac{a+b}{2} - s \right)^2 |f(s, r)| ds, \int_{\frac{a+b}{2}}^b \left( s - \frac{a+b}{2} \right)^2 (b-s)^2 |f(s, r)| ds \right\}. \end{aligned}$$

Denote by  $u_1$  and  $u_2$  the solutions of equations (3.5) and (3.6), respectively, satisfying the initial conditions

$$\begin{aligned} u_1(a+) &= 0, & u'_1(a+) &= 1, \\ u_2(b-) &= 0, & u'_2(b-) &= -1. \end{aligned}$$

(The condition  $\int_a^b (s-a)(b-s)|f(s,r)| ds < +\infty$  guarantees the existence of  $u_1$  and  $u_2$ —cf., for example, [16], [17].) Then there exist  $t_1 \in ]a, \frac{a+b}{2}[$  and  $t_2 \in ]\frac{a+b}{2}, b[$  such that

$$\begin{aligned} u'_1(t) &> 0 \quad \text{for } a < t < t_1, & u'_1(t_1) &= 0, \\ u'_2(t) &< 0 \quad \text{for } t_2 < t < b, & u'_2(t_2) &= 0. \end{aligned}$$

Let

$$\sigma_1(t) = \begin{cases} \frac{\delta}{u_1(t_1)} u_1(t) & \text{for } a \leq t \leq t_1 \\ \delta & \text{for } t_1 < t < t_2 \\ \frac{\delta}{u_2(t_2)} u_2(t) & \text{for } t_2 \leq t \leq b \end{cases}.$$

As is easily seen,  $\sigma_1 \in AC'_{\text{loc}}(]a, b[; \mathbb{R}_+)$ ,  $\sigma_1(t) \leq \delta < r$  for  $a \leq t \leq b$ , and  $\sigma_1$  is a lower function of equation (1.1).

Choose  $\mu \in ]0, r[$  so small that the equations

$$v'' = -\mu \left[ f\left(t, \frac{1}{r}\right) \right]_- v - q^*(t)v'$$

and

$$v'' = -\mu \left[ f\left(t, \frac{1}{r}\right) \right]_- v - q^*(t)v'$$

have respectively solutions  $v_1$  and  $v_2$  satisfying the conditions

$$\begin{aligned} v'_1(t) &> 0 \quad \text{for } a < t \leq \frac{a+b}{2}, & v_1(a+) &= 1, \\ v'_2(t) &< 0 \quad \text{for } \frac{a+b}{2} \leq t < b, & v_2(b-) &= 1. \end{aligned}$$

For this, by Lemma 2.5, it is sufficient to assume that

$$\begin{aligned} \mu &< \frac{b-1}{2 \exp\left[3 \int_a^b q^*(s) ds\right]} \\ &\times \min \left\{ \left( 1 + \int_a^{\frac{a+b}{2}} (s-a) \left[ f\left(s, \frac{1}{r}\right) \right]_- ds \right)^{-1}, \left( 1 + \int_{\frac{a+b}{2}}^b (b-s) \left[ f\left(s, \frac{1}{r}\right) \right]_- ds \right)^{-1} \right\}. \end{aligned}$$

Let

$$\sigma_2(t) = \begin{cases} \lambda_1 v_1(t) & \text{for } a \leq t < \frac{a+b}{2}, \\ \lambda_2 v_2(t) & \text{for } \frac{a+b}{2} \leq t < b, \end{cases}$$

where

$$\lambda_i = \frac{1}{v_i\left(\frac{a+b}{2}\right)} \left( v_1\left(\frac{a+b}{2}\right) + v_2\left(\frac{a+b}{2}\right) \right) \left( r + \frac{1}{\mu} \right), \quad i = 1, 2.$$

Clearly,  $\sigma_2$  is continuous,  $\sigma_1 \in AC'_{\text{loc}}(]a, b[\setminus\{\frac{a+b}{2}\}; \mathbb{R}_+)$ ,

$$r\sigma_2(t) > \mu\sigma_2(t) > 1, \quad \sigma_2(t) > r \quad \text{for } a \leq t \leq b, \quad \sigma_2'\left(\frac{a+b}{2}-\right) > \sigma_2'\left(\frac{a+b}{2}+\right),$$

and  $\sigma_2$  is an upper function of equation (1.1). Consequently, problem (1.1), (1.2) has at least one solution.

Now let us prove the necessity. Let  $u$  be a solution of problem (1.1), (1.2). Suppose  $0 < x \leq r$  and choose  $a_0 \in ]a, b[$  such that

$$(3.7) \quad u(t) < x \quad \text{for } a \leq t \leq a_0.$$

Since the function  $u$  is bounded in  $[a, a_0]$ , we have

$$(3.8) \quad \lim_{t \rightarrow a^+} \inf (t-a)|u'(t)| = 0.$$

Multiplying both sides of equation (1.1) by

$$q(t) = \int_a^t \exp \left[ \int_t^s g(\xi, u(\xi)) d\xi \right] ds \quad \text{for } a \leq t \leq b$$

and integrating from  $t$  to  $a_0$ , we obtain for  $t \in ]a, b[$  that

$$(3.9) \quad u'(t)q(t) = u'(a_0)q(a_0) + u(a_0) - u(t) - \int_a^{a_0} q(s)f(s, u(s)) ds.$$

Owing to (1.10), we can easily see that

$$\frac{1}{c}(t-a) \leq q(t) \leq c(t-a) \quad \text{for } a < t < b,$$

where  $c = \exp \left[ \int_a^b q^*(s) ds \right]$ . Moreover, taking into account condition (3.7) and the fact that the function  $f$  is monotone, from (3.9) we easily find

$$\begin{aligned} \int_t^{a_0} (s-a)|f(s, x)| ds &\leq c(u(t) - u(a_0) - u'(a_0)q(a_0)) \\ &\quad + (t-a)|u'(t)| \quad \text{for } a < t < a_0. \end{aligned}$$

This, due to (1.2) and (3.8), implies

$$(3.10) \quad \int_a^{a_0} (s-a)|f(s, x)| ds < +\infty.$$

Analogously we can show that  $\int_{b_0}^b (b-s)|f(s, x)| ds < +\infty$  for some  $b_0 \in ]a, b[$ . Consequently, conditions (1.11) are satisfied.  $\square$

Proof of Corollary 1.2. First we prove the sufficiency. To this end, by Theorem 1.1 it suffices to show that there exist lower and upper functions  $\sigma_1$  and  $\sigma_2$  of equation (1.1) satisfying conditions (1.4). According to Corollary 1.1, the equation

$$u'' = f(t, u)$$

has solutions  $u_1$  and  $u_2$  satisfying the conditions

$$\begin{aligned} u_1(t) > 0 & \quad \text{for } a < t < c, & u_1(a+) = 0, & u_1(c-) = 0, \\ u_2(t) > 0 & \quad \text{for } c < t < b, & u_2(c+) = 0, & u_2(b-) = 0. \end{aligned}$$

It is evident that there exist  $t_1 \in ]a, c[$  and  $t_2 \in ]c, b[$  such that

$$\begin{aligned} u_1'(t) > 0 & \quad \text{for } a < t < t_1, & u_1'(t_1) = 0, \\ u_2'(t) < 0 & \quad \text{for } t_2 < t < b, & u_2'(t_2) = 0. \end{aligned}$$

Set  $\delta = u_2(t_2)/u_1(t_1)$  and

$$\sigma_1(t) = \begin{cases} \min\{1, \delta\} \cdot u_1(t) & \text{for } a \leq t < t_1, \\ \min\{1, \delta\} \cdot u_1(t_1) & \text{for } t_1 \leq t \leq t_2, \\ \min\left\{1, \frac{1}{\delta}\right\} \cdot u_2(t) & \text{for } t_2 < t \leq b. \end{cases}$$

It is not difficult to see that  $\sigma_1$  is a lower function of equation (1.1).

Choose  $\nu > 0$  such that

$$\nu \int_a^c f(s, r) q_1(s) ds = \int_c^b f(s, r) q_2(s) ds,$$

where

$$q_1(t) = \int_a^t \exp\left[\int_c^s g(\xi, r) d\xi\right] ds, \quad q_2(t) = \int_t^b \exp\left[\int_c^s g(\xi, r) d\xi\right] ds \quad \text{for } a < t < b.$$

Suppose

$$h(t) = \begin{cases} \max\{1, \nu\} \cdot f(t, r) & \text{for } a < t < c \\ \max\left\{1, \frac{1}{\nu}\right\} \cdot f(t, r) & \text{for } c < t < b \end{cases}$$

and consider the equation

$$u'' = h(t) + g(t, r)u'.$$

We can readily see that the above equation has a solution  $\sigma_2$  satisfying the conditions

$$\sigma_2(t) > 1 + \max\{\sigma_1(t) : a \leq t \leq b\}, \quad \sigma_2'(t) \operatorname{sgn}(c-t) \geq 0 \quad \text{for } a < t < b.$$

Taking now into account the fact that the functions  $f$  and  $g \operatorname{sgn}(c-t)$  are monotone, we conclude that  $\sigma_2$  is an upper function of equation (1.1) and conditions (1.4) hold. Hence problem (1.1), (1.2) has at least one solution.

Let us now prove the necessity. Let  $u$  be a solution of problem (1.1), (1.2). Clearly, (3.8) holds. Suppose  $x \in ]0, r[$  and choose  $a_0 \in ]a, c[$  such that

$$(3.11) \quad u(t) < x, \quad u'(t) > 0 \quad \text{for } a < t < a_0.$$

Multiplying both sides of equation (1.1) by  $t-a$  and integrating from  $t$  to  $a_0$ , we get

$$\begin{aligned} & (t-a)u'(t) - (a_0-a)u'(a_0) + u(a_0) - u(t) \\ &= - \int_t^{a_0} (s-a)f(s, u(s)) \, ds + \int_t^{a_0} (s-a)g(s, u(s))u'(s) \, ds \quad \text{for } a < t < a_0. \end{aligned}$$

If we now take into consideration the fact that the functions  $f$  and  $g$  are monotone, then due to conditions (3.11) from the last equality we obtain

$$\int_t^{a_0} (s-a)|f(s, x)| \, ds \leq (t-a)u'(t) - (a_0-a)u'(a_0) + u(a_0) - u(t) \quad \text{for } a < t < a_0,$$

whence according to (1.2) and (3.8) we conclude that (3.10) is satisfied. Analogously we can see that  $\int_{b_0}^b (s-a)|f(s, x)| \, ds < +\infty$  for some  $b_0 \in ]c, b[$ . Consequently, (1.11) is valid.  $\square$

**P r o o f** of Corollary 1.3. Denote by  $v$  a solution of the problem

$$(3.12) \quad v'' = \frac{p(t)}{(t-a)^2(b-t)^2} v - \lambda q(t); \quad v(a+) = 0, \quad v(b-) = 0,$$

where

$$(3.13) \quad \lambda = \left[ \int_a^b (s-a)(b-s)q(s) \, ds \right]^{-\frac{1}{2}} \exp \left[ \frac{-1}{b-a} \int_a^b \frac{p(s) \, ds}{(s-a)(b-s)} \right] \sqrt{b-a}.$$

By Green's formula (cf., for example, [16], [17]),

$$v(t) = \lambda \int_a^b G_0(t, \tau) q(\tau) \, d\tau \quad \text{for } a \leq t \leq b,$$

where  $G_0$  is Green's function of the problem

$$v'' = \frac{p(t)}{(t-a)^2(b-t)^2} v; \quad v(a+) = 0, \quad v(b-) = 0.$$

Applying Lemma 2.6, we can easily find

$$(3.14) \quad v(t) = \frac{\lambda \varrho_0^2}{b-a} \int_a^b (s-a)(b-s)q(s) ds \quad \text{for } a \leq t \leq b,$$

$$(3.15) \quad v(t) \geq \frac{\lambda \varrho_0^{-1}}{b-a} \left[ (b-t) \int_a^t (s-a)q(s) ds \right. \\ \left. + (t-a) \int_t^b (b-s)q(s) ds \right] \quad \text{for } a \leq t \leq b,$$

where  $\varrho_0 = \exp[(b-a)^{-1} \int_a^b [(s-a)(b-s)]^{-1} p(s) ds]$ .

By (1.15) and (3.13), from (3.14) and (3.15) we obtain

$$(t-a)(b-t) \leq v(t) \leq \frac{1}{\lambda} \quad \text{for } a \leq t \leq b.$$

Owing to this, (3.12) results in

$$v''(t) \geq \frac{p(t) - q(t)}{v(t)} \quad \text{for } a < t < b.$$

Hence  $v$  is a lower function of equation (1.14).

Let  $w$  be a solution of the problem

$$w'' = -\frac{q(t)}{w}; \quad w(a+) = 1 + \max\{v(t) : a \leq t \leq b\} = w(b-).$$

Then it is clear that  $w$  is an upper function of equation (1.14). Hence, by Theorem 1.1, problem (1.14), (1.2) has at least one solution.  $\square$

Theorem 1.2 is proved similarly to Theorem 1.1.

**P r o o f** of Corollary 1.4. Suppose  $\delta = \frac{1}{4}(b-a)^2$  and  $k = 1 + \frac{n}{\mu}$ , and choose  $\varepsilon > 0$  such that

$$\varepsilon \delta^k < 1, \quad \varepsilon \delta^{k-1} < \frac{1}{2k}, \quad \varepsilon^\mu (r + \delta^n) \delta^\mu < p_0.$$

Then it can be readily verified that  $\sigma_1(t) = \varepsilon[(t-a)(b-t)]^k$  for  $a \leq t \leq b$  is a lower function of equation (1.17), and  $\sigma_1(t) < 1$  for  $a \leq t \leq b$ .



Assume now that  $\sigma_2$  is a solution of the problem

$$u'' = -\frac{p(t)}{u^\mu}; \quad u(a+) = 1, \quad u(b-) = 1.$$

Evidently,  $\sigma_2(t) > 1$  for  $a \leq t \leq b$ .

Thus  $\sigma_1$  and  $\sigma_2$  are respectively lower and upper functions of equation (1.17) satisfying conditions (1.4). Hence by Theorem 1.2, problem (1.17), (1.2) is solvable. The uniqueness follows from the fact that the right-hand side of equation (1.17) is a nondecreasing function.  $\square$

### References

- [1] *J. A. Ackroyd*: On the laminar compressible boundary layer with stationary origin on a moving flat wall. *Proc. Cambridge Phil. Soc.* 63 (1967), 871–888.
- [2] *R. P. Agarwal and D. O'Regan*: Singular boundary value problems for superlinear second order ordinary and delay differential equations. *J. Differential Equations* 130 (1996), 333–355.
- [3] *J. E. Bouillet and S. M. Gomes*: An equation with singular nonlinearity related to diffusion problems in one dimension. *Quart. Appl. Math.* 42 (1985), 395–402.
- [4] *J. V. Baxley*: A singular nonlinear boundary value problem: membrane response of a spherical cap. *SIAM J. Appl. Math.* 48 (1988), 497–505.
- [5] *L. E. Bobisud, D. O'Regan and W. D. Royalty*: Solvability of some nonlinear boundary value problems. *Nonlinear Anal.* 12 (1988), 855–869.
- [6] *A. J. Callegary and M. B. Friedman*: An analytic solution of a nonlinear singular boundary value problem in the theory of viscous fluids. *J. Math. Anal. Appl.* 21 (1968), 510–529.
- [7] *A. J. Callegary and A. Nachman*: Some singular nonlinear differential equations arising in boundary layer theory. *J. Math. Anal. Appl.* 64 (1978), 96–105.
- [8] *A. J. Callegary and A. Nachman*: A nonlinear singular boundary value problem in the theory of pseudoplastic fluids. *SIAM J. Appl. Math.* 38 (1980), 275–281.
- [9] *D. R. Dunninger and J. C. Kurtz*: A priori bounds and existence of positive solutions for singular nonlinear boundary value problems. *SIAM J. Math. Anal.* 17 (1986), 595–609.
- [10] *J. A. Gatika, V. Oliker and P. Waltman*: Singular nonlinear boundary value problems for second order ordinary differential equations. *J. Differential Equations* 79 (1989), 62–78.
- [11] *Z. Guo*: Solvability of some singular nonlinear boundary value problems and existence of positive radial solutions of some nonlinear elliptic problems. *Nonlinear Anal.* 16 (1991), 781–790.
- [12] *J. Janus and A. Myjak*: A generalized Emden–Fowler equation with a negative exponent. *Nonlinear Anal.* 23 (1994), 953–970.
- [13] *P. Habets and F. Zanolin*: Upper and lower solutions for a generalized Emden–Fowler equation. *J. Math. Anal. Appl.* 181 (1994), 684–700.
- [14] *P. Habets and F. Zanolin*: Positive solutions for a class of singular boundary value problems. *Boll. Un. Mat. Ital. A* 9 (1995), 273–286.
- [15] *I. T. Kiguradze*: On some singular boundary value problems for nonlinear differential equations of the second order. *Differentsial'nye Uravneniya* 4 (1968), 1753–1773. (In Russian.)

- [16] *I. T. Kiguradze and A. G. Lomtatidze*: On certain boundary value problems for second order linear ordinary differential equations with singularities. *J. Math. Anal. Appl.* 101 (1984), 325–347.
- [17] *I. T. Kiguradze and B. L. Shekhter*: Singular boundary value problems for second order ordinary differential equations. In: *Curent Problems in Mathematics: Newest Results*, Vol. 3. Itogi Nauki i Tekhniki, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1987, pp. 3-103.
- [18] *Yu. A. Klovok and A. I. Lomakina*: One a boundary value problem with singularities on the ends of the segment. *Differ. Equations Latv. Mat. Ezhegodnik* 17 (1976), 179–186.
- [19] *A. Lomtatidze*: Positive solutions of boundary value problems for second order ordinary differential equations with singular points. *Differentsial'nye Uravneniya* 23 (1987), 1685–1692.
- [20] *A. Lomtatidze*: Existence of conjugate points for second order linear differential equations. *Georgian Math. J.* 2 (1995), 93–98.
- [21] *C. D. Luning and W. L. Perry*: Positive solutions of negative exponent generalized Emden–Fowler boundary value problems. *SIAM J. Math. Anal.* 12 (1981), 874–879.
- [22] *N. F. Morozov*: On analytic structure of a solution of the membrane equation. *Dokl. Akad. Nauk SSSR* 152 (1963), 78–80.
- [23] *N. F. Morozov and L. S. Srubshchik*: Application of Chaplygin’s method to investigation of the membrane equation. *Differentsial'nye Uravneniya* 2 (1966), 425–427. (In Russian.)
- [24] *L. S. Srubshchik and V. I. Yudovich*: Asymptotics of equation of large deflection of circular symmetrically loaded plate. *Sibirsk. Mat. Zh.* 4 (1963), 657–672. (In Russian.)
- [25] *S. Taliaferro*: A nonlinear singular boundary value problem. *Nonlinear Anal.* 3 (1979), 897–904.
- [26] *A. Tineo*: Existence theorems for a singular two-point Dirichlet problem. *Nonlinear Anal.* 19 (1992), 323–333.
- [27] *J. Wang*: Solvability of singular nonlinear two-point boundary value problems. *Nonlinear Anal.* 24 (1995), 555–561.

*Authors' addresses*: A. Lomtatidze, Mathematical Institute of the Academy of Sciences of the Czech Republic, Žižkova 22, 616 62 Brno, Czech Republic, and Department of Mathematical Analysis, Faculty of Natural Sciences, Masaryk University, Janáčkovo nám. 2a, 662 95 Brno, Czech Republic, e-mail: [bacho@math.muni.cz](mailto:bacho@math.muni.cz); P. Torres, Departamento de Matemática Aplicada, Universidad de Granada, 18071 Granada, Spain, e-mail: [ptorres@ugr.es](mailto:ptorres@ugr.es).