

Nikola Tuneski

On some simple sufficient conditions for univalence

Mathematica Bohemica, Vol. 126 (2001), No. 1, 229–236

Persistent URL: <http://dml.cz/dmlcz/133914>

Terms of use:

© Institute of Mathematics AS CR, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON SOME SIMPLE SUFFICIENT CONDITIONS FOR UNIVALENCE

NIKOLA TUNESKI, Skopje

(Received June 8, 1999)

Abstract. In this paper some simple conditions on $f'(z)$ and $f''(z)$ which lead to some subclasses of univalent functions will be considered.

Keywords: univalent, starlike, sharp result

MSC 2000: 30C45

1. INTRODUCTION AND PRELIMINARIES

Let A denote the class of analytic functions $f(z)$ in the unit disc $U = \{z: |z| < 1\}$ and normalized so that $f(0) = f'(0) - 1 = 0$.

A function $f(z) \in A$ is said to be *starlike of order* α , i.e., to belong to $S^*(\alpha)$, $0 \leq \alpha < 1$, if and only if

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > \alpha$$

for all $z \in U$. Then $S^* = S^*(0)$ is the class of *starlike functions* in the unit disc U . Further, $\tilde{S}^*(\alpha)$, $0 < \alpha \leq 1$, is the class of *strongly starlike functions of order* α defined by

$$\tilde{S}^*(\alpha) = \left\{ f(z) \in A: \left| \arg \frac{zf'(z)}{f(z)} \right| < \frac{\alpha\pi}{2}, z \in U \right\}.$$

Also $K(\alpha)$, $0 \leq \alpha < 1$, is the class of *convex functions of order* α which consists of functions $f(z) \in A$ such that

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \alpha$$

for all $z \in U$, and $K = K(0)$ is the class of *convex functions* on the unit disc U .

In addition to these classes we will deal also with the following ones:

$$R(\alpha) = \{f(z) \in A : \operatorname{Re}\{f'(z)\} > \alpha, z \in U\}, 0 \leq \alpha < 1;$$

$$R_\alpha = \left\{f(z) \in A : |\arg f'(z)| < \frac{\alpha\pi}{2}, z \in U\right\}, 0 < \alpha \leq 1.$$

All of the above mentioned classes are subclasses of univalent functions in U and moreover $K \subset S^*$ (see [1]). Further, S^* does not contain R_1 and R_1 does not contain S^* ([2]).

Let $f(z)$ and $g(z)$ be analytic in the unit disc U . Then we say that $f(z)$ is *subordinate* to $g(z)$, and we write $f(z) \prec g(z)$, if there exists a function $\omega(z)$ analytic in U such that $\omega(0) = 0$, $|\omega(z)| < 1$ and $f(z) = g(\omega(z))$ for all $z \in U$. If $g(z)$ is univalent in U , $f(0) = g(0)$ and $f(U) \subseteq g(U)$ then $f(z) \prec g(z)$.

The problem of finding $\lambda > 0$ such that the condition $|f''(z)| \leq \lambda$, $z \in U$, implies $f(z) \in S^*$ was first considered by Mocanu in his paper [3] for $\lambda = 2/3$. Later, Ponnusamy and Singh found a better constant $\lambda = 2/\sqrt{5}$, and recently Obradović in [4] closed this problem with the constant $\lambda = 1$ by proving that this result is sharp. In this paper, using similar techniques as Obradović did in [4] we will study λ such that the condition $|f''(z)| \leq \lambda$, $z \in U$, implies that $f(z)$ belongs to one of the classes defined above.

We will also generalize the result that Mocanu gave in [5]: $|f'(z) - 1| < 2/\sqrt{5}$, $z \in U$, implies $f(z) \in S^*$.

For all of this we will need the following two lemmas.

Lemma 1 ([6]). *Let $G(z)$ be convex and univalent in U , $G(0) = 1$. Let $F(z)$ be analytic in U , $F(0) = 1$ and let $F(z) \prec G(z)$ in U . Then for all $n \in \mathbb{N}_0$,*

$$(n+1)z^{-n-1} \int_0^z t^n F(t) dt \prec (n+1)z^{-n-1} \int_0^z t^n G(t) dt.$$

Lemma 2 ([7]). *Let $F(z)$ and $G(z)$ be analytic functions in the unit disc U and $F(0) = G(0)$. If $H(z) = zG'(z)$ is a starlike function in U and $zF'(z) \prec zG'(z)$ then*

$$F(z) \prec G(z) = G(0) + \int_0^z \frac{H(t)}{t} dt.$$

2. CONDITIONS ON $f''(z)$

Theorem 1. *If $f(z) \in A$ and $|f''(z)| \leq k$, $z \in U$, $0 < k \leq 1$, then*

$$(1) \quad \frac{zf'(z)}{f(z)} \prec 1 + \frac{k}{2-k}z$$

Proof. Noting that the condition of the theorem is equivalent to $zf''(z) \prec kz$, from lemma 1, choosing $F(z) = zf''(z) + 1$, $G(z) = kz + 1$ and $n = 0$, we get

$$f'(z) - \frac{f(z)}{z} \prec \frac{kz}{2},$$

which is equivalent to

$$(2) \quad z\left(\frac{f(z)}{z}\right)' \prec z\left(1 + \frac{kz}{2}\right)'$$

and to

$$(3) \quad \frac{f(z)}{z}\left(\frac{zf'(z)}{f(z)} - 1\right) \prec \frac{k}{2}z$$

for $z \in U$. Now, from (2) and lemma 2, taking $F(z) = f(z)/z$ and $G(z) = 1 + kz/2$ we obtain $f(z)/z \prec 1 + kz/2$, which implies $1 - k/2 < |f(z)/z| < 1 + k/2$, $z \in U$. From this relation and from (3) we can conclude that

$$\left(1 - \frac{k}{2}\right)\left|\frac{zf'(z)}{f(z)} - 1\right| < \left|\frac{f(z)}{z}\right|\left|\frac{zf'(z)}{f(z)} - 1\right| < \frac{k}{2}, \quad z \in U,$$

i.e.,

$$\left|\frac{zf'(z)}{f(z)} - 1\right| < \frac{k}{2-k},$$

$z \in U$, and (1) follows. □

Corollary 1. *If $f(z) \in A$ and $|f''(z)| \leq 2(1 - \alpha)/(2 - \alpha) = k$, $z \in U$, $0 \leq \alpha < 1$, then $f(z) \in S^*(\alpha)$. The result is sharp.*

Proof. It is obvious that the conditions of Theorem 1 are satisfied, and so from (1) we obtain that $\operatorname{Re}\{zf'(z)/f(z)\} > 1 - k/(2 - k) = \alpha$, $z \in U$, i.e., $f(z) \in S^*(\alpha)$. Further, the function $f(z) = z + (k + \varepsilon)z^2/2$, $0 < k \leq 1$, $0 < \varepsilon < 1$, proves that the result is sharp, i.e., that k defined in the corollary is the biggest for a given α because $|f''(z)| = k + \varepsilon > k$ and

$$\frac{zf'(z)}{f(z)} = \frac{2(1 + (k + \varepsilon)z)}{2 + (k + \varepsilon)z}$$

is smaller than α when z is real and close to -1 . Hence $f(z) \notin S^*(\alpha)$. □

Remark 1. For $\alpha = 0$ ($k = 1$) in Corollary 1 we get Theorem 1 from [4].

Corollary 1.1. *Let $f(z) \in A$. Then*

- (i) $|f''(z)| \leq 4/5$ implies $f(z) \in S^*(1/3)$;
- (ii) $|f''(z)| \leq 2/3$ implies $f(z) \in S^*(1/2)$; and
- (iii) $|f''(z)| \leq 1/2$ implies $f(z) \in S^*(2/3)$.

Corollary 2. *If $f(z) \in A$ and $|f''(z)| \leq 2 \sin(\alpha\pi/2)/(1 + \sin(\alpha\pi/2)) = k$, $z \in U$, $0 < \alpha \leq 1$, then $f(z) \in \tilde{S}^*(\alpha)$.*

Proof. Because the conditions from Theorem 1 are fulfilled, from the subordination (1) we get that $|\arg\{zf'(z)/f(z)\}| < \arcsin(k/(2-k)) = \alpha\pi/2$, $z \in U$, i.e., $f(z) \in \tilde{S}^*(\alpha)$. \square

Remark 2. The question about the sharpness of the result from Corollary 2 is open. It can be subject to further investigation if for given α , $0 < \alpha < 1$, $k = 2 \sin(\alpha\pi/2)/(1 + \sin(\alpha\pi/2))$ is the biggest number for which $|f''(z)| \leq k$, $z \in U$, implies $f(z) \in \tilde{S}^*(\alpha)$ (in [4] Obradović showed that for $\alpha = 1$, $k = 1$ is the biggest number with this property). The function $f(z) = z + (k + \varepsilon)z^2/2$, $0 < k < 1$, $\varepsilon > 0$, for which $|f''(z)| = k + \varepsilon > k$ cannot be used for proving sharpness because for each k , $0 < k < 1$, there exists an $\varepsilon > 0$ small enough such that $f(z) \in \tilde{S}^*(\alpha)$. This follows from the fact that for $z = re^{i\theta}$

$$\arg \frac{zf'(z)}{f(z)} = \arctan \frac{r(k + \varepsilon) \sin \theta}{2 + 3r(k + \varepsilon) \cos \theta + r^2(k + \varepsilon)^2}$$

and

$$\sup_{z \in U} \left| \arg \frac{zf'(z)}{f(z)} \right| = \arcsin \frac{k + \varepsilon}{2 - (k + \varepsilon)^2},$$

which is smaller than $\arcsin(k/(2-k)) = \alpha\pi/2$ for $\varepsilon > 0$ small enough.

Corollary 2.1. *Let $f(z) \in A$. Then*

- (i) $|f''(z)| \leq 2/3$ implies $f(z) \in \tilde{S}^*(1/3)$;
- (ii) $|f''(z)| \leq 2(\sqrt{2} - 1) = 0,8284\dots$ implies $f(z) \in \tilde{S}^*(1/2)$; and
- (iii) $|f''(z)| \leq 2(2\sqrt{3} - 3) = 0,9282\dots$ implies $f(z) \in \tilde{S}^*(2/3)$.

Using the next theorem we will obtain some results on the classes $K(\alpha)$, $R(\alpha)$ and R_α .

Theorem 2. *If $f(z) \in A$ and $|f''(z)| \leq k$, $z \in U$, $0 < k \leq 1$, then*

$$(4) \quad f'(z) \prec 1 + kz.$$

P r o o f. The condition $|f''(z)| \leq k, z \in U$, is equivalent to

$$(5) \quad z f''(z) \prec kz$$

$z \in U$, and again, using Lemma 2 for $F(z) = f'(z)$ and $G(z) = 1 + kz$, we get that the subordination (4) is true. \square

Corollary 3. *If $f(z) \in A$ and $|f''(z)| \leq (1 - \alpha)/(2 - \alpha) = k, z \in U, 0 \leq \alpha < 1$, then $f(z) \in K(\alpha)$. The result is sharp.*

P r o o f. Because the conditions from Theorem 2 are fulfilled we get that (4) and (5) are true, and from (5) with $p(z) = 1 + z f''(z)/f'(z)$ we conclude

$$(6) \quad (p(z) - 1)f'(z) \prec kz$$

for $z \in U$. Now, let us suppose that there exists $z_0 \in U$ such that $p(z_0) = \alpha + ix$. So from (4) and (6) it follows that

$$(7) \quad 1 - k < |f'(z_0)| < 1 + k$$

and

$$(8) \quad |(p(z_0) - 1)f'(z_0)| < k.$$

Further, using (7) we obtain

$$\begin{aligned} |(p(z_0) - 1)f'(z_0)|^2 &= |\alpha - 1 + ix|^2 |f'(z_0)|^2 \\ &> [(\alpha - 1)^2 + x^2](1 - k)^2 \\ &= (\alpha - 1)^2(1 - k)^2 + x^2(1 - k)^2 \\ &\geq (\alpha - 1)^2(1 - k)^2 = k^2 \end{aligned}$$

for $\alpha = (1 - 2k)/(1 - k)$ ($\Leftrightarrow k = (1 - \alpha)/(2 - \alpha)$), which contradicts to (8). Therefore we have proved that under the conditions of Corollary 3 $\operatorname{Re}\{1 + z f'(z)/f(z)\} > \alpha$ is true for any $z \in U$, i.e., $f(z) \in K(\alpha)$.

The proof that the result is sharp is again done by the function $f(z) = z + (k + \varepsilon)z^2/2$, $0 < k \leq 1/2$ and $\varepsilon > 0$, for which $|f''(z)| = k + \varepsilon > k$ and

$$\operatorname{Re}\left\{1 + \frac{z f''(z)}{f'(z)}\right\} = \frac{1 + 2z(k + \varepsilon)}{1 + z(k + \varepsilon)}$$

is smaller than α when z is real and close to -1 , i.e., $f(z) \notin K(\alpha)$. \square

Remark 3. For $\alpha = 0$, i.e., $k = 1/2$, Corollary 3 is equivalent to Theorem 3 from [4].

Corollary 3.1. *Let $f(z) \in A$. Then*

- (i) $|f''(z)| \leq 2/5$ implies $f(z) \in K(1/3)$;
- (ii) $|f''(z)| \leq 1/3$ implies $f(z) \in K(1/2)$; and
- (iii) $|f''(z)| \leq 1/4$ implies $f(z) \in K(2/3)$.

Corollary 4. *If $f(z) \in A$ and $|f''(z)| \leq 1 - \alpha = k$, $z \in U$, $0 \leq \alpha < 1$, then $f(z) \in R(\alpha)$. The result is sharp.*

Proof. Subordination (4) is true because the conditions from Theorem 2 are fulfilled and hence we conclude that $\operatorname{Re}\{f'(z)\} > 1 - k = \alpha$ for $z \in U$, $f(z) \in R(\alpha)$. Once again, using the function $f(z) = z + (k + \varepsilon)z^2/2$, $0 < k \leq 1$ and $\varepsilon > 0$, for which $|f''(z)| = k + \varepsilon > k$ and $f'(z) = 1 + (k + \varepsilon)z$ is smaller than α when z is real and close to -1 , we prove that the result of the corollary is sharp. \square

Corollary 4.1. *Let $f(z) \in A$. Then*

- (i) $|f''(z)| \leq 2/3$ implies $f(z) \in R(1/3)$;
- (ii) $|f''(z)| \leq 1/2$ implies $f(z) \in R(1/2)$;
- (iii) $|f''(z)| \leq 1/3$ implies $f(z) \in R(2/3)$.

Corollary 5. *If $f(z) \in A$ and $|f''(z)| \leq \sin(\alpha\pi/2) = k$, $z \in U$, $0 < \alpha \leq 1$, then $f(z) \in R_\alpha$. The result is sharp.*

Proof. From the subordination (4), which is true because the conditions of Theorem 2 are fulfilled, we obtain that $|\arg f'(z)| < \arcsin k = \alpha\pi/2$, $z \in U$, i.e., $f(z) \in R_\alpha$. And in this case the proof that the result is sharp is done by considering the function $f(z) = z + (k + \varepsilon)z^2/2$, $0 < k \leq 1$ and $\varepsilon > 0$, for which $|f''(z)| = k + \varepsilon > k$ and $\sup_{z \in U} |\arg f'(z)| = \arcsin(k + \varepsilon) > \arcsin k = \alpha\pi/2$ for $\varepsilon > 0$ small enough. \square

Corollary 5.1. *Let $f(z) \in A$. Then*

- (i) $|f''(z)| \leq 1/2$ implies $f(z) \in R_{1/3}$;
- (ii) $|f''(z)| \leq \sqrt{2}/2 = 0,7071\dots$ implies $f(z) \in R_{1/2}$; and
- (iii) $|f''(z)| \leq \sqrt{3}/2 = 0,8660\dots$ implies $f(z) \in R_{2/3}$.

3. CONDITION ON $f'(z)$

Theorem 3. *Let $f(z) \in A$. If $|f'(z) - 1| < \lambda$ for some $0 < \lambda \leq 1$ and for all $z \in U$, then $f(z) \in \tilde{S}^*(\alpha)$, where*

$$\alpha = \frac{2}{\pi} \arcsin \left(\lambda \sqrt{1 - \frac{\lambda^2}{4}} + \frac{\lambda}{2} \sqrt{1 - \lambda^2} \right),$$

and $|f(z)| < 1 + \lambda/2$ for $z \in U$.

P r o o f. From the condition $f'(z) \prec 1 + \lambda z$ it follows that

$$(9) \quad |\arg f'(z)| < \arcsin \lambda, \quad z \in U.$$

From the same condition, using lemma 1 for $F(z) = f'(z)$, $G(z) = 1 + \lambda z$ and $n = 0$ we get that

$$(10) \quad \frac{f(z)}{z} \prec 1 + \frac{\lambda}{2}z.$$

Consequently,

$$(11) \quad \left| \arg \frac{f(z)}{z} \right| < \arcsin \frac{\lambda}{2}$$

for $z \in U$. Now from (9) and (11) we can conclude that

$$\begin{aligned} \left| \arg \frac{zf'(z)}{f(z)} \right| &= \left| \arg \frac{z}{f(z)} + \arg f'(z) \right| \leq \left| \arg \frac{z}{f(z)} \right| + |\arg f'(z)| \\ &< \arcsin \frac{\lambda}{2} + \arcsin \lambda = \arcsin \left(\lambda \sqrt{1 - \frac{\lambda^2}{4}} + \frac{\lambda}{2} \sqrt{1 - \lambda^2} \right), \end{aligned}$$

i.e., $f(z) \in \tilde{S}^*(\alpha)$ for

$$(12) \quad \alpha = \frac{2}{\pi} \arcsin \left(\lambda \sqrt{1 - \frac{\lambda^2}{4}} + \frac{\lambda}{2} \sqrt{1 - \lambda^2} \right).$$

Further, from (10) it is easy to infer that for $z \in U$

$$|f(z)| < \left| \frac{f(z)}{z} \right| < 1 + \frac{\lambda}{2}.$$

□

We can rewrite Theorem 3 in the following way.

Theorem 3'. Let $f(z) \in A$, $0 < \alpha \leq 1$ and let

$$(13) \quad |f'(z) - 1| < 2a \sqrt{\frac{5 - 4\sqrt{1 - a^2}}{16a^2 + 9}} = \lambda,$$

where $a = \sin(\alpha\pi/2)$. Then $f(z) \in \tilde{S}^*(\alpha)$ and $|f(z)| < 1 + \lambda/2$ for $z \in U$.

Proof. If we put λ from (13) to the right side of (12) we obtain α . □

Corollary 6. Let $f(z) \in A$ and $|f'(z) - 1| < \lambda$. Then

- (i) if $\lambda = 2\sqrt{5}/5 = 0,8944\dots$, then $f(z) \in \tilde{S}^*(1) = S^*$ and $|f(z)| < 1 + \sqrt{5}/5 = 1,4472\dots$, for $z \in U$;
- (ii) if $\lambda = \sqrt{21}/7 = 0,6546\dots$, then $f(z) \in \tilde{S}^*(2/3)$ and $|f(z)| < 1 + \sqrt{21}/14 = 1,3273\dots$, for $z \in U$;
- (iii) if $\lambda = \sqrt{(10 - 4\sqrt{2})}/17 = 0,5054\dots$, then $f(z) \in \tilde{S}^*(1/2)$ and $|f(z)| < 1 + \lambda/2 = 1,2527\dots$, for $z \in U$;
- (iv) if $\lambda = \sqrt{(5 - 2\sqrt{3})}/13 = 0,3437\dots$, then $f(z) \in \tilde{S}^*(1/3)$ and $|f(z)| < 1 + \lambda/2 = 1,1718\dots$, for $z \in U$;

Remark 4. The result from Corollary 6 (i) is the same as the result from Theorem 2 from [5], but it is obtained by a different method.

- [1] Goodman, A. W.: Univalent Functions, Vol. I and II. Tampa, Florida, 1983.
- [2] Krzyz, J.: A counter example concerning univalent functions. Folia Societatis Scientiarum Lubliniensis Mat. Fiz. Chem. 2 (1962), 57–58.
- [3] Mocanu, P. T.: Two simple conditions for starlikeness. Mathematica (Cluj) 34 (1992), 175–181.
- [4] Obradović, M.: Simple sufficient conditions for starlikeness. Matemacki Vesnik. 49 (1997), 241–244.
- [5] Mocanu, P. T.: Some starlikeness conditions for analytic functions. Rev. Roumaine Math. Pures. Appl. 33 (1988), 117–124.
- [6] Halenbeck, D. J.: Ruscheweh, S.: Subordination by convex functions. Proc. Amer. Math. Soc. 52 (1975), 191–195.
- [7] Miller, S. S., Mocanu P. T.: Subordination-preserving integral operators. Trans. Amer. Math. Soc. 283 (1984), 605–615.

Author's address: Nikola Tuneski, Faculty of Mechanical Engineering Karpos II b.b., 91000 Skopje, R. Macedonia, e-mail: nikolat@reb.mf.ukim.edu.mk.