

Márcia Federson

Substitution formulas for the Kurzweil and Henstock vector integrals

*Mathematica Bohemica*, Vol. 127 (2002), No. 1, 15–26

Persistent URL: <http://dml.cz/dmlcz/133979>

## Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SUBSTITUTION FORMULAS FOR THE KURZWEIL AND  
HENSTOCK VECTOR INTEGRALS

M. FEDERSON, Brazil

(Received December 8, 1999)

*Abstract.* Results on integration by parts and integration by substitution for the variational integral of Henstock are well-known. When real-valued functions are considered, such results also hold for the Generalized Riemann Integral defined by Kurzweil since, in this case, the integrals of Kurzweil and Henstock coincide. However, in a Banach-space valued context, the Kurzweil integral properly contains that of Henstock. In the present paper, we consider abstract vector integrals of Kurzweil and prove Substitution Formulas by functional analytic methods. In general, Substitution Formulas need not hold for Kurzweil vector integrals even if they are defined.

*Keywords:* Kurzweil-Henstock integrals, integration by parts, integration by substitution

*MSC 2000:* 26A39

## 1. PRELIMINARIES

Throughout this paper,  $[a, b]$  is a compact interval of the real line  $\mathbb{R}$ ,  $X$  and  $Y$  are Banach spaces and  $L(X, Y)$  is the Banach space of all linear continuous functions from  $X$  to  $Y$ .

Let  $C([a, b], X)$  and  $G([a, b], X)$  be respectively the Banach spaces of continuous and of regulated functions from  $[a, b]$  to  $X$  endowed with the supremum norm which we denote by  $\|\cdot\|_\infty$ . We recall that a function  $f: [a, b] \rightarrow X$  is regulated if it has only discontinuities of the first kind, i.e. if the onesided limits  $\lim_{t \rightarrow c+} f(t)$  and  $\lim_{t \rightarrow c-} f(t)$  exist for every  $c \in [a, b]$ . Let  $C^\sigma([a, b], L(X, Y))$  be the set of all functions  $\alpha: [a, b] \rightarrow L(X, Y)$  which are weakly continuous (i.e. the function  $t \in [a, b] \rightarrow \alpha(t)x \in Y$  is continuous for every  $x \in X$ ) and let  $G^\sigma([a, b], L(X, Y))$  be the set of all weakly regulated functions  $\alpha: [a, b] \rightarrow L(X, Y)$  (i.e. the function  $t \in [a, b] \rightarrow \alpha(t)x \in Y$  is regulated for every  $x \in X$ ). See [4].

Let  $a = t_0 < t_1 < \dots < t_n = b$  be a division of  $[a, b]$ . In this case we write  $d = (t_i) \in D_{[a,b]}$  and  $|d| = n$ . Given  $d = (t_i) \in D_{[a,b]}$  and functions  $\alpha: [a, b] \rightarrow L(X, Y)$  and  $f: [a, b] \rightarrow X$ , we define

$$V_d(f) = \sum_i \|f(t_i) - f(t_{i-1})\|$$

and

$$SV_d(\alpha) = \sup \left\{ \left\| \sum_i [\alpha(t_i) - \alpha(t_{i-1})] y_i \right\|; y_i \in Y, \|y_i\| \leq 1 \right\}.$$

Then  $V(f) = \sup \{V_d(f); d \in D_{[a,b]}\}$  is the variation of  $f$  and  $SV(\alpha) = \sup\{SV_d(\alpha); d \in D_{[a,b]}\}$  is the semivariation of  $\alpha$ . If  $V(f) < \infty$ , then  $f$  is of bounded variation and we write  $f \in BV([a, b], X)$ . If  $SV(\alpha) < \infty$ , then  $\alpha$  is of bounded semivariation and we write  $\alpha \in SV([a, b], L(X, Y))$ . Clearly  $BV([a, b], L(X, Y)) \subset SV([a, b], L(X, Y))$ . Let  $X' = L(X, \mathbb{R})$ . Then  $SV([a, b], L(X, \mathbb{R})) = BV([a, b], X')$ . Moreover, for  $L(X) = L(X, X)$ ,  $SV([a, b], L(X)) = BV([a, b], L(X))$  if and only if  $X$  is of finite dimension.

For more information about these spaces, see [4].

## 2. BASIC DEFINITIONS AND PROPERTIES

When  $d = (t_i) \in D_{[a,b]}$  and  $\xi_i \in [t_{i-1}, t_i]$  for  $i = 1, 2, \dots, |d|$ , then  $d = (\xi_i, t_i)$  is called a tagged division of  $[a, b]$ . We denote by  $TD_{[a,b]}$  the set of all tagged divisions of  $[a, b]$ . A gauge of  $[a, b]$  is a function  $\delta: [a, b] \rightarrow ]0, \infty[$  and  $d = (\xi_i, t_i) \in TD_{[a,b]}$  is called  $\delta$ -fine if  $[t_{i-1}, t_i] \subset \{t \in [a, b]; |t - \xi_i| < \delta(\xi_i)\}$  for  $i = 1, 2, \dots, |d|$ .

In what follows we consider functions  $f: [a, b] \rightarrow X$  and  $\alpha: [a, b] \rightarrow L(X, Y)$ .

We say that  $f$  is Kurzweil  $\alpha$ -integrable (we write  $f \in K^\alpha([a, b], X)$ ) and that  $I \in Y$  is its integral (we write  $I = \int_{[a,b]}^K d\alpha(t) f(t)$ ) if given  $\varepsilon > 0$ , there is a gauge  $\delta$  of  $[a, b]$  such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a,b]}$ ,

$$\left\| \sum_{i=1}^{|d|} [\alpha(t_i) - \alpha(t_{i-1})] f(\xi_i) - \int_{[a,b]}^K d\alpha(t) f(t) \right\| < \varepsilon.$$

In particular, if we consider only constant gauges, then we obtain the Riemann-Stieltjes integral  $\int_{[a,b]} d\alpha(t) f(t)$ . We denote by  $R^\alpha([a, b], X)$  the set of all  $f: [a, b] \rightarrow X$  for which the Riemann-Stieltjes integral with respect to  $\alpha$  exists.

In an analogous way we say that  $\alpha$  is Kurzweil  $f$ -integrable (we write  $\alpha \in K_f([a, b], L(X, Y))$ ) and that  $I \in Y$  is its integral (we write  $I = \int_{[a,b]}^K \alpha(t) df(t)$ )

if for every  $\varepsilon > 0$ , there is a gauge  $\delta$  of  $[a, b]$  such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a,b]}$ ,

$$\left\| \sum_{i=1}^{|d|} \alpha(\xi_i) [f(t_i) - f(t_{i-1})] - \int_{[a,b]}^K \alpha(t) df(t) \right\| < \varepsilon.$$

Again, when constant gauges are considered, we obtain the space of Riemann-Stieltjes integrals  $\int_{[a,b]} \alpha(t) df(t)$  denoted by  $R_f([a, b], L(X, Y))$ .

Let  $f \in K^\alpha([a, b], X)$ . Its indefinite integral  $\tilde{f}^\alpha: [a, b] \rightarrow Y$  is given by  $\tilde{f}^\alpha(t) = \int_{[a,t]}^K \alpha(s) f(s)$  for every  $t \in [a, b]$ . Similarly, we denote the indefinite integral of  $\alpha \in K_f([a, b], L(X, Y))$  by  $\tilde{\alpha}_f: [a, b] \rightarrow Y$ , that is,  $\tilde{\alpha}_f(t) = \int_{[a,t]}^K \alpha(s) df(s)$  for every  $t \in [a, b]$ . When  $\alpha(t) = t$ , we write simply  $K([a, b], X)$ ,  $R([a, b], X)$  and  $\tilde{f}$  instead of  $K^\alpha([a, b], X)$ ,  $R^\alpha([a, b], X)$  and  $\tilde{f}^\alpha$ , respectively.

We say that  $f$  is Henstock  $\alpha$ -integrable (we write  $f \in H^\alpha([a, b], X)$ ) if there is a function  $F^\alpha: [a, b] \rightarrow Y$  such that for every  $\varepsilon > 0$ , there is a gauge  $\delta$  of  $[a, b]$  such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a,b]}$ ,

$$\sum_{i=1}^{|d|} \|\alpha(t_i) - \alpha(t_{i-1})\| f(\xi_i) - [F^\alpha(t_i) - F^\alpha(t_{i-1})]\| < \varepsilon.$$

We set  $\int_{[a,b]}^H \alpha(t) f(t) = F^\alpha(b) - F^\alpha(a)$  in this case. Analogously,  $\alpha$  is Henstock  $f$ -integrable (we write  $\alpha \in H_f([a, b], L(X, Y))$ ) if there is a function  $A_f: [a, b] \rightarrow Y$  such that for every  $\varepsilon > 0$ , there is a gauge  $\delta$  of  $[a, b]$  such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a,b]}$ ,

$$\sum_{i=1}^{|d|} \|\alpha(\xi_i) [f(t_i) - f(t_{i-1})] - [A_f(t_i) - A_f(t_{i-1})]\| < \varepsilon.$$

In this case we set  $\int_{[a,b]}^H \alpha(t) df(t) = A_f(b) - A_f(a)$ .

The inclusions  $H^\alpha([a, b], X) \subset K^\alpha([a, b], X)$  and  $H_f([a, b], L(X, Y)) \subset K_f([a, b], L(X, Y))$  hold. Moreover, if  $f \in H^\alpha([a, b], X)$ , then  $F^\alpha(t) - F^\alpha(a) = \int_{[a,t]}^K \alpha(t) f(t) = \tilde{f}^\alpha(t) - \tilde{f}^\alpha(a)$  for every  $t \in [a, b]$ . Analogously, given  $\alpha \in H_f([a, b], L(X, Y))$ , then  $A_f(t) - A_f(a) = \int_{[a,t]}^K \alpha(t) df(t) = \tilde{\alpha}_f(t) - \tilde{\alpha}_f(a)$  for every  $t \in [a, b]$ . Similarly as above if  $\alpha(t) = t$ , we write  $H([a, b], X)$  instead of  $H^\alpha([a, b], X)$ .

**Theorem 1** ([2, Theorem 1.2]). *If  $\alpha \in G^\sigma([a, b], L(X, Y))$  (or  $\alpha \in C^\sigma([a, b], L(X, Y))$ ) and  $f \in K^\alpha([a, b], X)$ , then  $\tilde{f}^\alpha \in G([a, b], X)$  (respectively  $\tilde{f}^\alpha \in C([a, b], X)$ ). Analogously, if  $f \in G([a, b], X)$  (or  $f \in C([a, b], X)$ ) and  $\alpha \in K_f([a, b], L(X, Y))$ , then  $\tilde{\alpha}_f \in G([a, b], Y)$  (respectively  $\tilde{\alpha}_f \in C([a, b], Y)$ ).*

**Theorem 2** ([2, Theorem 1.5]). *If either  $\alpha \in SV([a, b], L(X, Y))$  and  $f \in C([a, b], X)$ , or  $\alpha \in C([a, b], L(X, Y))$  and  $f \in BV([a, b], X)$ , then the Riemann-Stieltjes integrals  $\int_{[a, b]} \alpha(t) f(t)$  and  $\int_{[a, b]} \alpha(t) df(t)$  exist and the Integration by Parts Formula holds:*

$$\int_{[a, b]} \alpha(t) f(t) = \alpha(b) f(b) - \alpha(a) f(a) - \int_{[a, b]} \alpha(t) df(t).$$

**Theorem 3** ([9, Theorem 15]). *If  $f \in G([a, b], X)$ ,  $\alpha \in SV([a, b], L(X, Y))$  and  $\alpha \in G^\sigma([a, b], L(X, Y))$ , then  $f \in K^\alpha([a, b], X)$ .*

By a proof analogous to that of Schwabik for Theorem 3 one can show that

**Theorem 4.** *If  $f \in BV([a, b], X)$  and  $\alpha \in G([a, b], L(X, Y))$ , then  $\alpha \in K_f([a, b], L(X, Y))$ .*

**Lemma 1 (Saks-Henstock)** ([9, Lemma 16]).

(i) *Let  $\alpha: [a, b] \rightarrow L(X, Y)$  and  $f \in K^\alpha([a, b], X)$ . Given  $\varepsilon > 0$ , let the gauge  $\delta$  of  $[a, b]$  be such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a, b]}$ ,*

$$\left\| \sum_{i=1}^{|d|} [\alpha(t_i) - \alpha(t_{i-1})] f(\xi_i) - \int_{[a, b]}^K \alpha(t) f(t) \right\| < \varepsilon.$$

*Then for  $a \leq c_1 \leq \eta_1 \leq d_1 \leq c_2 \leq \eta_2 \leq d_2 \leq \dots \leq c_k \leq \eta_k \leq d_k \leq b$  with  $[c_j, d_j] \subset ]\eta_j - \delta(\eta_j), \eta_j + \delta(\eta_j)[$  for every  $j$ ,*

$$\left\| \sum_{j=1}^k \left\{ [\alpha(d_j) - \alpha(c_j)] f(\eta_j) - \int_{[c_j, d_j]}^K \alpha(t) f(t) \right\} \right\| \leq \varepsilon.$$

(ii) *Let  $f: [a, b] \rightarrow X$  and  $\alpha \in K_f([a, b], L(X, Y))$ . If for  $\varepsilon > 0$ , the gauge  $\delta$  of  $[a, b]$  is such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a, b]}$ ,*

$$\left\| \sum_{i=1}^{|d|} \alpha(\xi_i) [f(t_i) - f(t_{i-1})] - \int_{[a, b]}^K \alpha(t) df(t) \right\| < \varepsilon,$$

*then for  $a \leq c_1 \leq \eta_1 \leq d_1 \leq c_2 \leq \eta_2 \leq d_2 \leq \dots \leq c_k \leq \eta_k \leq d_k \leq b$  with  $[c_j, d_j] \subset ]\eta_j - \delta(\eta_j), \eta_j + \delta(\eta_j)[$  for every  $j$ ,*

$$\left\| \sum_{j=1}^k \left\{ \alpha(\eta_j) [f(d_j) - f(c_j)] - \int_{[c_j, d_j]}^K \alpha(t) df(t) \right\} \right\| \leq \varepsilon.$$

### 3. INTEGRATION BY PARTS FORMULAS

Using the definitions it is not difficult to show that the following holds.

**Theorem 5.** *If  $f \in H([a, b], X)$  and  $\alpha \in K_{\tilde{f}}([a, b], L(X, Y))$  (or  $\alpha \in H_{\tilde{f}}([a, b], L(X, Y))$ ) is bounded, then  $\alpha f \in K([a, b], Y)$  (respectively  $\alpha f \in H([a, b], Y)$ ) and*

$$\int_{[a,b]}^K \alpha(t) f(t) dt = \int_{[a,b]}^K \alpha(t) d\tilde{f}(t).$$

**Corollary 1.** *Let  $\alpha \in G([a, b], L(X, Y))$  and  $f \in H([a, b], X)$  be such that  $\tilde{f} \in BV([a, b], X)$ . Then  $\alpha f \in K([a, b], Y)$  and*

$$\int_{[a,b]}^K \alpha(t) f(t) dt = \int_{[a,b]}^K \alpha(t) d\tilde{f}(t).$$

*Proof.* By Theorem 4, the Kurzweil vector integral  $\int_{[a,b]}^K \alpha(t) d\tilde{f}(t)$  exists. □

In the next corollaries, we use the fact that the Riemann-Stieltjes integrals are special cases of the Kurzweil vector integrals.

**Corollary 2.** *Let  $\alpha \in C([a, b], L(X, Y))$  and  $f \in H([a, b], X)$  be such that  $\tilde{f} \in BV([a, b], X)$ . Then  $\alpha f \in K([a, b], Y)$  with*

$$(1) \quad \int_{[a,b]}^K \alpha(t) f(t) dt = \int_{[a,b]} \alpha(t) d\tilde{f}(t)$$

and

$$(2) \quad \int_{[a,b]} \alpha(t) d\tilde{f}(t) = \tilde{\alpha}(b) f(b) - \tilde{\alpha}(a) f(a) - \int_{[a,b]} d\alpha(t) \tilde{f}(t).$$

*Proof.* Follows by Theorem 2. □

**Corollary 3.** *If  $f \in H([a, b], X)$  and  $\alpha \in SV([a, b], L(X, Y))$ , then  $\alpha f \in K([a, b], Y)$  and equations (1) and (2) hold.*

*Proof.* Follows by Theorems 1 and 2. □

It is even true that

**Theorem 6** ([5]). *If  $f \in K([a, b], X)$  and  $\alpha \in SV([a, b], L(X, Y))$ , then  $\alpha f \in K([a, b], Y)$  and equations (1) and (2) hold.*

The proof of the above theorem is due to Hönig. It follows the steps of the proof of Theorem 11 below applying Theorems 1 and 2. When the functions are real-valued,  $SV([a, b], L(\mathbb{R})) = BV([a, b], L(\mathbb{R}))$  and Theorem 6 is also proved in [7, Theorem 12.1 and Corollary 12.2].

On the other hand,

**Theorem 7.** *If  $f \in H([a, b], X)$  and  $\alpha \in BV([a, b], L(X, Y))$ , then  $\alpha f \in H([a, b], Y)$  and equations (1) and (2) hold.*

*Proof.* By hypothesis,  $f \in H([a, b], X)$ . By Theorem 1,  $\tilde{f}$  is continuous. Then for every  $\varepsilon > 0$ , there exists  $\delta^* > 0$  such that for the oscillation  $\omega(\tilde{f}, [c, d])$  of  $\tilde{f}$  over the interval  $[c, d]$  we have  $\omega(\tilde{f}, [c, d]) < \varepsilon$  if  $0 < d - c < \delta^*$  and there is a gauge  $\delta$  of  $[a, b]$  for which  $\delta(t) < \delta^*/2$ ,  $t \in [a, b]$ , such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a, b]}$ ,

$$\sum_{i=1}^{|d|} \left\| f(\xi_i)(t_i - t_{i-1}) - \int_{[t_{i-1}, t_i]}^K f(t) dt \right\| < \varepsilon.$$

Thus,

$$\begin{aligned} & \sum_{i=1}^{|d|} \left\| \alpha(\xi_i) f(\xi_i)(t_i - t_{i-1}) - \int_{[t_{i-1}, t_i]}^K \alpha(t) f(t) dt \right\| \\ & \leq \sum_{i=1}^{|d|} \left\| \alpha(\xi_i) \left\{ f(\xi_i)(t_i - t_{i-1}) - \int_{[t_{i-1}, t_i]}^K f(t) dt \right\} \right\| \\ & \quad + \sum_{i=1}^{|d|} \left\| \int_{[t_{i-1}, t_i]}^K [\alpha(t) - \alpha(\xi_i)] f(t) dt \right\| \\ & < \|\alpha\|_\infty \varepsilon + \sum_{i=1}^{|d|} \left\| \int_{[t_{i-1}, t_i]}^K [\alpha(t) - \alpha(\xi_i)] f(t) dt \right\|. \end{aligned}$$

However, by Corollary 3 we have  $\alpha f \in K([a, b], Y)$  where

$$\int_{[a, b]}^K \alpha(t) f(t) dt = \int_{[a, b]} \alpha(t) d\tilde{f}(t) = \alpha(b)\tilde{f}(b) - \alpha(a)\tilde{f}(a) - \int_{[a, b]} d\alpha(t)\tilde{f}(t)$$

and a similar formula holds also for every subinterval contained in  $[a, b]$ . Hence, for  $\beta_{t_i} = [\alpha(t_i) - \alpha(\xi_i)] \tilde{f}(t_i)$  and  $\beta_{t_{i-1}} = [\alpha(t_{i-1}) - \alpha(\xi_i)] \tilde{f}(t_{i-1})$  we have

$$\begin{aligned} \sum_{i=1}^{|d|} \left\| \int_{[t_{i-1}, t_i]}^K [\alpha(t) - \alpha(\xi_i)] f(t) dt \right\| &= \sum_{i=1}^{|d|} \left\| \beta_{t_i} - \beta_{t_{i-1}} - \int_{[t_{i-1}, t_i]} d\alpha(t) \tilde{f}(t) \right\| \\ &= \sum_{i=1}^{|d|} \left\| \beta_{t_i} - \int_{[\xi_i, t_i]} d\alpha(t) \tilde{f}(t) - \beta_{t_{i-1}} - \int_{[t_{i-1}, \xi_i]} d\alpha(t) \tilde{f}(t) \right\| \\ &= \sum_{i=1}^{|d|} \left\| \int_{[\xi_i, t_i]} d\alpha(t) [\tilde{f}(t_i) - \tilde{f}(t)] + \int_{[t_{i-1}, \xi_i]} d\alpha(t) [\tilde{f}(t_{i-1}) - \tilde{f}(t)] \right\| \\ &\leq V(\alpha) \varepsilon, \end{aligned}$$

because  $\|\tilde{f}(t_i) - \tilde{f}(t)\| \leq \omega(\tilde{f}, [t_{i-1}, t_i])$  and  $\|\tilde{f}(t_{i-1}) - \tilde{f}(t)\| \leq \omega(\tilde{f}, [t_{i-1}, t_i])$  for every  $t \in [t_{i-1}, t_i]$ . The proof is complete.  $\square$

The next result can be proved by appealing to the definitions.

**Theorem 8.** *Let  $\alpha \in H([a, b], L(X, Y))$  and  $f \in K^{\tilde{\alpha}}([a, b], X)$  (or  $f \in H^{\tilde{\alpha}}([a, b], X)$ ). If  $f$  is bounded, then  $\alpha f \in K([a, b], Y)$  (respectively  $\alpha f \in H([a, b], Y)$ ) and*

$$\int_{[a, b]}^K \alpha(t) f(t) dt = \int_{[a, b]}^K d\tilde{\alpha}(t) f(t).$$

**Corollary 4.** *If  $\alpha \in H([a, b], L(X, Y))$  with  $\tilde{\alpha} \in SV([a, b], L(X, Y))$  and  $f \in G([a, b], X)$ , then  $\alpha f \in K([a, b], Y)$  and*

$$\int_{[a, b]}^K \alpha(t) f(t) dt = \int_{[a, b]}^K d\tilde{\alpha}(t) f(t).$$

**Proof.** By Theorem 1,  $\tilde{\alpha} \in C([a, b], L(X, Y))$ . Then the result comes by Theorem 3, since  $C([a, b], L(X, Y)) \subset G^\sigma([a, b], L(X, Y))$ .  $\square$

**Corollary 5.** *If  $\alpha \in H([a, b], L(X, Y))$  with  $\tilde{\alpha} \in SV([a, b], L(X, Y))$  and  $f \in C([a, b], X)$ , then  $\alpha f \in K([a, b], Y)$  and we have*

$$(3) \quad \int_{[a, b]}^K \alpha(t) f(t) dt = \int_{[a, b]} d\tilde{\alpha}(t) f(t)$$

and

$$(4) \quad \int_{[a, b]} d\tilde{\alpha}(t) f(t) = \tilde{\alpha}(b) f(b) - \tilde{\alpha}(a) f(a) - \int_{[a, b]} \tilde{\alpha}(t) df(t).$$



**Corollary 6.** *If  $\alpha \in H([a, b], L(X, Y))$  and  $f \in BV([a, b], X)$ , then  $\alpha f \in K([a, b], Y)$  and equations (3) and (4) hold.*

*Proof.* Follows by Theorems 1 and 2. □

More generally, we have

**Theorem 9.** *If  $\alpha \in K([a, b], L(X, Y))$  and  $f \in BV([a, b], X)$ , then  $\alpha f \in K([a, b], Y)$  and both (3) and (4) hold.*

**Theorem 10.** *If  $\alpha \in H([a, b], L(X, Y))$  and  $f \in BV([a, b], X)$ , then  $\alpha f \in H([a, b], Y)$  and equations (3) and (4) hold.*

The proof of Theorem 9 follows the steps of the proof of Theorem 12 below applying Theorems 1 and 2. The proof of Theorem 10 is analogous to that of Theorem 7.

#### 4. THE SUBSTITUTION FORMULAS

In this section, in addition to  $X$  and  $Y$ ,  $W$  also denotes a Banach space.

##### 4.1. Substitution formulas for the Kurzweil vector integrals.

**Theorem 11.** *Let  $f: [a, b] \rightarrow X$ ,  $\alpha \in K_f([a, b], L(X, W))$ ,  $g = \tilde{\alpha}_f: [a, b] \rightarrow W$  (i.e.,  $g(t) = {}^K\int_{[a,t]} \alpha(s) df(s)$  for every  $t \in [a, b]$ ) and  $\gamma \in SV([a, b], L(W, Y))$ . Then  $\gamma \in K_g([a, b], L(W, Y))$  if and only if  $\gamma\alpha \in K_f([a, b], L(X, Y))$ . In this case,*

$$(5) \quad \int_{[a,b]}^K \gamma(t) \alpha(t) df(t) = \int_{[a,b]}^K \gamma(t) dg(t).$$

*Proof.* Since  $\alpha \in K_f([a, b], L(X, W))$ , then given  $\varepsilon > 0$ , there is a gauge  $\delta$  of  $[a, b]$  such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a,b]}$ ,

$$\left\| \sum_{i=1}^{|d|} \left\{ \alpha(\xi_i) [f(t_i) - f(t_{i-1})] - \int_{[t_{i-1}, t_i]}^K \alpha(t) df(t) \right\} \right\| < \varepsilon.$$

Taking approximated sums for  ${}^K\int_{[a,b]} \gamma(t) \alpha(t) df(t)$  and  ${}^K\int_{[a,b]} \gamma(t) dg(t)$  we have

$$\begin{aligned} & \left\| \sum_{i=1}^{|d|} \gamma(\xi_i) \alpha(\xi_i) [f(t_i) - f(t_{i-1})] - \sum_{i=1}^{|d|} \gamma(\xi_i) [g(t_i) - g(t_{i-1})] \right\| \\ &= \left\| \sum_{i=1}^{|d|} \gamma(\xi_i) \left\{ \alpha(\xi_i) [f(t_i) - f(t_{i-1})] - \int_{[t_{i-1}, t_i]}^K \alpha(t) df(t) \right\} \right\| = I. \end{aligned}$$

However, if  $\gamma_i \in L(X, Y)$  and  $x_i \in X$ , then

$$\sum_{i=i}^n \gamma_i x_i = \sum_{j=1}^n (\gamma_j - \gamma_{j-1}) \left( \sum_{i=j}^n x_i \right) + \gamma_0 \left( \sum_{i=j}^n x_i \right), \quad n \in \mathbb{N}.$$

Thus, for  $x_i = \alpha(\xi_i) [f(t_i) - f(t_{i-1})] - K \int_{[t_{i-1}, t_i]} \alpha(t) \, df(t)$ ,  $\gamma_i = \gamma(\xi_i)$ ,  $\gamma_0 = \gamma(a)$  and  $n = |d|$  we have

$$I = \left\| \sum_{j=1}^{|d|} [\gamma(\xi_j) - \gamma(\xi_{j-1})] \left( \sum_{i=j}^{|d|} x_i \right) + \gamma_0 \left( \sum_{i=j}^{|d|} x_i \right) \right\| < SV(\gamma)\varepsilon + \|\gamma(a)\|\varepsilon$$

provided  $\left\| \sum_{i=j}^{|d|} x_i \right\| \leq \varepsilon$  for every  $j \in \{1, 2, \dots, |d|\}$ , by the Saks-Henstock Lemma (Lemma 1). □

**Corollary 7.** *Let  $f: [a, b] \rightarrow X$ ,  $\alpha \in K_f([a, b], L(X, W))$ ,  $g = \tilde{\alpha}_f \in BV([a, b], W)$  and  $\gamma \in G([a, b], L(W, Y))$ . Then  $\gamma \in K_g([a, b], L(W, Y))$ ,  $\gamma\alpha \in K_f([a, b], L(X, Y))$  and equation (5) holds.*

*Proof.* By Theorem 4,  $\gamma \in K_g([a, b], L(W, Y))$ . □

**Corollary 8.** *If  $f \in C([a, b], X)$ ,  $\alpha \in K_f([a, b], L(X, W))$ ,  $g = \tilde{\alpha}_f: [a, b] \rightarrow W$  and  $\gamma \in SV([a, b], L(W, Y))$ , then  $\gamma \in R_g([a, b], L(W, Y))$ ,  $\gamma\alpha \in K_f([a, b], L(X, Y))$  and equation (5) holds.*

*Proof.*  $\gamma \in R_g([a, b], L(W, Y)) \subset K_g([a, b], L(W, Y))$  by Theorems 1 and 2. □

**Theorem 12.** *Let  $\gamma: [a, b] \rightarrow L(W, Y)$ ,  $\alpha \in K^\gamma([a, b], L(X, W))$ ,  $f \in BV([a, b], X)$  and  $\beta = \tilde{\alpha}^\gamma: [a, b] \rightarrow L(X, Y)$  (i.e.,  $\beta(t) = K \int_{[a, t]} d\gamma(s) \alpha(s)$  for every  $t \in [a, b]$ ). Then  $f \in K^\beta([a, b], X)$  if and only if  $\alpha f \in K^\gamma([a, b], Y)$ . In this case,*

$$(6) \quad \int_{[a, b]}^K d\gamma(t) \alpha(t) f(t) = \int_{[a, b]}^K d\beta(t) f(t).$$

*Proof.* By hypothesis,  $\alpha \in K^\gamma([a, b], L(X, W))$ . Hence, given  $\varepsilon > 0$ , there exists a gauge  $\delta$  of  $[a, b]$  such that for every  $\delta$ -fine  $d = (\xi_i, t_i) \in TD_{[a, b]}$ ,

$$\left\| \sum_{i=1}^{|d|} \left\{ [\gamma(t_i) - \gamma(t_{i-1})] \alpha(\xi_i) - \int_{[t_{i-1}, t_i]}^K d\gamma(t) \alpha(t) \right\} \right\| < \varepsilon.$$

Taking approximated sums for  $K\int_{[a,b]} d\gamma(t) \alpha(t) f(t)$  and  $K\int_{[a,b]} d\beta(t) f(t)$  we have

$$\begin{aligned} & \left\| \sum_{i=1}^{|d|} [\gamma(t_i) - \gamma(t_{i-1})] \alpha(\xi_i) f(\xi_i) - \sum_{i=1}^{|d|} [\beta(t_i) - \beta(t_{i-1})] f(\xi_i) \right\| \\ &= \left\| \sum_{i=1}^{|d|} \left\{ [\gamma(t_i) - \gamma(t_{i-1})] \alpha(\xi_i) - \int_{[t_{i-1}, t_i]}^K d\gamma(t) \alpha(t) \right\} f(\xi_i) \right\|. \end{aligned}$$

However, if  $\gamma_i \in L(X, Y)$  and  $x_i \in X$ , then

$$\sum_{i=1}^n \gamma_i x_i = \left( \sum_{i=1}^n \gamma_i \right) x_0 + \sum_{j=1}^n \left( \sum_{i=j}^n \gamma_i \right) (x_j - x_{j-1}), \quad n \in \mathbb{N}.$$

Hence, taking  $\gamma_i = [\gamma(t_i) - \gamma(t_{i-1})] \alpha(\xi_i) - K\int_{[t_{i-1}, t_i]} d\gamma(t) \alpha(t)$ ,  $x_i = f(\xi_i)$ ,  $x_0 = f(a)$  and  $n = |d|$ , we obtain

$$\begin{aligned} I &\leq \left\| \sum_{i=1}^{|d|} \left\{ [\gamma(t_i) - \gamma(t_{i-1})] \alpha(\xi_i) - \int_{[t_{i-1}, t_i]}^K d\gamma(t) \alpha(t) \right\} \right\| \|f(a)\| \\ &\quad + \sum_{j=1}^{|d|} \left\| \sum_{i=j}^{|d|} \gamma_i \right\| \|f(t_i) - f(t_{i-1})\| < \varepsilon \|f(a)\| + \varepsilon V(f), \end{aligned}$$

since  $\left\| \sum_{i=j}^{|d|} \gamma_i \right\| = \left\| \sum_{i=j}^{|d|} \{ [\gamma(t_i) - \gamma(t_{i-1})] \alpha(\xi_i) - K\int_{[t_{i-1}, t_i]} d\gamma(t) \alpha(t) \} \right\| \leq \varepsilon$  for every  $j \in \{1, 2, \dots, |d|\}$ , by the Saks-Henstock Lemma (Lemma 1).  $\square$

**Corollary 9.** *If  $\gamma \in G^\sigma([a, b], L(W, Y))$ ,  $\alpha \in K^\gamma([a, b], L(X, W))$ ,  $\beta = \tilde{\alpha}^\gamma \in SV([a, b], L(X, Y))$  and  $f \in G([a, b], X)$ , then  $f \in K^\beta([a, b], X)$ ,  $\alpha f \in K^\gamma([a, b], Y)$  and equation (6) holds.*

*Proof.* By Theorem 1,  $\beta \in G([a, b], L(X, Y))$ . Then Theorem 3 implies that  $f \in K^\beta([a, b], X)$ .  $\square$

**Corollary 10.** *If  $\gamma \in C^\sigma([a, b], L(W, Y))$ ,  $\alpha \in K^\gamma([a, b], L(X, W))$ ,  $f \in BV([a, b], X)$  and  $\beta = \tilde{\alpha}^\gamma: [a, b] \rightarrow L(X, Y)$ , then  $f \in R^\beta([a, b], X)$ ,  $\alpha f \in K^\gamma([a, b], Y)$  and equation (6) holds.*

*Proof.* We have  $f \in R^\beta([a, b], X)$  by Theorems 1 and 2, and  $R^\beta([a, b], X) \subset K^\beta([a, b], X)$ .  $\square$

**Remark 1.** Similar results are valid for the Riemann-Stieltjes integrals, since the Saks-Henstock Lemma holds for  $R^\alpha([a, b], X)$  and  $R_f([a, b], L(X, Y))$  instead of  $K^\alpha([a, b], X)$  and  $K_f([a, b], L(X, Y))$ , respectively.

## 4.2. Substitution formulas for the Henstock vector integrals.

The results of this subsection are straightforward applications of the definitions.

**Theorem 13.** *Let  $f: [a, b] \rightarrow X$ ,  $\alpha \in H_f([a, b], L(X, W))$ ,  $g = \tilde{\alpha}f: [a, b] \rightarrow W$  (i.e.,  $g(t) = \int_{[a,t]}^K \alpha(s) df(s)$  for every  $t \in [a, b]$ ) and let  $\gamma: [a, b] \rightarrow L(W, Y)$  be bounded. Then  $\gamma \in H_g([a, b], L(W, Y))$  if and only if  $\gamma\alpha \in H_f([a, b], L(X, Y))$ . In this case, equation (5) holds.*

**Corollary 11.** *Let  $f \in H([a, b], X)$ . If  $\alpha: [a, b] \rightarrow L(X, Y)$  is bounded, then  $\alpha \in H_{\tilde{f}}([a, b], L(X, Y))$  if and only if  $\alpha f \in H([a, b], Y)$ . In this case,*

$$\int_{[a,b]}^K \alpha(t) f(t) dt = \int_{[a,b]}^K \alpha(t) d\tilde{f}(t).$$

**Theorem 14.** *Let  $\gamma: [a, b] \rightarrow L(W, Y)$ ,  $\alpha \in H^\gamma([a, b], L(X, W))$ ,  $\beta = \tilde{\alpha}\gamma: [a, b] \rightarrow L(X, Y)$  (i.e.,  $\beta(t) = \int_{[a,t]}^K d\gamma(s) \alpha(s)$  for every  $t \in [a, b]$ ) and let  $f: [a, b] \rightarrow X$  be bounded. Then  $f \in H^\beta([a, b], X)$  if and only if  $\alpha f \in H^\gamma([a, b], Y)$ . In this case, equation (6) holds.*

**Corollary 12.** *Let  $\alpha \in H([a, b], L(X, Y))$  and  $f: [a, b] \rightarrow X$  be bounded. Then  $f \in H^{\tilde{\alpha}}([a, b], X)$  if and only if  $\alpha f \in H([a, b], Y)$ . In this case,*

$$\int_{[a,b]}^K \alpha(t) f(t) dt = \int_{[a,b]}^K d\tilde{\alpha}(t) f(t).$$

**Remark 2.** Similar results hold for the vector integrals of McShane, which gives a Riemannian definition of the Bochner-Lebesgue-Stieltjes integrals when we introduce McShane's modification in the Henstock vector integrals ([8], [6]).

**Acknowledgement.** The author thanks Prof. Dr. Chaim Samuel Hönig for the suggestion of the subject.

## References

- [1] *Federson, M.*: The Fundamental Theorem of Calculus for multidimensional Banach space-valued Henstock vector integrals. *Real Anal. Exchange* 25 (2000), 469–480.
- [2] *Federson, M., Bianconi, R.*: Linear integral equations of Volterra concerning the integral of Henstock. *Real Anal. Exchange* 25 (2000), 389–417.
- [3] *Henstock, R.*: The General Theory of Integration. Oxford Math. Monog., Clarendon Press, Oxford, 1991.
- [4] *Hönig, C. S.*: Volterra-Stieltjes Integral Equations. *Math. Studies*, 16, North-Holland Publ. Comp., Amsterdam, 1975.
- [5] *Hönig, C. S.*: There is no natural Banach space norm on the space of Kurzweil-Henstock-Denjoy-Perron integrable functions. *Seminário Brasileiro de Análise* 30 (1989), 387–397.
- [6] *Hönig, C. S.*: A Riemanniann characterization of the Bochner-Lebesgue integral. *Seminário Brasileiro de Análise* 35 (1992), 351–358.
- [7] *Lee, P. Y.*: Lanzhou Lectures on Henstock Integration. World Sci., Singapore, 1989.
- [8] *McShane, E. J.*: A unified theory of integration. *Amer. Math. Monthly* 80 (1973), 349–359.
- [9] *Schwabik, Š.*: Abstract Perron-Stieltjes integral. *Math. Bohem.* 121 (1996), 425–447.

*Author's address:* *M. Federson*, Universidade de São Paulo, Departamento de Matemática, São Carlos, CP 688, CEP 13560-970, São Carlos-SP, Brazil, e-mail: `federson@icmc.sc.usp.br`.