

Zafer Ercan

A remark on supra-additive and supra-multiplicative operators on $C(X)$

Mathematica Bohemica, Vol. 132 (2007), No. 1, 55–58

Persistent URL: <http://dml.cz/dmlcz/133995>

Terms of use:

© Institute of Mathematics AS CR, 2007

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A REMARK ON SUPRA-ADDITIVE AND SUPRA-MULTIPLICATIVE
OPERATORS ON $C(X)$

Z. ERCAN, Ankara

(Received September 9, 2005)

Abstract. M. Radulescu proved the following result: Let X be a compact Hausdorff topological space and $\pi: C(X) \rightarrow C(X)$ a supra-additive and supra-multiplicative operator. Then π is linear and multiplicative. We generalize this result to arbitrary topological spaces.

Keywords: $C(X)$ -space, supra-additive, supra-multiplicative operator, realcompact

MSC 2000: 46J10, 46E25

1. THE RESULT

We follow the terminology of [1]. As usual for a topological space X , the space of real valued continuous (bounded) functions on X is denoted by $C(X)$ ($C_b(X)$). For each $x \in X$, $\delta_x: C(X) \rightarrow \mathbb{R}$ is defined by $\delta_x(f) = f(x)$. For $B \subset X$, χ_B denotes the characteristic function of B . For each $n \in \mathbb{R}$, \mathbf{n} denotes the constant function with value \mathbf{n} . A map $\pi: C(X) \rightarrow C(X)$ is called

- (i) *supra-additive* if $\pi(f + g) \geq \pi(f) + \pi(g)$ for each $f, g \in C(X)$,
- (ii) *supra-multiplicative* if $\pi(fg) \geq \pi(f)\pi(g)$ for each $f, g \in C(X)$.

The following theorem is the main result of [4].

Theorem 1. *Let X be a compact Hausdorff space and $\pi: C(X) \rightarrow C(X)$ a supra-additive and supra-multiplicative map. Then π is multiplicative and linear.*

The main result of this note is to generalize the above theorem as follows.

Theorem 2. Let X and Y be topological spaces and $\pi: C(X) \rightarrow C(Y)$ a supra-additive and supra-multiplicative map. Then the following statements are equivalent.

- (i) $\pi(f^+ \wedge \mathbf{n} - f^- \wedge \mathbf{n})(y) \rightarrow \pi(f)(y)$ for each $f \in C(X)$ and $y \in Y$.
- (ii) π is linear and multiplicative.

Proof. (ii) \implies (i): For each $y \in T$, $\delta_y \circ \pi$ is a Riesz homomorphism, so

$$\pi(f \wedge \mathbf{n})(y) = \delta_y \circ \pi(f \wedge \mathbf{n}) = \delta_y \circ \pi(f) \wedge n \rightarrow \delta_y \circ \pi(f) = \pi(f)(y)$$

(i) \implies (ii):

Claim 1. Let K be a compact Hausdorff space and let $T: C(K) \rightarrow \mathbb{R}$ be supra-additive and supra-multiplicative. Then T is linear and multiplicative.

Indeed, let $T^\sim: C(K) \rightarrow C(K)$ be defined by $T^\sim(f) = T(f)\mathbf{1}$. Then T^\sim is supra-additive and supra-multiplicative, so by Theorem 1, T^\sim is linear and multiplicative, so T is linear and multiplicative.

Claim 2. For each topological space M there exists a compact Hausdorff space K_M such that $C(K_M)$ and $C_b(M)$ are Riesz and algebraic isomorphic spaces.

As $C_b(M)$ is an AM-space with order unit $\mathbf{1}$, this follows from the Kakutani-Krein Representation Theorem (see [1]).

Claim 3. Let $\pi^\sim = \pi|_{C_b(X)}$. Then for each $y \in Y$, $\delta_y \circ \pi^\sim: C_b(X) \rightarrow \mathbb{R}$ is linear and multiplicative.

This follows from Theorem 1 and from the above claims.

Claim 4. π is linear.

To see this we use the linearity of $\delta_y \circ \pi^\sim$ as follows. Let $f, g \geq 0$ be given. Then

$$\pi(f + g)(y) = \lim \delta_y \circ \pi^\sim((f + g) \wedge \mathbf{n}) \leq \lim \delta_y \circ \pi^\sim(f \wedge \mathbf{n} + g \wedge \mathbf{n}).$$

Since $\delta_y \circ \pi^\sim$ is linear and π is supra-additive we have

$$\pi(f + g) \leq \pi(f) + \pi(g) \leq \pi(f + g),$$

so π is additive on $C(X)^+$. Now by the Kantorovic Theorem (see Theorem 1.7. [1]), $\varphi: C(X) \rightarrow C(Y)$ defined by $\varphi(f) = \pi(f^+) - \pi(f^-)$ is linear and from the second assumption it is clear that $\varphi = \pi$, so π is linear.

Claim 5. π is multiplicative.

Indeed, let $0 \leq f \in C(X)$ be given. As for each $y \in Y$, $\delta_y \circ \pi^\sim$ is multiplicative, we have

$$\begin{aligned} \pi(f^2)(y) &= \delta_y \circ \pi(f^2) = \lim \delta_y \circ \pi^\sim(f^2 \wedge \mathbf{n}) = \lim \delta_y \circ \pi^\sim((f \wedge \mathbf{n}^{\frac{1}{2}})^2) \\ &= (\lim \delta_y \circ \pi^\sim(f \wedge \mathbf{n}^{\frac{1}{2}}))^2 = \pi(f)^2(y), \end{aligned}$$

so $\pi(f^2) = \pi(f)^2$. Let $f \in C(X)$ be given. As $\pi(f^+)\pi(f^-) = 0$, due to the linearity of π we have $\pi(f^2) = \pi(f)^2$. Now the multiplicativity follows from the equality

$$fg = \frac{1}{4}((f+g)^2 - (f-g)^2).$$

□

Recall that a topological space X is called pseudocompact if $C(X) = C_b(X)$ ([3]). It is clear that any countable compact space is pseudocompact. Now the following corollary immediately follows from the above theorem.

Corollary 3. *Let X be a pseudocompact space and Y a topological space. A map $\pi: C(X) \rightarrow C(Y)$ is supra-additive and supra-multiplicative if and only if it is linear and multiplicative.*

Recall that a topological space is called *realcompact* if it is homeomorphic to a closed subspace of the product space of \mathbb{R} . It is well known that a Hausdorff space is compact if and only if it is realcompact and pseudocompact (see [3]). If K is a realcompact space and $T: C(K) \rightarrow \mathbb{R}$ is nonzero linear and multiplicative then there exists $k \in K$ such that $T(f) = f(k)$ for each $f \in C(K)$ (see [2] for a simple proof). By using this fact we have the following theorem.

Theorem 4. *Let X be a realcompact space and let Y be an arbitrary topological space. Let $\pi: C(X) \rightarrow C(Y)$ be a supra-additive and supra-multiplicative map. Then the following assertions are equivalent.*

- (i) $\pi(f^+ \wedge \mathbf{n} - f^- \wedge \mathbf{n})(y) \rightarrow \pi(f)(y)$ for each $f \in C(X)$ and $y \in Y$
- (ii) *There exists a clopen subset $B \subset Y$ and a continuous function $\sigma: Y \rightarrow X$ such that*

$$\pi(f)(y) = \chi_B(y)f(\sigma(y))$$

for each $y \in Y, f \in C(X)$.

Proof. It is clear that (ii) \implies (i). Suppose that (i) holds. Then from Theorem 2, π is linear and multiplicative. The fact that $\pi(\mathbf{1})^2 = \pi(\mathbf{1})$ for each $y \in Y$ implies that either $\pi(\mathbf{1})(y) = 0$ or $\pi(\mathbf{1})(y) = 1$, so $B = \{y \in Y: \pi(\mathbf{1})(y) = 1\}$ is clopen in Y . Let $y \in Y$ be given. As X is realcompact and $\delta_y \circ \pi: C(X) \rightarrow \mathbb{R}$ is linear and multiplicative there exists $\alpha(y)$ such that

$$\pi(f)(y) = \pi(\mathbf{1})(y)f(\alpha(y)) = \chi_B(y)f(\alpha(y)).$$

Since X is completely regular Hausdorff space, $\alpha(y)$ must be unique for each $y \in B$. Let $x_0 \in Y$ be fixed and let $\sigma: Y \rightarrow X$ be defined by $\sigma(y) = \alpha(y)$ when $y \in B$ and $\sigma(y) = x_0$ otherwise. It is clear that $\sigma|_B: B \rightarrow X$ is continuous. Since B is clopen, actually σ itself is continuous. This completes the proof. □

References

- [1] *C. D. Aliprantis, O. Burkinshaw*: Positive Operators. Academic Press, New York, 1985. [Zbl 0608.47039](#)
- [2] *Z. Ercan, S. Önal*: A remark on the homomorphism on $C(X)$. Proc. Amer. Math. Soc. *133* (2005), 3609–3611. [Zbl 1087.46038](#)
- [3] *K. P. Hart, J. Nagata, J. E. Vaughan*: Encyclopedia of General Topology. Elsevier, Amsterdam, 2004. [Zbl 1059.54001](#)
- [4] *M. Radulescu*: On a supra-additive and supra-multiplicative operator of $C(X)$. Bull. Math. Soc. Sci. Math. Répub. Soc. Roum., Nouv. Sér. *24* (1980), 303–305. [Zbl 0463.47034](#)

Author's address: Z. Ercan, Middle East Technical University, Department of Mathematics, 065 31 Ankara, Turkey, e-mail: zercan@metu.edu.tr.