

A. Ebadian; A. A. Shokri

On the Lipschitz operator algebras

Archivum Mathematicum, Vol. 45 (2009), No. 3, 213--222

Persistent URL: <http://dml.cz/dmlcz/134233>

Terms of use:

© Masaryk University, 2009

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON THE LIPSCHITZ OPERATOR ALGEBRAS

A. EBADIAN AND A. A. SHOKRI

ABSTRACT. In a recent paper by H. X. Cao, J. H. Zhang and Z. B. Xu an α -Lipschitz operator from a compact metric space into a Banach space A is defined and characterized in a natural way in the sense that $F : K \rightarrow A$ is a α -Lipschitz operator if and only if for each $\sigma \in X^*$ the mapping $\sigma \circ F$ is a α -Lipschitz function. The Lipschitz operators algebras $L^\alpha(K, A)$ and $l^\alpha(K, A)$ are developed here further, and we study their amenability and weak amenability of these algebras. Moreover, we prove an interesting result that $L^\alpha(K, A)$ and $l^\alpha(K, A)$ are isometrically isomorphic to $L^\alpha(K) \otimes A$ and $l^\alpha(K) \otimes A$ respectively. Also we study homomorphisms on the $L_A^\alpha(X, B)$.

1. INTRODUCTION

Let (K, d) be compact metric space with at least two elements and $(X, \|\cdot\|)$ be a Banach space over the scalar field \mathbf{F} ($= \mathbf{R}$ or \mathbf{C}). For a constant $\alpha > 0$ and an operator $T : K \rightarrow X$, set

$$(1) \quad L_\alpha(T) := \sup_{s \neq t} \frac{\|T(t) - T(s)\|}{d(s, t)^\alpha},$$

which is called the Lipschitz constant of T . Define

$$T_\alpha(x, y) = \frac{T(x) - T(y)}{d(x, y)^\alpha}, \quad x \neq y$$
$$L^\alpha(K, X) = \{T : K \rightarrow X : L_\alpha(T) < \infty\}$$

and

$$l^\alpha(K, X) = \{T : K \rightarrow X : \|T_\alpha(x, y)\| \rightarrow 0 \text{ as } d(x, y) \rightarrow 0\}.$$

The elements of $L^\alpha(K, X)$ and $l^\alpha(K, X)$ are called big and little Lipschitz operators, respectively [1].

Let $C(K, X)$ be the set of all continuous operators from K into X and for each $T \in C(K, X)$, define

$$\|T\|_\infty = \sup_{x \in K} \|T(x)\|.$$

For S, T in $C(K, X)$ and λ in \mathbf{F} , define

$$(S + T)(x) = S(x) + T(x), \quad (\lambda T)(x) = \lambda T(x), \quad (x \in X).$$

2000 *Mathematics Subject Classification*: primary 47B48; secondary 46J10.

Key words and phrases: Lipschitz algebras, amenability, homomorphism.

Received October 7, 2008. Editor A. Pultr.

It is easy to see that $(C(K, X), \|\cdot\|_\infty)$ becomes a Banach space over F and $L^\alpha(K, X)$ is a linear subspace of $C(K, X)$. For each element T of $L^\alpha(K, X)$, define $\|T\|_\alpha = L_\alpha(T) + \|T\|_\infty$.

In their papers [3, 4], Cao, Zhang and Xu proved that $(L^\alpha(K, X), \|\cdot\|_\alpha)$ is a Banach space over F and $l^\alpha(K, X)$ is a closed linear subspace of $(L^\alpha(K, X), \|\cdot\|_\alpha)$. Now, let $(A, \|\cdot\|)$ be a unital Banach algebra with unit e . In this paper, we show that $(L^\alpha(K, A), \|\cdot\|_\alpha)$ is a Banach algebra under pointwise and scalar multiplication and $l^\alpha(K, A)$ is a closed linear subalgebra of $(L^\alpha(K, A), \|\cdot\|_\alpha)$ and study many aspects of these algebras. The spaces $L^\alpha(K, A)$ and $l^\alpha(K, A)$ are called big and little Lipschitz operators algebras. Note that Lipschitz operators algebras are, in fact, extensions of Lipschitz algebras. Sherbert [11, 12], Weaver [13, 14], Honary and Mahyar [7], Johnson [8, 9], Alimohammadi and Ebadian [1], Ebadian [6], Bade, Curtis and Dales [2], studied some properties of Lipschitz algebras. We will study (weak) amenability of Lipschitz operators algebras. Also we study homomorphisms on the $L_A^\alpha(X, B)$.

2. CHARACTERIZATIONS OF LIPSCHITZ OPERATORS ALGEBRAS

In this section, let (K, d) be a compact metric space which has at least two elements and $(A, \|\cdot\|)$ to denote a unital Banach algebra over the scalar field F ($= \mathbb{R}$ or \mathbb{C}).

Theorem 2.1. *$(L^\alpha(K, A), \|\cdot\|_\alpha)$ is a Banach algebra over F and $l^\alpha(K, A)$ is a closed linear subspace of $(L^\alpha(K, A), \|\cdot\|_\alpha)$.*

Proof. As we have already $L^\alpha(K, A)$ is a Banach space and $l^\alpha(K, A)$ is a closed linear subspace if it. Now let $T, S \in L^\alpha(K, A)$, and define

$$(TS)(t) = T(t)S(t) \quad (t \in K).$$

Then

$$\begin{aligned} \|TS\|_\alpha &= \|TS\|_\infty + L_\alpha(TS) \\ &\leq \|T\|_\infty \|S\|_\infty + \sup_{t \neq s} \frac{\|T(t)S(t) - T(s)S(s)\|}{d(t, s)^\alpha} \\ &\leq \|T\|_\infty \|S\|_\infty + \|T\|_\infty L_\alpha(S) + \|S\|_\infty L_\alpha(T) \\ &\leq (\|T\|_\infty + L_\alpha(T)) (\|S\|_\infty + L_\alpha(S)) \\ &= \|T\|_\alpha \|S\|_\alpha. \end{aligned}$$

So that we see that $(L^\alpha(K, A), \|\cdot\|_\alpha)$ is a Banach algebra and $l^\alpha(K, A)$ is a closed linear subspace of $(L^\alpha(K, A), \|\cdot\|_\alpha)$. □

Theorem 2.2. *Let (K, d) be a compact metric space. Then $L^\alpha(K, A)$ is uniformly dense in $C(K, A)$.*

Proof. Let $f \in C(K, A)$. Then for every $\sigma \in A^*$ we have $\sigma \circ f \in C(K)$, so that there is $g \in L^\alpha(K)$ such that $\|g - \sigma \circ f\|_\infty < \varepsilon$. We define, the map $\eta: C \rightarrow A$ by

$\eta(\lambda) = \lambda \cdot e$. It is easy to see that $\eta \circ g \in L^\alpha(K, A)$, and for every $\sigma \in A^*$, we have

$$|\sigma(g(x) \cdot e - f(x))| = |g(x) - (\sigma \circ f)(x)| < \varepsilon, \quad (x \in K).$$

Therefore $|\sigma(\eta \circ g - f)(x)| < \varepsilon$ for every $\sigma \in A^*$ and $x \in K$. This implies that $\|(\eta \circ g - f)(x)\| < \varepsilon$ for every $x \in K$. Therefore, $\|\eta \circ g - f\|_\infty < \varepsilon$ and the proof is complete. \square

Remark 2.3. Let A, B be unital Banach algebras over \mathbb{F} . Then the injective tensor $A \check{\otimes} B$ is a unital Banach algebra under norm $\|\cdot\|_\varepsilon$, [10].

Theorem 2.4. $L^\alpha(K, A) = \{F : K \rightarrow A \mid \sigma \circ F \in L^\alpha(K, \mathbb{C}), (\forall \sigma \in A^*)\}$

Proof. Use the principle of Uniform Boundedness. \square

Lemma 2.5. Let $(E_1, \|\cdot\|_1), (E_2, \|\cdot\|_2)$ be Banach spaces. Then for $G \in E_1 \check{\otimes} E_2$

$$\|G\|_\varepsilon = \sup \{ \|(\text{id} \otimes \phi)(G)\|_1 : \phi \in E_2^*, \|\phi\| \leq 1 \}.$$

Proof. See [10]. \square

Theorem 2.6. Let (K, d) be a compact metric space and A be a unital commutative Banach algebra. Then $L^\alpha(K, A)$ is isometrically isomorphic to $L^\alpha(K) \check{\otimes} A$.

Proof. It is straightforward to prove that the mapping $V : L^\alpha(K) \times A \rightarrow L^\alpha(K, A)$ defined by

$$\begin{aligned} V(f, a) &= fa \quad (f \in L^\alpha(K), a \in A), \\ (fa)(x) &:= f(x)a \quad (x \in K), \end{aligned}$$

is bilinear. Therefore there exists a unique linear map $T : L^\alpha(K) \check{\otimes} A \rightarrow L^\alpha(K, A)$ such that $T(f \otimes a) = V(f, a) = fa$, [10]. For every $G \in L^\alpha(K) \check{\otimes} A$, there is $m \in \mathbb{N}$, $f_j \in L^\alpha(K)$ and $a_j \in A$ ($1 \leq j \leq m$) such that $G = \sum_{j=1}^m f_j \otimes a_j$, so we have

$$\begin{aligned} \|G\|_\varepsilon &= \sup_{\phi \in A^*, \|\phi\| \leq 1} \|(\text{id} \otimes \phi)(G)\|_\alpha = \sup_{\phi \in A^*, \|\phi\| \leq 1} \left\| (\text{id} \otimes \phi) \left(\sum_{j=1}^m f_j \otimes a_j \right) \right\| \\ &= \sup_{\phi \in A^*, \|\phi\| \leq 1} \left\| \sum_{j=1}^m f_j \phi(a_j) \right\|_\alpha = \sup_{\phi \in A^*, \|\phi\| \leq 1} \left[\sup_{x \in K} \left| \sum_{j=1}^m f_j(x) \phi(a_j) \right| \right. \\ &\quad \left. + \sup_{x \neq y} \frac{|\sum_{j=1}^m f_j(x) \phi(a_j) - \sum_{j=1}^m f_j(y) \phi(a_j)|}{d^\alpha(x, y)} \right] \\ &= \sup_{\phi \in A^*, \|\phi\| \leq 1} \left[\sup_{x \in K} \left| \phi \left(\sum_{j=1}^m f_j(x) a_j \right) \right| \right. \\ &\quad \left. + \sup_{x \neq y} \frac{|\phi(\sum_{j=1}^m (f_j(x) a_j - f_j(y) a_j))|}{d^\alpha(x, y)} \right] \end{aligned}$$

$$\begin{aligned}
&\leq \sup_{\phi \in A^*, \|\phi\| \leq 1} \left[\sup_{x \in K} \|\phi\| \left\| \sum_{j=1}^m f_j(x) a_j \right\| \right. \\
&\quad \left. + \sup_{x \neq y} \|\phi\| \frac{\left\| \sum_{j=1}^m f_j(x) a_j - \sum_{j=1}^m f_j(y) a_j \right\|}{d^\alpha(x, y)} \right] \\
&\leq \sup_{x \in K} \left\| \sum_{j=1}^m f_j(x) a_j \right\| + \sup_{x \neq y} \frac{\left\| \sum_{j=1}^m f_j(x) a_j - \sum_{j=1}^m f_j(y) a_j \right\|}{d^\alpha(x, y)} \\
&= \left\| \sum_{j=1}^m f_j a_j \right\|_\infty + p_\alpha \left(\sum_{j=1}^m f_j a_j \right) = \left\| \sum_{j=1}^m f_j a_j \right\|_\alpha = \left\| T \left(\sum_{j=1}^m f_j \otimes a_j \right) \right\|_\alpha \\
&= \|TG\|_\alpha \implies \|G\|_\varepsilon \leq \|TG\|_\alpha.
\end{aligned}$$

Now let $\gamma > 0$ be arbitrary, such that $\|TG\|_\alpha > \gamma$. Then $\left\| \sum_{j=1}^m f_j a_j \right\|_\alpha > \gamma$, and so we have

$$\begin{aligned}
&\left\| \sum_{j=1}^m f_j a_j \right\|_\infty + p_\alpha \left(\sum_{j=1}^m f_j a_j \right) > \gamma \\
\Rightarrow \sup_{x \in K} \left\| \sum_{j=1}^m f_j(x) a_j \right\| + \sup_{x \neq y} \frac{\left\| \sum_{j=1}^m f_j(x) a_j - \sum_{j=1}^m f_j(y) a_j \right\|}{d^\alpha(x, y)} > \gamma \\
\Rightarrow \sup_{\phi \in A^*, \|\phi\| \leq 1} \left[\sup_{x \in K} \left| \sum_{j=1}^m f_j(x) \phi(a_j) \right| \right. \\
&\quad \left. + \sup_{x \neq y} \frac{\left| \sum_{j=1}^m f_j(x) \phi(a_j) - \sum_{j=1}^m f_j(y) \phi(a_j) \right|}{d^\alpha(x, y)} \right] > \gamma \\
\Rightarrow \sup_{\phi \in A^*, \|\phi\| \leq 1} \left[\left\| \sum_{j=1}^m f_j \phi(a_j) \right\|_\infty + p_\alpha \left(\sum_{j=1}^m f_j \phi(a_j) \right) \right] > \gamma \\
\Rightarrow \sup_{\phi \in A^*, \|\phi\| \leq 1} \left[\left\| (\text{id} \otimes \phi) \left(\sum_{j=1}^m f_j \otimes a_j \right) \right\|_\infty + p_\alpha \left((\text{id} \otimes \phi) \left(\sum_{j=1}^m f_j \otimes a_j \right) \right) \right] > \gamma \\
\Rightarrow \sup_{\phi \in A^*, \|\phi\| \leq 1} \left\| (\text{id} \otimes \phi) \left(\sum_{j=1}^m f_j \otimes a_j \right) \right\|_\alpha > \gamma \\
\Rightarrow \left\| \sum_{j=1}^m f_j \otimes a_j \right\|_\varepsilon > \gamma \implies \|G\|_\varepsilon > \gamma.
\end{aligned}$$

Since $\gamma > 0$ is arbitrary, then we have $\|TG\|_\alpha \leq \|G\|_\varepsilon$. Therefore $\|TG\|_\alpha = \|G\|_\varepsilon$, and this implies that T is a linear isometry map. So T one-one and continuous map. Now, we show that T is a onto map. For this, we show that the range of T , R_T is a closed and dense subset of $L^\alpha(K, A)$. It is easy to see that R_T is closed. Let $f \in L^\alpha(K, A)$ and $\gamma > 0$. There exist $a_1, \dots, a_n \in A$ such that $X := f(K) \subset \bigcup_{i=1}^n B(a_i, \gamma)$. Set $U_j = f^{-1}(B(a_j, \gamma))$ where $j = 1, \dots, n$. Then there

exist $f_1, \dots, f_n \in L^\alpha(K, A)$ and $\sigma \in A^*$ such that $\text{supp}(f_j) \subset U_j$ for $j = 1, \dots, n$ and $\sigma \circ (f_1 + \dots + f_n) = 1$. For every $x \in K$ we have,

$$\begin{aligned} & \|f(x) - ((\sigma \circ f_1)a_1 + \dots + (\sigma \circ f_n)a_n)(x)\| \\ &= \|f(x)((\sigma \circ f_1)(x) + \dots + (\sigma \circ f_n)(x)) \\ &\quad - ((\sigma \circ f_1)(x)a_1 + \dots + (\sigma \circ f_n)(x)a_n)\| \\ &= \|(\sigma \circ f_1)(x)(f(x) - a_1) + \dots + (\sigma \circ f_n)(x)(f(x) - a_n)\| \\ &\leq \sum_{i=1}^n |(\sigma \circ f_i)(x)| \|f(x) - a_i\| < \gamma, \end{aligned}$$

since $\text{supp } f_j \subset U_j$. Therefore,

$$\|f - ((\sigma \circ f_1)a_1 + \dots + (\sigma \circ f_n)a_n)\|_\alpha < \gamma.$$

This implies that

$$\left\| f - \sum_{i=1}^n T(\sigma \circ f_i \otimes a_i) \right\|_\alpha < \gamma.$$

We conclude that $\bar{R}_T = L^\alpha(K, A)$. So $R_T = L^\alpha(K, A)$, since R_T is closed. Hence T is a onto map. Also by product \bullet on $L^\alpha(K) \check{\otimes} A$

$$(f \otimes a) \bullet (g \otimes b) = fg \otimes ab \quad (f, g \in L^\alpha(K), a, b \in A),$$

clearly T is homomorphism. □

Furthermore T is open map, for this purpose, let τ and τ' be topologies on $L^\alpha(K) \check{\otimes} A$ and $L^\alpha(K, A)$ respectively. Let $U \in \tau$, we show that $T(U) \in \tau'$. Let p be a limit point in $L^\alpha(K, A) \setminus T(U)$. Then there exists a sequence $\{p_n\}$ in $L^\alpha(K, A) \setminus T(U)$ converges to p . Since T is onto, there is a sequence $\{q_n\}$ and q in $L^\alpha(K) \check{\otimes} A$ such that $T(q_n) = p_n$ and $Tq = p$. Therefore $T(q_n)$ converges to p in $L^\alpha(K)$. Since $q_n \in L^\alpha(K) \check{\otimes} A$, we can find $m \in \mathbb{N}$, $f_j^{(n)} \in L^\alpha(K)$ and $a_j^{(n)} \in A$ such that whenever $1 \leq j \leq m$ we have

$$(1) \quad T(q_n) = \sum_{j=1}^m f_j^{(n)} a_j^{(n)}.$$

Also, since $q \in L^\alpha(K) \check{\otimes} A$ there exist $r \in \mathbb{N}$, $g_i \in L^\alpha(K)$ and $b_i \in A$ such that

$$(2) \quad p = T(q) = \sum_{i=1}^r g_i b_i.$$

Since $\|T(q_n) - p\|_\alpha \rightarrow 0$ as $n \rightarrow \infty$, for every positive number γ there exists a positive integer N such that

$$(3) \quad \left\| \sum_{j=1}^m f_j^{(n)} a_j^{(n)} - \sum_{i=1}^r g_i b_i \right\|_\alpha < \gamma,$$

when $n \geq N$. By applying (3), we have

$$\begin{aligned} & \sup_{(x \in K)} \left\| \sum_{j=1}^m f_j^{(n)}(x) a_j^{(n)} - \sum_{i=1}^r g_i(x) b_i \right\| + \sup_{(x \neq y)} \frac{1}{d(x, y)^\alpha} \\ & \times \left\| \sum_{j=1}^m f_j^{(n)}(x) a_j^{(n)} - \sum_{i=1}^r g_i(x) b_i - \sum_{j=1}^m f_j^{(n)}(y) a_j^{(n)} + \sum_{i=1}^r g_i(y) b_i \right\| < \gamma. \end{aligned}$$

Therefore if $\sigma \in A^*$ with $\|\sigma\| \leq 1$ then

$$\begin{aligned} & \sup_{(x \in K)} \left\| \sum_{j=1}^m f_j^{(n)}(x) \sigma(a_j^{(n)}) - \sum_{i=1}^r g_i(x) \sigma(b_i) \right\| + \sup_{(x \neq y)} \frac{1}{d(x, y)^\alpha} \\ & \times \left\| \sum_{j=1}^m f_j^{(n)}(x) \sigma(a_j^{(n)}) - \sum_{i=1}^r g_i(x) \sigma(b_i) - \sum_{j=1}^m f_j^{(n)}(y) \sigma(a_j^{(n)}) + \sum_{i=1}^r g_i(y) \sigma(b_i) \right\| < \gamma. \end{aligned}$$

This implies that

$$(4) \quad \left\| \sum_{j=1}^m f_j^{(n)} \sigma(a_j^{(n)}) - \sum_{i=1}^r g_i \sigma(b_i) \right\|_\alpha < \gamma$$

Now by using (4), for every $\phi \in L^\alpha(K)^*$ with $\|\phi\|_\alpha \leq 1$ we have,

$$\left| \phi \left(\sum_{j=1}^m f_j^{(n)} \sigma(a_j^{(n)}) - \sum_{i=1}^r g_i \sigma(b_i) \right) \right| < \gamma,$$

hence

$$(5) \quad \left| \sum_{j=1}^m \phi(f_j^{(n)}) \sigma(a_j^{(n)}) - \sum_{i=1}^r \phi(g_i) \sigma(b_i) \right| < \gamma,$$

By (5), we conclude

$$(6) \quad \sup \left| \sum_{j=1}^m \phi(f_j^{(n)}) \sigma(a_j^{(n)}) - \sum_{i=1}^r \phi(g_i) \sigma(b_i) \right| < \gamma, \quad \|\sigma\| \leq 1, \|\phi\|_\alpha \leq 1.$$

Therefore $\|q_n - q\|_\epsilon \leq \gamma$ and hence $q_n \rightarrow q$ or $q_n \rightarrow T^{-1}p$ in $L^\alpha(K) \check{\otimes} A$. This show that $p \in T(U)^c$.

Remark 2.7. By using the above theorem we can prove that $l^\alpha(K, A) \cong l^\alpha(K) \check{\otimes} A$.

3. (WEAK) AMENABILITY OF $L^\alpha(K, A)$

Let A be a Banach algebra and X be a Banach A -module over F . The linear map $D: A \rightarrow X$ is called an X -derivation on A , if $D(ab) = D(a) \cdot b + a \cdot D(b)$, for every $a, b \in A$. The set of all continues X -derivations on A is a vector space over F which is denoted by $Z^1(A, X)$. For each $x \in X$ the map $\delta_x: A \rightarrow X$, defined by $\delta_x(a) = a \cdot x - x \cdot a$, is a continues X -derivation on A . The X -derivation $D: A \rightarrow X$ is called an inner derivation on A if there exists an $x \in X$ such that $D = \delta_x$. The set of all inner X -derivations on A is a linear subspace of $Z^1(A, X)$

which is denoted by $B^1(A, X)$. The quotient space $Z^1(A, X)/B^1(A, X)$ is denoted by $H^1(A, X)$ and is called the first cohomology group of A with coefficients in X .

Definition 3.1. The Banach algebra A over F is called amenable if for every Banach A -module X over F , $H^1(A, X^*) = \{0\}$. The Banach algebra A over F is called weakly amenable if $H^1(A, A^*) = \{0\}$.

The notion of amenability of Banach algebras were first introduced by B. E. Johnson in 1972 [8]. Bade, Curtis and Dales [2], studied the (weak) amenability of Lipschitz algebras in 1987 [2]. In this section, we study the (weak) amenability of $L^\alpha(K, A)$.

For every Banach algebra B , let Φ_B be the space of maximal ideal of B .

Definition 3.2. Let A be a commutative Banach algebra and let $\phi \in \Phi_A \cup \{0\}$. The non-zero linear functional D on A is called point derivation at ϕ if

$$D(ab) = \phi(a)D(b) + \phi(b)D(a), \quad (a, b \in A).$$

Lemma 3.3. For each non-isolated point $x \in K$ and $\sigma \in A^*$, if the map $\phi: L^\alpha(K, A) \rightarrow \mathbb{C}$ is given by

$$\phi(f) = (\sigma \circ f)(x), \quad (f \in L^\alpha(K, A))$$

then $\phi \in \Phi_{L^\alpha(K, A)}$.

Proof. Obvious. □

Let (K, d) be a fixed non-empty compact metric space, set

$$\Delta = \{(x, y) \in K \times K : x = y\}, \quad W = K \times K - \Delta.$$

We now examine the amenability and weak amenability of Lipschitz operators algebras $L^\alpha(K, A)$ and $l^\alpha(K, A)$.

Theorem 3.4. Let (K, d) be an infinite compact metric space and take $\alpha \in (0, 1]$. Then $L^\alpha(K, A)$ is not weakly amenable.

Proof. Let x be a non-isolated point in K . We define

$$W_x := \left\{ \{(x_n, y_n)\}_{n=1}^\infty : (x_n, y_n) \in W, (x_n, y_n) \rightarrow (x, x) \text{ as } n \rightarrow \infty \right\}.$$

For the net $w = \{(x_n, y_n)\}_{n=1}^\infty$ in W_x and $\sigma \in A^*$, we put

$$\bar{w}(f) = \frac{(\sigma \circ f)(x_n) - (\sigma \circ f)(y_n)}{d(x_n, y_n)^\alpha}, \quad (f \in L^\alpha(K, A))$$

then $\|\bar{w}(f)\|_\infty \leq \|\sigma\| \|f\|_\alpha$. Hence, \bar{w} is continous. Now set

$$D_w(f) = \text{LIM} (\bar{w}(f)), \quad (f \in L^\alpha(K, A)),$$

where $\text{LIM}(\cdot)$ is Banach limit [12]. We show that the linear map D_w is a non-zero point derivation at ϕ , which ϕ is given by Lemma 6. We have,

$$\begin{aligned} D_w(fg) &= \text{LIM}(\overline{w}(fg)) \\ &= \text{LIM} \frac{(\sigma \circ fg)(x_n) - (\sigma \circ fg)(y_n)}{d(x_n, y_n)^\alpha} \\ &= \text{LIM} \frac{1}{d(x_n, y_n)^\alpha} [\sigma \circ (f(x_n)g(x_n) - f(x_n)g(y_n))] \\ &= \text{LIM} \frac{1}{d(x_n, y_n)^\alpha} [\sigma \circ (f(x_n)(g(x_n) - g(y_n)) + g(y_n)(f(x_n) - f(y_n)))] \\ &= (\sigma \circ f)(x) \text{LIM}(\overline{w}(g)) + (\sigma \circ g)(x) \text{LIM}(\overline{w}(f)) \\ &= \phi(f)D_w(g) + \phi(g)D_w(f) \end{aligned}$$

Therefore, by the continuity f, g and properties of Banach limit we conclude D_w is a non-zero, continues point derivation at ϕ on $L^\alpha(K, A)$, an so by [5], $L^\alpha(K, A)$ is not weakly amenable. \square

Corollary 3.5. $L^\alpha(K, A)$ is not amenable.

Theorem 3.6. Let $K \subseteq \mathbb{C}$ be an infinite compact set, and take $\alpha \in (0, 1)$. Then $l^\alpha(K, A)$ is not amenable.

Proof. Let $x_0 \in K$. We define

$$M_{x_0} := \{f \in l^\alpha(K, A) : (\sigma \circ f)(x_0) = 0 \quad \forall \sigma \in A^*\}.$$

If $\sigma \in A^*$, then for each $f \in M_{x_0}^2$ we have

$$\frac{(\sigma \circ f)(x)}{d(x, x_0)^{2\alpha}} \longrightarrow 0 \quad \text{as} \quad d(x, x_0) \longrightarrow 0.$$

For $\beta \in (\alpha, 2\alpha)$, set $f_\beta(x) := \eta(d(x, x_0)^\beta)$, $x \in K$ where, the map $\eta: \mathbb{C} \rightarrow A$ defined by $\eta(\lambda) = \lambda \cdot e$. Then $f_\beta \in M_{x_0}$ and $\{f_\beta + M_{x_0}^2 : \beta \in (\alpha, 2\alpha)\}$ is a linearly independent set in $\frac{M_{x_0}}{M_{x_0}^2}$ because x_0 is non-isolated in K . Therefore $M_{x_0}^2$ has infinite codimension in M_{x_0} , and so $M_{x_0} \neq M_{x_0}^2$ then by [5] M_{x_0} has not a bounded approximate identity, and since M_{x_0} is closed ideal in $l^\alpha(K, A)$, then $l^\alpha(K, A)$ is not amenable. \square

Theorem 3.7. Let (K, d) be a compact metric space and A be a unital commutative Banach algebra. If $\frac{1}{2} < \alpha < 1$, then $l^\alpha(\mathbb{T}, A)$ is not weakly amenable, where \mathbb{T} is unit circle in complex plane.

Proof. By Remark 2.7, we have $l^\alpha(\mathbb{T}, A) \cong l^\alpha(\mathbb{T}) \check{\otimes} A$. Since by [5], $l^\alpha(\mathbb{T})$ is not weakly amenable, hence $l^\alpha(\mathbb{T}, A)$ is not weakly amenable. \square

Corollary 3.8. Let A be a finite-dimensional weakly amenable Banach algebra. If $0 < \alpha < \frac{1}{2}$, then $l^\alpha(K, A)$ is weakly amenable.

Proof. By [10], $l^\alpha(K) \hat{\otimes} A$ is weakly amenable. Now by [10], we have $l^\alpha(K) \hat{\otimes} A \cong l^\alpha(K) \check{\otimes} A$ and this implies that $l^\alpha(K) \check{\otimes} A$ is weakly amenable and so $l^\alpha(K, A)$ is weakly amenable. \square

4. HOMOMORPHISMS ON THE $L_A^\alpha(X, B)$

Definition 4.1. Let (X, d) be a compact metric space in C , $\alpha \in (0, 1]$, $(B, \|\cdot\|)$ be a commutative Banach algebra with unit \mathbf{e} , and B^* be the dual space of B , define

$$A(X, B) = \{f \in C(X, B) : \Lambda \circ f \text{ is analytic in interior of } X, \Lambda \in B^*\}$$

$$L_A^\alpha(X, B) = \{f \in L^\alpha(X, B) : \Lambda \circ f \text{ is analytic in interior of } X, \Lambda \in B^*\}$$

$$l_A^\alpha(X, B) = \{f \in l^\alpha(X, B) : \Lambda \circ f \text{ is analytic in interior of } X, \Lambda \in B^*\}$$

In this case, we have

$$L_A^\alpha(X, B) = L^\alpha(X, B) \cap A(X, B)$$

and

$$l_A^\alpha(X, B) = l^\alpha(X, B) \cap A(X, B).$$

So $L_A^\alpha(X, B) \cong L_A^\alpha(X) \check{\otimes} B$ and $l_A^\alpha(X, B) \cong l_A^\alpha(X) \check{\otimes} B$.

Theorem 4.2. Every character χ on $L_A^\alpha(X, B)$ (and $l_A^\alpha(X, B)$) is of form $\chi = \psi \circ \delta_z$ for some character ψ on B and some $z \in X$.

Proof. Since $L_A^\alpha(X, B) \cong L_A^\alpha(X) \check{\otimes} B$, let $j: L^\alpha(X) \rightarrow L_A^\alpha(X, B)$, $h \mapsto h \otimes \mathbf{e}$, be the canonical embedding. Then there is $z \in X$ such that $\chi \circ j$ is the evaluation in z , that is $\chi \circ j = \delta_z$ where $\delta_z(\varphi) = \varphi(z)$. Consider the ideal

$$I := \{f \in L_A^\alpha(X, B) : f(z) = 0\}.$$

We will show that I is contained in the kernel of χ . Given $f \in I$ we define,

$$\varphi(\omega) := \begin{cases} \omega - z & \text{if } \omega \neq z; \\ 0 & \text{if } \omega = z. \end{cases}$$

and

$$g(\omega) := \begin{cases} \frac{f(\omega)}{\omega - z} & \text{if } \omega \neq z; \\ f'(z) & \text{if } \omega = z. \end{cases}$$

Since f has a Taylor series expansion

$$f(\omega) = \sum_{n=1}^{\infty} \frac{f^{(n)}(z)}{n!} (\omega - z)^n$$

around z , it is easy to see that $\Lambda \circ g$ is holomorphic ($\Lambda \in B^*$), and hence $g \in L_A^\alpha(X, B)$. We have

$$\chi(f) = \chi(j(\varphi)g) = (\chi \circ j)(\varphi)\chi(g) = \delta_z(\varphi)\chi(g) = \varphi(z)\chi(g) = 0.$$

The evaluation δ_z is an epimorphism and since $\ker \delta_z = I \subset \ker \chi$, we obtain the desired factorization $\chi = \psi \circ \delta_z$ for some character ψ on B . □

REFERENCES

- [1] Alimohammadi, D., Ebadian, A., *Headberg's theorem in real Lipschitz algebras*, Indian J. Pure Appl. Math. **32** (2001), 1479–1493.
- [2] Bade, W. G., Curtis, P. C., Dales, H. G., *Amenability and weak amenability for Berurling and Lipschitz algebras*, Proc. London Math. Soc. **55** (3) (1987), 359–377.
- [3] Cao, H. X., Xu, Z. B., *Some properties of Lipschitz- α operators*, Acta Math. Sin. (Engl. Ser.) **45** (2) (2002), 279–286.
- [4] Cao, H. X., Zhang, J. H., Xu, Z. B., *Characterizations and extentions of Lipschitz- α operators*, Acta Math. Sin. (Engl. Ser.) **22** (3) (2006), 671–678.
- [5] Dales, H. G., *Banach Algebras and Automatic Continuity*, Clarendon Press, Oxford, 2000.
- [6] Ebadian, A., *Prime ideals in Lipschitz algebras of finite differentiable function*, Honam Math. J. **22** (2000), 21–30.
- [7] Honary, T. G., Mahyar, H., *Approximation in Lipschitz algebras*, Quaest. Math. **23** (2000), 13–19.
- [8] Johnson, B. E., *Cohomology in Banach algebras*, Mem. Amer. Math. Soc. **127** (1972).
- [9] Johnson, B. E., *Lipschitz spaces*, Pacific J. Math. **51** (1975), 177–186.
- [10] Runde, V., *Lectures on Amenability*, Springer, 2001.
- [11] Sherbert, D. R., *Banach algebras of Lipschitz functions*, Pacific J. Math. **3** (1963), 1387–1399.
- [12] Sherbert, D. R., *The structure of ideals and point derivations in Banach algebras of Lipschitz functions*, Trans. Amer. Math. Soc. **111** (1964), 240–272.
- [13] Weaver, N., *Subalgebras of little Lipschitz algebras*, Pacific J. Math. **173** (1996), 283–293.
- [14] Weaver, N., *Lipschitz Algebras*, World Scientific Publishing Co., Inc., River Edge, NJ, 1999.

FACULTY OF BASIC SCIENCE, SCIENCE AND RESEARCH BRUNCH
ISLAMIC AZAD UNIVERSITY (IAU), TEHRAN, IRAN
E-mail: a-shokri@iau-ahar.ac.ir

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE
URMIA UNIVERSITY, URMIA, IRAN
E-mail: a.ebadian@urmia.ac.ir