

Ahmad T. Ali; Rafael López

Timelike B_2 -slant helices in Minkowski space E_1^4

Archivum Mathematicum, Vol. 46 (2010), No. 1, 39--46

Persistent URL: <http://dml.cz/dmlcz/139994>

Terms of use:

© Masaryk University, 2010

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

TIMELIKE B_2 -SLANT HELICES IN MINKOWSKI SPACE \mathbf{E}_1^4

AHMAD T. ALI AND RAFAEL LÓPEZ

ABSTRACT. We consider a unit speed timelike curve α in Minkowski 4-space \mathbf{E}_1^4 and denote the Frenet frame of α by $\{\mathbf{T}, \mathbf{N}, \mathbf{B}_1, \mathbf{B}_2\}$. We say that α is a generalized helix if one of the unit vector fields of the Frenet frame has constant scalar product with a fixed direction U of \mathbf{E}_1^4 . In this work we study those helices where the function $\langle \mathbf{B}_2, U \rangle$ is constant and we give different characterizations of such curves.

1. INTRODUCTION AND STATEMENT OF RESULTS

A helix in Euclidean 3-space \mathbf{E}^3 is a curve where the tangent lines make a constant angle with a fixed direction. A helix curve is characterized by the fact that the ratio τ/κ is constant along the curve, where τ and κ denote the torsion and the curvature, respectively. Helices are well known curves in classical differential geometry of space curves [8] and we refer to the reader for recent works on this type of curves [4, 12]. Recently, Izumiya and Takeuchi have introduced the concept of slant helix by saying that the normal lines make a constant angle with a fixed direction [5]. They characterize a slant helix if and only if the function

$$(1) \quad \frac{\kappa^2}{(\kappa^2 + \tau^2)^{3/2}} \left(\frac{\tau}{\kappa} \right)'$$

is constant. The article [5] motivated generalizations in a twofold sense: first, by considering arbitrary dimension of Euclidean space [7, 9]; second, by considering analogous problems in other ambient spaces, for example, in Minkowski space \mathbf{E}_1^n [1, 3, 6, 11, 13].

In this work we consider the generalization of the concept of helix in Minkowski 4-space, when the helix is a timelike curve. We denote by \mathbf{E}_1^4 the Minkowski 4-space, that is, \mathbf{E}_1^4 is the real vector space \mathbb{R}^4 endowed with the standard Lorentzian metric

$$\langle \cdot, \cdot \rangle = -dx_1^2 + dx_2^2 + dx_3^2 + dx_4^2,$$

where (x_1, x_2, x_3, x_4) is a rectangular coordinate system of \mathbb{R}^4 . An arbitrary vector $v \in \mathbf{E}_1^4$ is said spacelike (resp. timelike, lightlike) if $\langle v, v \rangle > 0$ or $v = 0$ (resp. $\langle v, v \rangle < 0$, $\langle v, v \rangle = 0$ and $v \neq 0$). Let $\alpha: I \subset \mathbb{R} \rightarrow \mathbf{E}_1^4$ be a (differentiable) curve

2000 *Mathematics Subject Classification*: primary 53C50; secondary 53B30.

Key words and phrases: Minkowski space, timelike curve, Frenet equations, slant helix.

Partially supported by MEC-FEDER grant no. MTM2007-61775 and Junta de Andalucía grant no. P06-FQM-01642.

Received January 27, 2009, revised October 2009. Editor J. Slovák.

with $\alpha'(t) \neq 0$, where $\alpha'(t) = d\alpha/dt(t)$. The curve α is said timelike if all its velocity vectors $\alpha'(t)$ are timelike. Then it is possible to re-parametrize α by a new parameter s , in such way that $\langle \alpha'(s), \alpha'(s) \rangle = -1$, for any $s \in I$. We say then that α is a unit speed timelike curve.

Consider $\alpha = \alpha(s)$ a unit speed timelike curve in \mathbf{E}_1^4 . Let $\{\mathbf{T}(s), \mathbf{N}(s), \mathbf{B}_1(s), \mathbf{B}_2(s)\}$ be the moving frame along α , where $\mathbf{T}, \mathbf{N}, \mathbf{B}_1$ and \mathbf{B}_2 denote the tangent, the principal normal, the first binormal and second binormal vector fields, respectively. Here $\mathbf{T}(s), \mathbf{N}(s), \mathbf{B}_1(s)$ and $\mathbf{B}_2(s)$ are mutually orthogonal vectors satisfying

$$\langle \mathbf{T}, \mathbf{T} \rangle = -1, \langle \mathbf{N}, \mathbf{N} \rangle = \langle \mathbf{B}_1, \mathbf{B}_1 \rangle = \langle \mathbf{B}_2, \mathbf{B}_2 \rangle = 1.$$

Then the Frenet equations for α are given by

$$(2) \quad \begin{bmatrix} \mathbf{T}' \\ \mathbf{N}' \\ \mathbf{B}_1' \\ \mathbf{B}_2' \end{bmatrix} = \begin{bmatrix} 0 & \kappa_1 & 0 & 0 \\ \kappa_1 & 0 & \kappa_2 & 0 \\ 0 & -\kappa_2 & 0 & \kappa_3 \\ 0 & 0 & -\kappa_3 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B}_1 \\ \mathbf{B}_2 \end{bmatrix}.$$

Recall the functions $\kappa_1(s), \kappa_2(s)$ and $\kappa_3(s)$ are called respectively, the first, the second and the third curvatures of α . If $\kappa_3(s) = 0$ for any $s \in I$, then $\mathbf{B}_2(s)$ is a constant vector B and the curve α lies in a three-dimensional affine subspace orthogonal to B , which is isometric to the Minkowski 3-space \mathbf{E}_1^3 .

We will assume throughout this work that all the three curvatures satisfy $\kappa_i(s) \neq 0$ for any $s \in I, 1 \leq i \leq 3$.

Definition 1.1. A unit speed timelike curve $\alpha: I \rightarrow \mathbf{E}_1^4$ is said to be a generalized (timelike) helix if there exists a constant vector field U different from zero and a vector field $X \in \{\mathbf{T}, \mathbf{N}, \mathbf{B}_1, \mathbf{B}_2\}$ such that the function

$$s \longmapsto \langle X(s), U \rangle, \quad s \in I$$

is constant.

In this work we are interested in generalized timelike helices in \mathbf{E}_1^4 where the function $\langle \mathbf{B}_2, U \rangle$ is constant. Motivated by the concept of slant helix in \mathbf{E}^4 [9], we give the following

Definition 1.2. A unit speed timelike curve α is called a B_2 -slant helix if there exists a constant vector field U such that the function $\langle \mathbf{B}_2(s), U \rangle$ is constant.

Our main result in this work follows similar ideas as in [6] for timelike helices in \mathbf{E}_1^4 . In this sense, we have the following characterization of B_2 -slant helices in the spirit of the one given in equation (1) for a slant helix in \mathbf{E}^3 :

A unit speed timelike curve in \mathbf{E}_1^4 is a B_2 -slant helix if and only if the function

$$\frac{1}{\kappa_1^2} \left(\frac{\kappa_3}{\kappa_2} \right)' ^2 - \left(\frac{\kappa_3}{\kappa_2} \right)^2$$

is constant.

When α is a lightlike curve, similar computations have been given by Erdogan and Yilmaz in [2].

2. BASIC EQUATIONS OF TIMELIKE HELICES

Let α be a unit speed timelike curve in \mathbf{E}_1^4 and let U be a unit constant vector field in \mathbf{E}_1^4 . For each $s \in I$, the vector U is expressed as linear combination of the orthonormal basis $\{\mathbf{T}(s), \mathbf{N}(s), \mathbf{B}_1(s), \mathbf{B}_2(s)\}$. Consider the differentiable functions a_i , $1 \leq i \leq 4$,

$$(3) \quad U = a_1(s)\mathbf{T}(s) + a_2(s)\mathbf{N}(s) + a_3(s)\mathbf{B}_1(s) + a_4(s)\mathbf{B}_2(s), \quad s \in I,$$

that is,

$$a_1 = -\langle \mathbf{T}, U \rangle, \quad a_2 = \langle \mathbf{N}, U \rangle, \quad a_3 = \langle \mathbf{B}_1, U \rangle, \quad a_4 = \langle \mathbf{B}_2, U \rangle.$$

Because the vector field U is constant, a differentiation in (3) together (2) gives the following ordinary differential equation system

$$(4) \quad \begin{cases} a_1' + \kappa_1 a_2 = 0 \\ a_2' + \kappa_1 a_1 - \kappa_2 a_3 = 0 \\ a_3' + \kappa_2 a_2 - \kappa_3 a_4 = 0 \\ a_4' + \kappa_3 a_3 = 0 \end{cases}$$

In the case that U is spacelike (resp. timelike), we will assume that $\langle U, U \rangle = 1$ (resp. -1). This means that the constant M defined by

$$(5) \quad M := \langle U, U \rangle = -a_1^2 + a_2^2 + a_3^2 + a_4^2$$

is 1, -1 or 0 depending if U is spacelike, timelike or lightlike, respectively.

We now suppose that α is a generalized helix. This means that there exists i , $1 \leq i \leq 4$, such that the function $a_i = a_i(s)$ is constant. Thus in the system (4) we have four differential equations and three derivatives of functions.

The first case that appears is that the function a_1 is constant, that is, the function $\langle \mathbf{T}(s), U \rangle$ is constant. If U is timelike, that is, the tangent lines of α make a constant (hyperbolic) angle with a fixed timelike direction, the curve α is called a timelike cylindrical helix [6]. Then it is known that α is timelike cylindrical helix if and only if the function

$$\frac{1}{\kappa_3^2} \left(\frac{\kappa_1}{\kappa_2} \right)' ^2 + \left(\frac{\kappa_1}{\kappa_2} \right)^2$$

is constant [6].

However the hypothesis that U is timelike can be dropped and we can assume that U has any causal character, as for example, spacelike or lightlike. We explain this situation. In Euclidean space one speaks on the angle that makes a fixed direction with the tangent lines (cylindrical helices) or the normal lines (slant helices). In Minkowski space, one can only speak about the angle between two vectors $\{u, v\}$ if both are timelike and the belong to the same timecone (hyperbolic angle). See [10, page 144]. This is the reason to avoid any reference about ‘angles’ in Definition 1.1.

Suppose now that the function $\langle \mathbf{T}(s), U \rangle$ is constant, independent on the causal character of U . From the expression of U in (3), we know that $a_1' = 0$ and by

using (4), we obtain $a_2 = 0$ and

$$a_3 = \frac{\kappa_1}{\kappa_2} a_1, \quad a_3' = \kappa_3 a_4, \quad a_4' + \kappa_3 a_3 = 0.$$

Consider the change of variable $t(s) = \int_0^s \kappa_3(x) dx$. Then $\frac{dt}{ds}(s) = \kappa_3(s)$ and the last two above equations write as $a_3''(t) + a_3(t) = a_4''(t) + a_4(t) = 0$. Then one obtains that there exist constants A and B such that

$$\begin{aligned} a_3(s) &= A \cos \int_0^s \kappa_3(s) ds + B \sin \int_0^s \kappa_3(s) ds \\ a_4(s) &= -A \sin \int_0^s \kappa_3(s) ds + B \cos \int_0^s \kappa_3(s) ds. \end{aligned}$$

Since $a_3^2 + a_4^2 = \langle U, U \rangle + a_1^2$ is constant, and

$$a_4 = \frac{1}{\kappa_3} a_3' = \frac{1}{\kappa_3} \left(\frac{\kappa_1}{\kappa_2} \right)' a_1,$$

it follows that

$$\frac{1}{\kappa_3^2} \left(\frac{\kappa_1}{\kappa_2} \right)'^2 + \left(\frac{\kappa_1}{\kappa_2} \right)^2 = \text{constant}.$$

Thus we have proved the following theorem.

Theorem 2.1. *Let α be a unit speed timelike curve in \mathbf{E}_1^4 . Then the function $\langle \mathbf{T}(s), U \rangle$ is constant for a fixed constant vector field U if and only if the function*

$$\frac{1}{\kappa_3^2} \left(\frac{\kappa_1}{\kappa_2} \right)'^2 + \left(\frac{\kappa_1}{\kappa_2} \right)^2$$

is constant.

When U is a timelike constant vector field, we re-discover the result given in [6].

3. TIMELIKE B_2 -SLANT HELICES

Let α be a B_2 -slant helix, that is, a unit speed timelike curve in \mathbf{E}_1^4 such that the function $\langle \mathbf{B}_2(s), U \rangle$, $s \in I$, is constant for a fixed constant vector field U . We point out that U can be of any causal character. In the particular case that U is spacelike, and since \mathbf{B}_2 is too, we can say that a B_2 -slant helix is a timelike curve whose second binormal lines make a constant angle with a fixed (spacelike) direction.

Using the system (3), the fact that α is a B_2 -slant helix means that the function a_4 is constant. Then (4) gives $a_3 = 0$ and (3) writes as

$$(6) \quad U = a_1(s) \mathbf{T}(s) + a_2(s) \mathbf{N}(s) + a_4 \mathbf{B}_2(s), \quad a_4 \in \mathbb{R}$$

where

$$(7) \quad a_2 = \frac{\kappa_3}{\kappa_2} a_4 = -\frac{1}{\kappa_1} a_1', \quad a_4' + \kappa_1 a_1 = 0.$$

We remark that $a_4 \neq 0$: on the contrary, and from (4), we conclude $a_i = 0$, $1 \leq i \leq 4$, that is, $U = 0$: contradiction.

It follows from (7) that the function a_1 satisfies the following second order differential equation:

$$\frac{1}{\kappa_1} \frac{d}{ds} \left(\frac{1}{\kappa_1} a_1' \right) - a_1 = 0.$$

If we change variables in the above equation as $\frac{1}{\kappa_1} \frac{d}{ds} = \frac{d}{dt}$, that is, $t = \int_0^s \kappa_1(s) ds$, then we get

$$\frac{d^2 a_1}{dt^2} - a_1 = 0.$$

The general solution of this equation is

$$(8) \quad a_1(s) = A \cosh \int_0^s \kappa_1(s) ds + B \sinh \int_0^s \kappa_1(s) ds,$$

where A and B are arbitrary constants. From (7) and (8) we have

$$(9) \quad a_2(s) = -A \sinh \int_0^s \kappa_1(s) ds - B \cosh \int_0^s \kappa_1(s) ds.$$

The above expressions of a_1 and a_2 give

$$(10) \quad \begin{aligned} A &= - \left[\frac{\kappa_3}{\kappa_2} \sinh \int_0^s \kappa_1(s) ds + \frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' \cosh \int_0^s \kappa_1(s) ds \right] a_4, \\ B &= - \left[\frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' \sinh \int_0^s \kappa_1(s) ds + \frac{\kappa_3}{\kappa_2} \cosh \int_0^s \kappa_1(s) ds \right] a_4. \end{aligned}$$

From (10),

$$A^2 - B^2 = \left[\frac{1}{\kappa_1^2} \left(\frac{\kappa_3}{\kappa_2} \right)' ^2 - \frac{\kappa_3^2}{\kappa_2^2} \right] a_4^2.$$

Therefore

$$(11) \quad \frac{1}{\kappa_1^2} \left(\frac{\kappa_3}{\kappa_2} \right)' ^2 - \frac{\kappa_3^2}{\kappa_2^2} = \text{constant} := m.$$

Conversely, if the condition (11) is satisfied for a timelike curve, then we can always find a constant vector field U such that the function $\langle \mathbf{B}_2(s), U \rangle$ is constant: it suffices if we define

$$U = \left[- \frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' \mathbf{T} + \frac{\kappa_3}{\kappa_2} \mathbf{N} + \mathbf{B}_2 \right].$$

By taking account of the differentiation of (11) and the Frenet equations (2), we have that $\frac{dU}{ds} = 0$ and this means that U is a constant vector. On the other hand, $\langle \mathbf{B}_2(s), U \rangle = 1$. The above computations can be summarized as follows:

Theorem 3.1. *Let α be a unit speed timelike curve in \mathbf{E}_1^4 . Then α is a B_2 -slant helix if and only if the function*

$$\frac{1}{\kappa_1^2} \left(\frac{\kappa_3}{\kappa_2} \right)' ^2 - \left(\frac{\kappa_3}{\kappa_2} \right)^2$$

is constant.

From (5), (8) and (9) we get

$$A^2 - B^2 = a_4^2 - M = a_4^2 m.$$

Thus, the sign of the constant m agrees with the one $A^2 - B^2$. So, if U is timelike or lightlike, m is positive. If U is spacelike, then the sign of m depends on $a_4^2 - 1$. For example, $m = 0$ if and only if $a_4^2 = 1$. With similar computations as above, we have

Corollary 3.2. *Let α be a unit speed timelike curve in \mathbf{E}_1^4 and let U be a unit spacelike constant vector field. Then $\langle \mathbf{B}_2(s), U \rangle^2 = 1$ for any $s \in I$ if and only if there exists a constant A such that*

$$\frac{\kappa_3}{\kappa_2}(s) = A \exp\left(\int_0^s \kappa_1(t) dt\right).$$

As a consequence of Theorem 3.1, we obtain other characterization of B_2 -slant helices. The first one is the following

Corollary 3.3. *Let α be a unit speed timelike curve in \mathbf{E}_1^4 . Then α is a B_2 -slant helix if and only if there exists real numbers C and D such that*

$$(12) \quad \frac{\kappa_3}{\kappa_2}(s) = C \sinh \int_0^s \kappa_1(s) ds + D \cosh \int_0^s \kappa_1(s) ds,$$

Proof. Assume that α is a B_2 -slant helix. From (7) and (9), the choice $C = -A/a_4$ and $D = -B/a_4$ yields (12).

We now suppose that (12) is satisfied. A straightforward computation gives

$$\frac{1}{\kappa_1^2} \left(\frac{\kappa_3}{\kappa_2} \right)' ^2 - \left(\frac{\kappa_3}{\kappa_2} \right)^2 = C^2 - D^2.$$

We now use Theorem 3.1. □

We end this section with a new characterization for B_2 -slant helices. Let now assume that α is a B_2 -slant helix in \mathbf{E}_1^4 . By differentiation (11) with respect to s we get

$$(13) \quad \frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' \left[\frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' \right]' - \left(\frac{\kappa_3}{\kappa_2} \right) \left(\frac{\kappa_3}{\kappa_2} \right)' = 0,$$

and hence

$$\frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' = \frac{\left(\frac{\kappa_3}{\kappa_2} \right) \left(\frac{\kappa_3}{\kappa_2} \right)'}{\left[\frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' \right]'},$$

If we define a function $f(s)$ as

$$f(s) = \frac{\left(\frac{\kappa_3}{\kappa_2} \right) \left(\frac{\kappa_3}{\kappa_2} \right)'}{\left[\frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)' \right]'},$$

then

$$(14) \quad f(s) \kappa_1(s) = \left(\frac{\kappa_3}{\kappa_2} \right)'.$$

By using (13) and (14), we have

$$f'(s) = \frac{\kappa_1 \kappa_3}{\kappa_2}.$$

Conversely, consider the function $f(s) = \frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)'$ and assume that $f'(s) = \frac{\kappa_1 \kappa_3}{\kappa_2}$. We compute

$$(15) \quad \frac{d}{ds} \left[\frac{1}{\kappa_1^2} \left(\frac{\kappa_3}{\kappa_2} \right)' ^2 - \frac{\kappa_3^2}{\kappa_2^2} \right] = \frac{d}{ds} \left[f(s)^2 - \frac{f'(s)^2}{\kappa_1^2} \right] := \varphi(s).$$

As $f(s)f'(s) = \left(\frac{\kappa_3}{\kappa_2} \right) \left(\frac{\kappa_3}{\kappa_2} \right)'$ and $f''(s) = \kappa_1' \left(\frac{\kappa_3}{\kappa_2} \right) + \kappa_1 \left(\frac{\kappa_3}{\kappa_2} \right)'$ we obtain

$$f'(s)f''(s) = \kappa_1 \kappa_1' \left(\frac{\kappa_3}{\kappa_2} \right)^2 + \kappa_1^2 \left(\frac{\kappa_3}{\kappa_2} \right) \left(\frac{\kappa_3}{\kappa_2} \right)'.$$

As consequence of above computations

$$\varphi(s) = 2 \left(f(s)f'(s) - \frac{f'(s)f''(s)}{\kappa_1^2} + \frac{\kappa_1' f'(s)^2}{\kappa_1^3} \right) = 0,$$

that is, the function $\frac{1}{\kappa_1^2} \left(\frac{\kappa_3}{\kappa_2} \right)' ^2 - \left(\frac{\kappa_3}{\kappa_2} \right)^2$ is constant. Therefore we have proved the following

Theorem 3.4. *Let α be a unit speed timelike curve in \mathbf{E}_1^4 . Then α is a B_2 -slant helix if and only if the function $f(s) = \frac{1}{\kappa_1} \left(\frac{\kappa_3}{\kappa_2} \right)'$ satisfies $f'(s) = \frac{\kappa_1 \kappa_3}{\kappa_2}$.*

REFERENCES

- [1] Barros, M., *General helices and a theorem of Lancret*, Proc. Amer. Math. Soc. **125** (1997), 1503–1509.
- [2] Erdoğan, M., Yılmaz, G., *Null generalized and slant helices in 4-dimensional Lorentz-Minkowski space*, Int. J. Contemp. Math. Sci. **3** (2008), 1113–1120.
- [3] Ferrandez, A., Gimenez, A., Luca, P., *Null helices in Lorentzian space forms*, Int. J. Mod. Phys. A **16** (2001), 4845–4863.
- [4] Gluck, H., *Higher curvatures of curves in Euclidean space*, Amer. Math. Monthly **73** (1996), 699–704.
- [5] Izumiya, S., Takeuchi, N., *New special curves and developable surfaces*, Turkish J. Math. **28** (2004), 531–537.
- [6] Kocayiğit, H., Önder, M., *Timelike curves of constant slope in Minkowski space \mathbf{E}_1^4* , J. Science Techn. Beykent Univ. **1** (2007), 311–318.
- [7] Kula, L., Yaylı, Y., *On slant helix and its spherical indicatrix*, Appl. Math. Comput. **169** (2005), 600–607.
- [8] Millman, R. S., Parker, G. D., *Elements of differential geometry*, Prentice-Hall Inc., Englewood Cliffs, N. J., 1977.
- [9] Önder, M., Kazaz, M., Kocayiğit, H., Kilic, O., *B_2 -slant helix in Euclidean 4-space E^4* , Int. J. Contemp. Math. Sci. **3** (29) (2008), 1433–1440.
- [10] O’Neill, B., *Semi-Riemannian geometry. With applications to relativity. Pure and Applied Mathematics*, vol. 103, Academic Press, Inc., New York, 1983.
- [11] Petrovic-Torgasev, M., Sucurovic, E., *W-curves in Minkowski spacetime*, Novi Sad J. Math. **32** (2002), 55–65.

- [12] Scofield, P. D., *Curves of constant precession*, Amer. Math. Monthly **102** (1995), 531–537.
- [13] Synge, J. L., *Timelike helices in flat space-time*, Proc. Roy. Irish Acad. Sect. A **65** (1967), 27–42.

AHMAD T. ALI,
AL-AZHAR UNIVERSITY, FACULTY OF SCIENCE, MATHEMATICS DEPARTMENT,
NASR CITY, 11448, CAIRO, EGYPT
E-mail: atali71@yahoo.com

RAFAEL LÓPEZ,
UNIVERSIDAD DE GRANADA,
DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA,
18071 GRANADA, SPAIN
E-mail: rcamino@ugr.es