

A. V. Harutyunyan; Wolfgang Lusky

ω -weighted holomorphic Besov spaces on the unit ball in C^n

Commentationes Mathematicae Universitatis Carolinae, Vol. 52 (2011), No. 1, 37--56

Persistent URL: <http://dml.cz/dmlcz/141427>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2011

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ω -weighted holomorphic Besov spaces on the unit ball in C^n

A.V. HARUTYUNYAN, W. LUSKY

Abstract. The ω -weighted Besov spaces of holomorphic functions on the unit ball B^n in C^n are introduced as follows. Given a function ω of regular variation and $0 < p < \infty$, a function f holomorphic in B^n is said to belong to the Besov space $B_p(\omega)$ if

$$\|f\|_{B_p(\omega)}^p = \int_{B^n} (1 - |z|^2)^p |Df(z)|^p \frac{\omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\nu(z) < +\infty,$$

where $d\nu(z)$ is the volume measure on B^n and D stands for the fractional derivative of f .

The holomorphic Besov space is described in the terms of the corresponding $L_p(\omega)$ space. Some projection theorems and theorems on existence of the inversions of these projections are proved. Also, explicit descriptions of the duals of the considered Besov spaces are given.

Keywords: weighted Besov spaces, unit ball, projection

Classification: 32C37, 47B38, 46T25, 46E15

1. Introduction and basic constructions

Let C^n denote the complex Euclidean space of dimension n . For any points $z = (z_1, \dots, z_n)$, $\zeta = (\zeta_1, \dots, \zeta_n)$ in C^n , we define the inner product as $\langle z, \zeta \rangle = z_1 \bar{\zeta}_1 + \dots + z_n \bar{\zeta}_n$ and note that $|z|^2 = |z_1|^2 + \dots + |z_n|^2$. By $B^n = \{z \in C^n, |z| < 1\}$ and $C^n : S^n = \{z \in C^n, |z| = 1\}$ we denote the open unit ball and its boundary, i.e. the unit sphere, in C^n . Further, by $H(B^n)$ we denote the set of holomorphic functions on B^n and by $H^\infty(B^n)$ the set of bounded holomorphic functions on B^n .

If $f \in H(B^n)$, then $f(z) = \sum_m a_m z^m$ ($z \in B^n$), where the sum is taken over all multiindices $m = (m_1, \dots, m_n)$ with nonnegative integer components m_k and $z^m = z_1^{m_1} \dots z_n^{m_n}$. Assuming that $|m| = m_1 + \dots + m_n$ and putting $f_k(z) = \sum_{|m|=k} a_m z^m$ for any $k \geq 0$, one can rewrite the Taylor expansion of f as

$$(1) \quad f(z) = \sum_{k=0}^{\infty} f_k(z),$$

which is called homogeneous expansion of f , since each f_k is a homogeneous polynomial of the degree k .

Further, for a holomorphic function f the fractional differential D^α is defined as

$$D^\alpha f(z) = \sum_{k=0}^{\infty} (k+1)^\alpha f_k(z),$$

$$D^\alpha f(\bar{z}) = \sum_{k=0}^{\infty} (k+1)^\alpha f_k(\bar{z}), \quad \alpha = (\alpha_1, \dots, \alpha_n), \quad z \in B^n.$$

We consider the inverse operator $D^{-\alpha}$ defined in the standard way:

$$D^{-\alpha} D^\alpha f(z) = f(z).$$

Particularly, if $\alpha = 1$ we set $D^1 f(z) := Df(z)$.

By $d\nu$ we denote the volume measure on B^n , normalized so that $\nu(B^n) = 1$, and by $d\sigma$ the surface measure on S^n , normalized so that $\sigma(S^n) = 1$. Then following lemma, the proof of which can be found in [12] or [15], reveals the connection between these measures.

Lemma 1. *If f is a measurable function with summable modulus over B^n , then*

$$\int_{B^n} f(z) d\nu(z) = 2n \int_0^1 r^{2n-1} dr \int_{S^n} f(r\zeta) d\sigma(\zeta).$$

Definition 1. By S we denote the well-known class of all non-negative measurable functions ω on $(0, 1)$

$$\omega(x) = \exp\left\{\int_x^1 \frac{\varepsilon(u)}{u} du\right\}, \quad x \in (0, 1),$$

where $\varepsilon(u)$ is a bounded measurable function on $(0, 1)$ and $-\alpha_\omega \leq \varepsilon(u) \leq \beta_\omega$.

Note that the functions of S are called *functions of regular variation* (see [13]). Throughout the paper, we shall assume that $\omega \in S$. Besides, for any functions f and g by $f \preceq g$ ($f \succeq g$) we shall mean that $|f(z)| \leq C|g(z)|$ ($|g(z)| \leq C|f(z)|$) and by $f \asymp g$ that $C_1|f(z)| \leq |g(z)| \leq C_2|f(z)|$ for some positive constants C , C_1 , C_2 independent of z .

Proposition 1. *If $1 - |z| \asymp 1 - |w|$, then $\omega(1 - |z|) \asymp \omega(1 - |w|)$.*

PROOF: Let $C_1(1 - r) \leq 1 - |z| \leq C_2(1 - r)$ and $1 - r = \rho$, $1 - |z| = t$. Then we get

$$\omega(\rho) = \omega(t) \exp\left(\int_\rho^t \frac{\varepsilon(u)}{u} du\right) \leq \omega(t) \exp\left(\beta_\omega \int_\rho^t \frac{du}{u}\right) = \left(\frac{t}{\rho}\right)^{\beta_\omega} \omega(t)$$

and

$$\omega(\rho) \geq \omega(t) \exp\left(-\alpha_\omega \int_\rho^t \frac{du}{u}\right) = \left(\frac{t}{\rho}\right)^{-\alpha_\omega} \omega(t)$$

which proves our statement. \square

We define the holomorphic Besov spaces on the unit ball as follows.

Definition 2. Let $p > n + \beta_\omega$. Then a function $f \in H(B^n)$ is said to be in $B_p(\omega)$ if

$$M_f^p(\omega) = \int_{B^n} (1 - |z|^2)^p |Df(z)|^p \frac{\omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\nu(z) < +\infty.$$

We introduce the norm on $H(B^n)$ as $\|f\|_{B_p(\omega)} = M_f(\omega)$ ($|f(0)|$ needs not to be added since $Df = 0$ implies $f = 0$ for a holomorphic function f). Besides, it is easy to check that if $p > 1$, $n = 1$ and $\omega(t) = 1$, then $B_p(\omega)$ becomes the classical Besov space (see [1], [2], [8], [14]).

In particular, for $p = +\infty$ we shall write $B_\infty(\omega) = B_\omega$, where B_ω denotes the ω -weighted Bloch space on the ball (see [4]).

In [9], [10], [11], one can see some other definitions and some characterizations of holomorphic Besov spaces on B^n . For a holomorphic Besov space on the polydisc of C^n , see [5], [6].

Proposition 2. $H^\infty(B^n) \subset B_p(\omega)$ for all $0 < p < \infty$.

PROOF: Let $f \in H^\infty(B^n)$. Then, using the Cauchy inequality in the ball $\tilde{B}(z) = \{\zeta, |\zeta - z| < (1 - |z|)/2\}$ we get $|Df(z)| \preceq (1 - |z|)^{-1}$, and hence $|Df(z)|(1 - |z|) \leq \text{const}$. Thus,

$$\int_{B^n} (1 - |z|^2)^p |Df(z)|^p \frac{\omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\nu(z) < \infty$$

for $p > n + \beta_\omega$, and hence $f \in B_p(\omega)$. \square

By $L_p(\omega)$ we denote the class of all measurable functions on B^n , for which

$$\|f\|_{L_p(\omega)}^p = \int_{B^n} |f(z)|^p \frac{\omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\nu(z) < +\infty$$

and we shall assume that $n + \beta_\omega < 0$. Further, by L_∞ we denote the set of all measurable functions f for which

$$\|f\|_\infty = \sup_{z \in B^n} \{|f(z)|\} < \infty.$$

Besides, we shall consider also the set of holomorphic functions f for which

$$\|f\|_{A^p(\omega)}^p = \int_{B^n} |f(z)|^p \omega(1 - |z|) d\nu(z) < +\infty, \quad \alpha > -1, \quad 0 < p < \infty.$$

The following lemmas will be used for the proof of the main results of the paper.

Lemma 2. *Let $\omega \in S$ and let $f \in B_p(\omega)$ for some $0 < p < \infty$. Then*

$$|Df(z)| \leq \frac{\|f\|_{B_p(\omega)}}{\omega^{1/p}(1-|z|)(1-|z|^2)}, \quad z \in B^n.$$

PROOF: Let $z \in B^n$, and let $B_z^n(r)$ be the disc centered at z , with the radius $r = (1-|z|)/2$. If $w \in B_z^n(r)$, then

$$|w| \leq |w-z| + |z| \leq \frac{1-|z|}{2} + |z| = \frac{1+|z|}{2} < 1.$$

Hence $B_z^n(r) \subset B^n$. Besides, the function $|Df|^p$ is subharmonic and hence

$$|Df(z)|^p \leq \frac{1}{|B_z^n(r)|} \int_{B_z^n(r)} |Df(w)|^p d\nu(w).$$

On the other hand, it is not difficult to show that $1-|z| \asymp 1-|w|$. Therefore, $\omega(1-|z|) \asymp \omega(1-|w|)$ by Proposition 1. Consequently,

$$\begin{aligned} & (1-|z|^2)^p |Df(z)|^p \frac{\omega(1-|z|)}{(1-|z|^2)^{n+1}} \\ & \leq \frac{1}{|B_z^n(r)|} \int_{B_z^n(r)} (1-|w|^2)^p |Df(w)|^p \frac{\omega(1-|w|)}{(1-|w|)^{n+1}} d\nu(w) \leq \frac{\|f\|_{B_p(\omega)}^p}{|B_z^n(r)|}, \end{aligned}$$

and

$$|Df(z)| \leq \frac{\|f\|_{B_p(\omega)}}{\omega^{1/p}(1-|z|)(1-|z|)}$$

since $|B_z^n(r)| \asymp (1-|z|)^{n+1}$. □

Lemma 3. *Let $\omega \in S$ and let $f \in B_p(\omega)$ for some $0 < p < \infty$. Then*

$$|f(z)| \leq C(n, \omega, p) \frac{\|f\|_{B_p(\omega)}}{(1-|z|)^\gamma}, \quad z \in B^n,$$

where $\gamma = \frac{2n-1}{p}$, if $2n \geq p+1$; and $\gamma = 0$ if $2n < p+1$.

PROOF: The function $|Df(z)|^p$ is subharmonic in B^n and, hence,

$$|Df(rz)|^p \leq \int_{S^n} |Df(r\zeta)|^p P(z, \zeta) d\sigma(\zeta),$$

where $P(z, \zeta)$ is the Poisson kernel which satisfies the estimate

$$P(z, \zeta) = \frac{1-|z|^2}{|\zeta-z|^{2n}} \leq \frac{2}{(1-|z|)^{2n-1}}, \quad \zeta \in S^n.$$

Consequently,

$$|Df(rz)|^p \leq \frac{2}{(1-|z|)^{2n-1}} \int_{S^n} |Df(r\zeta)|^p d\sigma(\zeta),$$

and it is clear that

$$\int_{S^n} |Df(r_1\zeta)|^p d\sigma(\zeta) \leq \int_{S^n} |Df(r_2\zeta)|^p d\sigma(\zeta), \quad r_1 \leq r_2.$$

Hence

$$\begin{aligned} & \int_r^1 \frac{(1-t^2)^p \omega(1-t^2)t^{2n-1}}{(1-t^2)^{n+1}} \int_{S^n} |Df(r\zeta)|^p d\sigma(\zeta) dt \\ & \leq \int_r^1 \frac{(1-t^2)^p \omega(1-t^2)t^{2n-1}}{(1-t^2)^{n+1}} \int_{S^n} |Df(t\zeta)|^p d\sigma(\zeta) dt \\ & = \int_r^1 \int_{S^n} |Df(t\zeta)|^p d\sigma(\zeta) \frac{(1-t^2)^p \omega(1-t^2)t^{2n-1}}{(1-t^2)^{n+1}} dt \\ & \leq \int_{B^n} |Df(w)|^p \frac{(1-|w|^2)^p \omega(1-|w|^2)}{(1-|w|^2)^{n+1}} d\nu(w) = \|f\|_{B_p(\omega)}. \end{aligned}$$

Consequently, the following estimate is true:

$$|Df(rz)|^p \leq \frac{2}{(1-|z|^2)^{2n-1}} \left(\int_r^1 \frac{(1-t^2)^p \omega(1-t^2)t^{2n-1}}{(1-t^2)^{n+1}} dt \right)^{-1} \|f\|_{B_p(\omega)}^p.$$

Changing here $rz \mapsto z$ and putting $r = (1+2|z|)/3$, we get

$$|Df(z)| \leq C(n, \omega, p) \frac{\|f\|_{B_p(\omega)}^p}{(1-|z|^2)^{\frac{2n-1}{p}}}.$$

Therefore

$$|f(z)| \leq C(n, \omega, p) \|f\|_{B_p(\omega)} \int_0^1 \frac{dr}{(1-r|z|)^{\frac{2n-1}{p}}} \leq C(n, \omega, p) \frac{\|f\|_{B_p(\omega)}}{(1-|z|)^\gamma},$$

where $\gamma = \frac{2n-1}{p}$, if $2n \geq p+1$; and $\gamma = 0$, if $2n < p+1$. □

Lemma 4. *Let $\omega \in S$ and let $f \in B_p(\omega)$ for some $0 < p \leq 1$. Then*

$$\left(\int_{B^n} |Df(z)| \frac{\omega^{1/p}(1-|z|)}{(1-|z|)^n} d\nu(z) \right)^p \leq \int_{B^n} |Df(z)|^p \frac{(1-|z|)^p \omega(1-|z|)}{(1-|z|)^{n+1}} d\nu(z).$$

PROOF: Noting that $|Df(z)| = |Df(z)|^p |Df(z)|^{1-p}$ and using Lemma 2 we get

$$|Df(z)| \leq |Df(z)|^p \frac{\|f\|_{B_p(\omega)}^{1-p}}{\omega^{(1-p)/p}(1-|z|)(1-|z|)^{1-p}}.$$

Therefore

$$|Df(z)| \frac{(1-|z|)\omega^{1/p}(1-|z|)}{(1-|z|)^{n+1}} \leq |Df(z)|^p \|f\|_{B^p(\omega)}^{1-p} \frac{\omega(1-|z|)(1-|z|)^p}{(1-|z|)^{n+1}},$$

and the proof is completed by integration over B^n . \square

Corollary 1. *If $0 < p < 1$, then $B_p(\omega) \subset B_1(\omega^*)$, where $\omega^*(t) = \omega^{1/p}(t)$.*

Lemma 5. *Let $1 \leq p < \infty$ and let $f \in B_p(\omega)$. Further, let $\alpha > -n/p + \beta_\omega/p$. Then*

$$\int_{B^n} (1-|z|^2)^\alpha |Df(z)| d\nu(z) < \infty.$$

PROOF: For $1 < p < \infty$ Hölder's inequality gives

$$\begin{aligned} & \int_{B^n} (1-|z|^2)^\alpha |Df(z)| d\nu(z) \\ &= \int_{B^n} (1-|z|^2) |Df(z)| \frac{\omega(1-|z|)(1-|z|^2)^{\alpha+n}}{(1-|z|^2)^{n+1}\omega(1-|z|)} d\nu(z) \\ &\leq \left(\int_{B^n} (1-|z|^2)^p |Df(z)|^p \frac{\omega(1-|z|)}{(1-|z|^2)^{n+1}} d\nu(z) \right)^{1/p} \\ &\quad \times \left(\int_{B^n} (1-|z|^2)^{\alpha q+nq-n-1} \omega^{1-q} (1-|z|) d\nu(z) \right)^{1/q}. \end{aligned}$$

It is obvious that if $\alpha > -n/p - \beta_\omega/p$, then

$$\begin{aligned} & \int_{B^n} \omega^{1-q} (1-|z|)(1-|z|)^{\alpha q+n(q-1)-1} d\nu(z) \\ &\leq \int_0^1 (1-r)^{\alpha q+n(q-1)-1-\beta_\omega(1-q)} dr < \infty. \end{aligned}$$

Now, let $p = 1$. Then, evidently,

$$\begin{aligned} & \int_{B^n} (1-|z|^2) |Df(z)| \frac{\omega(1-|z|)(1-|z|^2)^{\alpha+n}}{(1-|z|^2)^{n+1}\omega(1-|z|)} d\nu(z) \\ &\leq \int_{B^n} (1-|z|^2) |Df(z)| \frac{\omega(1-|z|)}{(1-|z|^2)^{n+1}} d\nu(z) = \|f\|_{B_1(\omega)}. \end{aligned}$$

\square

Corollary 2. *Let $1 \leq p < \infty$ and let $f \in B_p(\omega)$. Further, let $\alpha > -n/p - \beta_\omega/p$. Then the function $Df(z)$ lies in the space $A^1(\alpha)$ and can be represented as*

$$(2) \quad Df(z) = C(n, \alpha) \int_{B^n} \frac{(1-|\zeta|^2)^\alpha Df(\zeta)}{(1-\langle z, \zeta \rangle)^{n+1+\alpha}} d\nu(\zeta), \quad z \in B^n,$$

where $C(n, \alpha) = \frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)\Gamma(\alpha+1)}$.

PROOF: (2) is a simple consequence of the well known representation in the one-dimensional case (for details, see [3], [15]). \square

Also the following auxiliary lemma will be used.

Lemma 6. *If $0 < p < \infty$ and $f \in B_p(\omega)$, then*

$$|f(z)| \preceq \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha}{|1 - \langle z, \zeta \rangle|^{n+\alpha}} |Df(\zeta)| d\nu(\zeta)$$

for sufficiently great α .

PROOF: Obviously, $f(z) = \int_0^1 Df(rz) dr$, and by Corollary 2

$$\begin{aligned} f(z) &= C(n, \alpha) \int_0^1 \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha Df(\zeta)}{(1 - r\langle z, \zeta \rangle)^{n+1+\alpha}} d\nu(\zeta) dr \\ &= C(n, \alpha) \int_{B^n} (1 - |\zeta|^2)^\alpha Df(\zeta) \int_0^1 \frac{dr}{(1 - r\langle z, \zeta \rangle)^{n+1+\alpha}} d\nu(\zeta) \\ &= \tilde{C}(n, \alpha) \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha ((1 - \langle z, \zeta \rangle)^{n+\alpha} - 1)}{\langle z, \zeta \rangle (1 - \langle z, \zeta \rangle)^{n+\alpha}} Df(\zeta) d\nu(\zeta). \end{aligned}$$

Hence the desired statement follows. \square

To prove the other main results, we need also the following lemma.

Lemma 7. *Let $D^m f \in B_p(\omega)$ for some $0 < p < +\infty$ and $m \in \mathbb{N}$. Then*

$$|f(z)| \preceq \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha}{|1 - \langle z, \zeta \rangle|^{n+1+\alpha-m}} |D^m f(\zeta)| d\nu(\zeta)$$

for sufficiently great α .

Lemma 8. *For any numbers $\alpha \in \mathbb{N}$ and $\beta > 0$*

$$D^\alpha \left(\frac{1}{(1 - \langle z, \zeta \rangle)^\beta} \right) \asymp \frac{1}{(1 - \langle z, \zeta \rangle)^{\beta+\alpha}}, \quad z, \zeta \in B^n.$$

PROOF: One can see that $D^\alpha f(z) = D^{\alpha-1} Df(z)$ and $Df(z) = Rf(z) + f(z)$, where

$$Rf(z) = \sum_{k=1}^n z_k \frac{\partial f(z)}{\partial z_k}.$$

On the other hand, $R(1 - \langle z, \zeta \rangle)^{-\alpha} = \alpha \langle z, \zeta \rangle (1 - \langle z, \zeta \rangle)^{-\alpha-1}$ and we get the proof of the lemma. \square

2. Description of the spaces $B_p(\omega)$

The following theorem is one of the main results of the paper.

Theorem 1. *For any $0 < p < \infty$, the space $B_p(\omega)$ is a closed subspace of $L_p(\omega)$.*

PROOF: First, we show that if $f \in B_p(\omega)$, then $f \in L_p(\omega) \cap H(B^n)$. Indeed, if $1 < p < \infty$ and $\gamma > 0$, then by Lemma 4 we obtain

$$\begin{aligned} |f(z)|^p &\leq \left(\int_{B^n} \frac{(1 - |\zeta|^2)^{\alpha-\gamma-1} d\nu(\zeta)}{|1 - \langle z, \zeta \rangle|^{\alpha+n}} \right)^{p/q} \\ &\quad \times \int_{B^n} \frac{(1 - |\zeta|^2)^{\alpha-\gamma-1} (1 - |\zeta|^2)^{p+\gamma p}}{|1 - \langle z, \zeta \rangle|^{\alpha+n}} |Df(\zeta)| d\nu(\zeta) \\ &\leq (1 - |z|^2)^{-\gamma p/q} \int_{B^n} \frac{(1 - |\zeta|^2)^{\alpha-\gamma-1} (1 - |\zeta|^2)^{p+\gamma p}}{|1 - \langle z, \zeta \rangle|^{\alpha+n}} |Df(\zeta)| d\nu(\zeta). \end{aligned}$$

Hence

$$\begin{aligned} \|f\|_{L_p(\omega)} &\leq \int_{B^n} |Df(\zeta)|^p (1 - |\zeta|^2)^{(n+1)(p-1)} \int_{B^n} \frac{\omega(1 - |z|) d\nu(z) d\nu(\zeta)}{|1 - \langle z, \zeta \rangle|^{(\alpha+n)p} (1 - |z|^2)^{n+1}} \\ &\leq \int_{B^n} |Df(\zeta)|^p (1 - |\zeta|^2)^p \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|^2)^{n+1}} d\nu(\zeta) = \|f\|_{B_p(\omega)}. \end{aligned}$$

If $0 < p \leq 1$, then by Lemma 4

$$|f(z)|^p \leq \int_{B^n} \frac{|Df(\zeta)|^p (1 - |\zeta|)^{pn+\alpha p+p}}{|1 - \langle z, \zeta \rangle|^{(\alpha+n)p} (1 - |\zeta|^2)^{n+1}} d\nu(\zeta).$$

Consequently,

$$\begin{aligned} \|f\|_{L_p(\omega)} &\leq \int_{B^n} |Df(\zeta)|^p (1 - |\zeta|^2)^{(n+1)(p-1)} \int_{B^n} \frac{\omega(1 - |z|) d\nu(z) d\nu(\zeta)}{|1 - \langle z, \zeta \rangle|^{(\alpha+n)p} (1 - |z|^2)^{n+1}} \\ &\leq \|f\|_{B_p(\omega)} \end{aligned}$$

by [15, Theorem 1.12] and [7, Lemma 1.6].

Next, we show that if $f \in L_p(\omega) \cap H(B^n)$, then $f \in B_p(\omega)$. Indeed, using (2) we obtain

$$|Df(z)|^p \leq \left(\int_{B^n} \frac{(1 - |\zeta|^2)^m |f(\zeta)|}{|1 - \langle z, \zeta \rangle|^{m+n+2}} d\nu(\zeta) \right)^p$$

for sufficiently great m . If $1 < p < \infty$, then by Hölder's inequality

$$\|f\|_{B_p(\omega)} \leq \int_{B^n} (1 - |\zeta|^2)^m |f(\zeta)|^p \int_{B^n} \frac{\omega(1 - |z|) d\nu(z)}{|1 - \langle z, \zeta \rangle|^{m+n+2}} \leq \|f\|_{L_p(\omega)}.$$

If $0 < p \leq 1$, then by Lemma 4 we obtain

$$|Df(z)|^p \preceq \int_{B^n} \frac{|f(\zeta)|^p (1 - |\zeta|)^{p(n+1)+mp}}{|1 - \langle z, \zeta \rangle|^{(m+n+2)p} (1 - |\zeta|^2)^{n+1}} d\nu(\zeta),$$

and hence

$$\begin{aligned} \|f\|_{B_p(\omega)} &\preceq \int_{B^n} (1 - |\zeta|^2)^{(n+1)(p-1)+mp} |f(\zeta)| \\ &\quad \times \int_{B^n} \frac{\omega(1 - |z|)(1 - |z|^2)^p d\nu(z)}{|1 - \langle z, \zeta \rangle|^{(m+n+2)p} (1 - |z|^2)^{n+1}} d\nu(\zeta) \\ &\preceq \int_{B^n} |f(\zeta)|^p \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|^2)^{n+1}} d\nu(\zeta) = \|f\|_{L_p(\omega)}. \end{aligned}$$

Now, we shall show that for any sequence $\{f_k\} \subset B_p(\omega)$ such that $\|f_k - f\|_{L_p(\omega)} \rightarrow 0$ as $k \rightarrow \infty$, the limit function $f \in L_p(\omega)$ is holomorphic in B^n . To this end, suppose that K is a compact set in B^n . Then by Lemma 3 there exists a constant $C(K, n, \omega, p)$ such that $\max_{z \in K} |f(z)| \leq C(K, n, \omega, p) \|f\|_{B_p(\omega)}$. Hence

$$\max_{z \in K} |f_k(z) - f_m(z)| \leq C(K, n, \omega, p) \|f_k - f_m\|_{B_p(\omega)}$$

for all k, m . Thus, $\{f_k\}$ uniformly converges to a holomorphic function $g(z)$ on all compact subsets of B^n . Since the compact sets are arbitrary, g is holomorphic on B^n . By Riesz' theorem, some subsequence of $\{f_k(z)\}$ pointwise converges to f for almost all $z \in B^n$. Hence $f = g$ for almost all $z \in B^n$, and the desired statement follows. \square

Corollary 3. $B_p(\omega)$ is a Banach space for $1 \leq p < \infty$, and a complete metric space for $0 < p < 1$.

Theorem 2. The following statements are true for any $0 < p < +\infty$:

1. If $f \in B_p(\omega)$ and $f_r(z) = f(rz)$, $0 < r < 1$, then $\|f - f_r\|_{B_p(\omega)} \rightarrow 0$ as $r \rightarrow 1 - 0$.
2. The set of polynomials is dense in $B_p(\omega)$, i.e. for any $f \in B_p(\omega)$ there is a sequence $\{P_n\}$ of polynomials such that $\|P_n - f\|_{B_p(\omega)} \rightarrow 0$ as $n \rightarrow \infty$.

PROOF: For completeness, we give a full proof, although it is based on a standard argument.

1. By the inequality $(a + b)^p \leq 2^p(a^p + b^p)$ ($a, b > 0$), for $\delta \in (0, 1)$ we get

$$\begin{aligned} \|f - f_r\|_{B_p(\omega)} &\leq \int_0^\delta \int_{S^n} |Df(z) - Df_r(z)|^p \frac{(1 - |z|^2)^p \omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\sigma(z) dr \\ &\quad + \int_\delta^1 \int_{S^n} |Df(z) - Df_r(z)|^p \frac{(1 - |z|^2)^p \omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\sigma(z) dr. \end{aligned}$$

The function $|Df|^p$ is subharmonic in B^n , and hence

$$\int_{S^n} |Df_r(z)|^p d\sigma(z) \leq \int_{S^n} |Df(z)|^p d\sigma(z).$$

Therefore,

$$\begin{aligned} \|f - f_r\|_{B_p(\omega)} &\leq \int_0^\delta \int_{S^n} |Df(z) - Df_r(z)|^p \frac{(1 - |z|^2)^p \omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\sigma(z) dr \\ &\quad + 2^p \int_\delta^1 \int_{S^n} |Df(z)|^p \frac{(1 - |z|^2)^p \omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\sigma(z) dr. \end{aligned}$$

The first integral in the right-hand side of this inequality vanishes as $r \rightarrow 1 - 0$, and the second one can be made arbitrarily small by choosing δ close enough to 1.

2. Let $f \in B_p(\omega)$ be an arbitrary function. Then

$$\lim_{r \rightarrow 1-0} \|f - f_r\|_{B_p(\omega)} = 0.$$

Further, the function f_r can be uniformly approximated by its Taylor polynomials in a neighborhood of \overline{B}^n . Therefore, the function f can be uniformly approximated in norm by a sequence of polynomials. \square

The following theorem gives a description of the space $B_p(\omega)$ in the terms of $L_p(\omega)$ ($0 < p < +\infty$).

Theorem 3. *Let $f \in H(B^n)$. Then for any $0 < p < +\infty$ the inclusion $f \in B_p(\omega)$ is true if and only if the function $g(z) = (1 - |z|^2)^\beta D^\beta f(z)$ is in $L_p(\omega)$ for some $\beta \geq 1$. Moreover, there are some constants C_1, C_2 such that*

$$(3) \quad C_1 \|g\|_{L_p(\omega)} \leq \|f\|_{B_p(\omega)} \leq C_1 \|g\|_{L_p(\omega)}.$$

PROOF: If $f \in B_p(\omega)$, then by (2) and Lemmas 8, 7

$$|D^\beta f(z)| \leq C(n, m) \int_{B^n} \frac{(1 - |\zeta|^2)^m |Df(\zeta)|}{|1 - \langle z, \zeta \rangle|^{m+n+\beta}} d\nu(\zeta).$$

Let $p > 1$. Then

$$\begin{aligned} &\int_{S^n} |D^\beta f(r\zeta)|^p d\sigma(\zeta) \\ &\leq C(m, n) \int_{S^n} \left| \int_0^1 (1 - \rho^2)^m \rho^{2n-1} \int_{S^n} \frac{|Df(\rho\zeta)| d\sigma(\zeta)}{|1 - r\rho\langle z, \zeta \rangle|^{m+n+\beta}} \right|^p d\sigma(z) d\rho \\ &\leq C(m, n) \int_0^1 \int_{S^n} \left| \int_{S^n} \frac{|Df(\rho\zeta)| d\sigma(\zeta)}{|1 - r\rho\langle z, \zeta \rangle|^{m+n+\beta}} \right|^p d\sigma(z) (1 - \rho^2)^{mp} \rho^{(2n-1)p} d\rho \end{aligned}$$

$$\begin{aligned} &\leq C(m, n) \int_0^1 \int_{S^n} \int_{S^n} \frac{|Df(\rho\zeta)|^p d\sigma(\zeta)}{|1 - r\rho\langle z, \zeta \rangle|^{m+n+\beta}} \\ &\quad \times \left(\int_{S^n} \frac{d\sigma(\zeta)}{|1 - r\rho\langle z, \zeta \rangle|^{m+n+\beta}} \right)^{p/q} d\sigma(z) (1 - \rho^2)^{mp} \rho^{(2n-1)p} d\rho. \end{aligned}$$

Further, using [15, Theorem 1.12] we get

$$\begin{aligned} &\int_{S^n} |D^\beta f(r\zeta)|^p d\sigma(\zeta) \\ &\leq C(m, n) \int_0^1 \int_{S^n} \int_{S^n} \frac{|Df(\rho\zeta)|^p d\sigma(\zeta)}{|1 - r\rho\langle z, \zeta \rangle|^{m+n+\beta}} d\sigma(z) \frac{(1 - \rho^2)^{mp} \rho^{(2n-1)p}}{(1 - r\rho)^{(m+\beta)p/q}} d\rho \\ &= C(m, n) \int_0^1 \int_{S^n} |Df(\rho\zeta)|^p d\sigma(\zeta) \int_{S^n} \frac{d\sigma(z)}{|1 - r\rho\langle z, \zeta \rangle|^{m+n+\beta}} \\ &\quad \times \frac{(1 - \rho^2)^{mp} \rho^{(2n-1)p}}{(1 - r\rho)^{(m+\beta)p/q}} d\rho \leq \int_0^1 \int_{S^n} |Df(\rho\zeta)|^p d\sigma(\zeta) \frac{(1 - \rho^2)^{mp} \rho^{(2n-1)p}}{(1 - r\rho)^{(m+\beta)p}} d\rho. \end{aligned}$$

Therefore,

$$\begin{aligned} &\int_0^1 \frac{(1 - r^2)^{\beta p} \omega(1 - r)}{(1 - r^2)^{n+1}} \int_{S^n} |D^\beta f(\rho\zeta)|^p d\sigma(\zeta) r^{2n-1} dr \\ &\leq \int_0^1 \int_0^1 \int_{S^n} |D^\beta f(\rho\zeta)|^p d\sigma(\zeta) \frac{(1 - \rho^2)^{mp} \rho^{(2n-1)p} r^{2n-1} \omega(1 - r) (1 - r^2)^{\beta p}}{(1 - \rho r)^{(m+\beta)p} (1 - r^2)^{n+1}} dr d\rho \\ &= \int_0^1 \int_{S^n} |D^\beta f(\rho\zeta)|^p (1 - \rho^2)^{mp} \rho^{(2n-1)p} \\ &\quad \times \int_0^1 \frac{r^{2n-1} \omega(1 - r) (1 - r^2)^{\beta p}}{(1 - \rho r)^{(m+\beta)p} (1 - r^2)^{n+1}} dr d\rho d\sigma(\zeta). \end{aligned}$$

Further, by [7, Lemma 1.6] we obtain

$$\int_0^1 \frac{r^{2n-1} \omega(1 - r) (1 - r^2)^{\beta p}}{(1 - \rho r)^{(m+\beta)p} (1 - r^2)^{n+1}} dr \preceq \frac{\omega(1 - \rho)}{(1 - \rho)^{mp+n}}.$$

Consequently,

$$\int_{B^n} |g(w)|^p \frac{\omega(1 - |w|)}{(1 - |w|^2)^{n+1}} d\nu(w) \preceq \|f\|_{B_p(\omega)}.$$

Conversely, if $f \in H(U^n)$ and $g \in L^p(\omega)$, then $f \in B_p(\omega)$. Indeed, using Lemma 7 we obtain

$$|Df(z)| \preceq \int_{B^n} \frac{(1 - |\zeta|^2)^m |D^\beta f(\zeta)|}{|1 - \langle z, \zeta \rangle|^{m-\beta+n+2}} d\nu(\zeta)$$

(recall that m is assumed to be sufficiently great). Using Hölder's inequality, we get

$$\begin{aligned} |Df(z)|^p &\preceq \int_{B^n} \frac{(1-|\zeta|^2)^{m+p\beta}}{|1-\langle z, \zeta \rangle|^{m+n+2-\beta}} |D^\beta f(\zeta)|^p d\nu(\zeta) \\ &\quad \times \left(\int_{B^n} \frac{(1-|\zeta|^2)^{m-\beta q} d\nu(\zeta)}{|1-\langle z, \zeta \rangle|^{m+n+2-\beta}} \right)^{p/q} \\ &\preceq (1-|z|^2)^{-(\beta q+1-\beta)p/q} \int_{B^n} \frac{(1-|\zeta|^2)^{m+p\beta}}{|1-\langle z, \zeta \rangle|^{m+n+2-\beta}} |D^\beta f(\zeta)|^p d\nu(\zeta), \end{aligned}$$

where [15, Theorem 1.12] is used for obtaining the last inequality. Consequently,

$$\begin{aligned} \|f\|_{B_p(\omega)} &\preceq \int_{B^n} |D^\beta f(\zeta)|^p (1-|\zeta|^2)^{\beta p+m} \\ &\quad \times \int_0^1 \omega(1-r)(1-r^2)^{p-n-1-(\beta q+1-\beta)p/q} \int_{S^n} \frac{d\nu(z)}{|1-r\langle z, \zeta \rangle|^{m+n+2-\beta}} r^{2n-1} d\nu(\zeta) \\ &\preceq \int_{B^n} |D^\beta f(\zeta)|^p (1-|\zeta|^2)^{\beta p+m} \int_0^1 \frac{\omega(1-r)(1-r^2)^{p-n-1-(\beta q+1-\beta)p/q} r^{2n-1}}{(1-r|\zeta|^2)^{m+2-\beta}} dr \\ &\preceq \int_{B^n} |D^\beta f(\zeta)|^p (1-|\zeta|^2)^{\beta p+m} \frac{\omega(1-|\zeta|)(1-|\zeta|^2)^{p-n-1-(\beta q+1-\beta)p/q} r^{2n-1}}{(1-|\zeta|^2)^{m+1-\beta}} d\nu(\zeta) \\ &= \int_{B^n} |D^\beta f(\zeta)|^p (1-|\zeta|^2)^{\beta p} \frac{\omega(1-|\zeta|)}{(1-|\zeta|^2)^{n+1}} d\nu(\zeta) = \|g\|_{L_p(\omega)}, \end{aligned}$$

where $g(z) = (1-|z|^2)^\beta D^\beta f(z)$. Summing up, we get the proof of (3).

The case $0 < p \leq 1$ requires a different proof. Let $f \in B_p(\omega)$. Then by Lemmas 8, 7

$$|D^\beta f(z)|^p \preceq \int_{B^n} \frac{(1-|\zeta|^2)^{mp+(n+1)p-n-1}}{|1-\langle z, \zeta \rangle|^{(m+n+\beta)p}} |Df(\zeta)|^p d\nu(\zeta).$$

Therefore,

$$\begin{aligned} \|g\|_{L_p(\omega)} &= \int_{B^n} (1-|z|^2)^{\beta p} |D^\beta f(z)|^p \frac{\omega(1-|z|)}{(1-|z|^2)^{n+1}} d\nu(z) \\ &\preceq \int_{B^n} (1-|\zeta|^2)^{mp+(n+1)p-n-1} |Df(\zeta)| \\ &\quad \times \int_{B^n} \frac{(1-|z|^2)^{\beta p-n-1} \omega(1-|z|)}{|1-\langle z, \zeta \rangle|^{(m+n+\beta)p}} d\nu(\zeta) d\nu(z) \\ &\preceq \int_{B^n} (1-|\zeta|^2)^p |Df(\zeta)|^p \frac{\omega(1-|\zeta|)}{(1-|\zeta|^2)^{n+1}} d\nu(\zeta) = \|f\|_{B_p(\omega)} < +\infty. \end{aligned}$$

Conversely, if $g \in L_p(\omega)$, then using Lemmas 8 and 7 one more time we get

$$|Df(z)|^p \preceq \int_{B^n} \frac{(1 - |\zeta|^2)^{mp+(n+1)p-n-1}}{|1 - \langle z, \zeta \rangle|^{(m+n+2-\beta)p}} |D^\beta f(\zeta)|^p d\nu(\zeta).$$

Therefore,

$$\begin{aligned} & \int_{B^n} (1 - |z|^2)^p |Df(z)|^p \frac{\omega(1 - |z|)}{(1 - |z|)^{n+1}} d\nu(z) \\ & \leq \int_{B^n} (1 - |\zeta|^2)^{mp+(n+1)p-n-1} |D^\beta f(\zeta)|^p \int_{B^n} \frac{(1 - |z|^2)^{p-n-1} \omega(1 - |z|)}{|1 - \langle z, \zeta \rangle|^{(m+n+2-\beta)p}} d\nu(\zeta) \\ & \preceq \int_{B^n} (1 - |\zeta|^2)^{\beta p} |D^\beta f(\zeta)|^p \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|)^{n+1}} d\nu(\zeta) = \|g\|_{L_p(\omega)}, \end{aligned}$$

where [7, Lemma 1.7] and [15, Theorem 1.12] were used for the last inequality. \square

3. Bounded and inverse operators on $B_p(\omega)$

Let us consider the linear operator

$$(4) \quad P_\alpha(f)(z) := \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha}{(1 - \langle z, \zeta \rangle)^{n+1+\alpha}} f(\zeta) d\nu(\zeta) \quad (\alpha > -1).$$

If $f \in L_p(\omega)$, then obviously $P_\alpha(f)$ is holomorphic on B^n . For finding the class of functions to which it belongs, we prove the following theorem stating that P_α is a bounded operator on $L_p(\omega)$.

Theorem 4. *If $\alpha > -\alpha_\omega - n - 1$ and $1 \leq p < \infty$, then $P_\alpha : L_p(\omega) \rightarrow B_p(\omega)$ boundedly.*

PROOF: We shall prove that if $f \in L_p(\omega)$ and $F = P_\alpha(f)$, then $F \in B_p(\omega)$. It is obvious that

$$|D^\beta F(z)| \preceq \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha |f(\zeta)|}{|1 - \langle z, \zeta \rangle|^{n+1+\alpha+\beta}} d\nu(\zeta),$$

where we assume that $\alpha + \beta + 1 > 0$ and $\beta > n + \beta_\omega$.

If $p > 1$, then by Hölder's inequality

$$\begin{aligned} & \int_{B^n} (1 - |z|^2)^{\beta p} |D^\beta F(z)|^p \frac{\omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\nu(z) \\ & \preceq \int_{B^n} (1 - |\zeta|^2)^\alpha |f(\zeta)|^p \int_{B^n} \frac{(1 - |z|^2)^{\beta p - \beta p/q - n - 1} \omega(1 - |z|)}{|1 - \langle z, \zeta \rangle|^{n+1+\alpha+\beta}} d\nu(\zeta) d\nu(z) \\ & = \int_{B^n} (1 - |\zeta|^2)^\alpha |f(\zeta)|^p \int_0^1 \int_{S^n} \frac{r^{2n-1} (1 - r^2)^{\beta - n - 1} \omega(1 - r) d\sigma(z)}{|1 - r\langle z, \zeta \rangle|^{n+1+\alpha+\beta}} dr d\nu(\zeta) \\ & \preceq \int_{B^n} (1 - |\zeta|^2)^\alpha |f(\zeta)|^p \int_0^1 \frac{r^{2n-1} (1 - r^2)^{\beta - n - 1} \omega(1 - r)}{(1 - r|\zeta|)^{\alpha+\beta+1}} dr d\nu(\zeta) \end{aligned}$$

$$\leq \int_{B^n} (1 - |\zeta|^2)^\alpha |f(\zeta)|^p \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|)^{n+\alpha+1}} d\nu(\zeta) = \|f\|_{L_p(\omega)},$$

where [15, Theorem 1.12] and [7, Lemma 1.7] were used for the last inequality.

If $p = 1$, then

$$\begin{aligned} & \int_{B^n} (1 - |z|^2)^\beta |D^\beta F(z)| \frac{\omega(1 - |z|)}{(1 - |z|^2)^{n+1}} d\nu(z) \\ & \leq \int_{B^n} (1 - |\zeta|^2)^\alpha |f(\zeta)| \int_{B^n} \frac{(1 - |z|^2)^\beta \omega(1 - |z|) d\nu(z) d\nu(\zeta)}{|1 - \langle z, \zeta \rangle|^{n+1+\beta+\alpha} (1 - |z|^2)^{n+1}} \leq \|f\|_{L_1(\omega)}. \end{aligned}$$

□

For the case when $1 \leq p < \infty$ and $\alpha > -1$, $\gamma > 0$, we define the inverse mapping $R_{\alpha,\gamma}$ of P_α by the formula

$$(5) \quad R_{\alpha,\gamma}(f)(z) := (1 - |z|^2)^\gamma \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha f(\zeta)}{(1 - \langle z, \zeta \rangle)^{n+1+\alpha+\gamma}} d\nu(\zeta), \quad z \in B^n.$$

One can prove that $P_\alpha R_{\alpha,\gamma}(f)(z) \equiv f(z)$ for all $f \in B_p(\omega)$ if $\alpha > 2n - 2$. To this end, observe that a change of integration order gives

$$\begin{aligned} P_\alpha R_{\alpha,\gamma}(f)(z) &= \int_{B^n} \frac{(1 - |t|^2)^\alpha f(t) d\nu(t)}{(1 - \langle \zeta, t \rangle)^{n+1+\alpha+\gamma}} d\nu(\zeta) \\ &= \int_{B^n} (1 - |t|^2)^\alpha f(t) \int_{B^n} \frac{(1 - |\zeta|^2)^{\alpha+\gamma} d\nu(\zeta)}{(1 - \langle t, \zeta \rangle)^{n+1+\alpha+\gamma} (1 - \langle \zeta, z \rangle)^{n+1+\alpha}} d\nu(t) \\ &= \int_{B^n} \frac{(1 - |t|^2)^\alpha f(t)}{(1 - \langle \zeta, t \rangle)^{n+1+\alpha}} d\nu(t) \equiv f(z). \end{aligned}$$

Further, we show that $R_{\alpha,\gamma}(f) \in L_p(\omega)$ for all $f \in L_p(\omega)$. Indeed, if $1 < p < \infty$, then by Hölder's inequality

$$\begin{aligned} |f(z)| &\leq \int_{B^n} \frac{(1 - |\zeta|^2)^m |Df(\zeta)|}{|1 - \langle z, \zeta \rangle|^{m+n}} d\nu(\zeta) \\ &\leq (1 - |z|^2)^{\frac{1-m}{q}} \left(\int_{B^n} \frac{(1 - |\zeta|^2)^{mp} |Df(\zeta)|^p}{|1 - \langle z, \zeta \rangle|^{m+n}} d\nu(\zeta) \right)^{1/p}. \end{aligned}$$

Consequently,

$$\begin{aligned} & \left(\int_{B^n} \frac{(1 - |\zeta|^2)^\alpha f(\zeta)}{(1 - \langle z, \zeta \rangle)^{n+1+\alpha+\gamma}} d\nu(\zeta) \right)^p \\ & \leq (1 - |z|^2)^{-\gamma p/q} \int_{B^n} \frac{(1 - |\zeta|^2)^\alpha |f(\zeta)|}{(1 - \langle z, \zeta \rangle)^{n+1+\alpha+\gamma}} d\nu(\zeta) \\ & \preceq \int_{B^n} \frac{(1 - |\zeta|^2)^{\alpha-(m-1)p/q}}{(1 - \langle z, \zeta \rangle)^{n+1+\alpha+\gamma}} \int_{B^n} \frac{(1 - |t|^2)^{mp} |Df(t)|^p}{|1 - \langle \zeta, t \rangle|^{m+n}} d\nu(t) d\nu(\zeta), \end{aligned}$$

and hence

$$\begin{aligned}
 & \int_{B^n} \frac{\omega(1-|z|)}{(1-|z|^2)^{n+1}} |R_\alpha(f)(z)|^p d\nu(z) \leq \int_{B^n} (1-|t|^2)^{mp} |Df(t)|^p \\
 & \quad \times \int_{B^n} \frac{(1-|\zeta|^2)^{\alpha-(m-1)p/q}}{|1-\langle \zeta, t \rangle|^{m+n}} \int_{B^n} \frac{(1-|z|)^{\gamma p - \gamma p/q - n - 1}}{|1-\langle z, \zeta \rangle|^{n+1+\alpha+\gamma}} d\nu(\zeta) d\nu(t) \\
 & \leq \int_{B^n} (1-|t|^2)^{mp} |Df(t)|^p \\
 & \quad \times \int_{B^n} \frac{(1-|\zeta|^2)^{\alpha-(m-1)p/q + p\gamma - \gamma p/q} \omega(1-|\zeta|)}{|1-\langle \zeta, t \rangle|^{m+n} (1-|\zeta|^2)^{n+1+\alpha+\gamma}} d\nu(\zeta) d\nu(t) \\
 & \leq \int_{B^n} |Df(t)|^p \frac{(1-|t|^2)^{mp+\alpha-(m-1)p/q} \omega(1-|t|)}{(1-|t|^2)^{m+n+\alpha}} d\nu(t) \\
 & = \int_{B^n} (1-|t|^2)^p |Df(t)|^p \frac{\omega(1-|t|)}{(1-|t|^2)^{n+1}} d\nu(t) = \|f\|_{B_p(\omega)}.
 \end{aligned}$$

If $p = 1$, then

$$\begin{aligned}
 & \int_{B^n} \frac{(1-|\zeta|^2)^\alpha |f(\zeta)|}{|1-\langle z, \zeta \rangle|^{n+1+\alpha+\gamma}} \\
 & \leq \int_{B^n} \frac{(1-|\zeta|^2)^\alpha}{|1-\langle z, \zeta \rangle|^{n+1+\alpha+\gamma}} \int_{B^n} \frac{(1-|t|^2)^m |Df(t)|}{|1-\langle t, \zeta \rangle|^{m+n}} d\nu(t) d\nu(\zeta).
 \end{aligned}$$

Hence

$$\begin{aligned}
 & \int_{B^n} |R_{\alpha,\gamma}(f)(z)| \frac{\omega(1-|z|)}{(1-|z|^2)^{n+1}} d\nu(z) \leq \int_{B^n} (1-|t|^2)^m |Df(t)| \\
 & \quad \times \int_{B^n} \frac{(1-|\zeta|^2)^\alpha}{|1-\langle t, \zeta \rangle|^{n+m}} \int_{B^n} \frac{\omega(1-|\zeta|)(1-|\zeta|)^\gamma d\nu(\zeta)}{|1-\langle t, \zeta \rangle|^{n+1+\alpha+\gamma} (1-|\zeta|^2)^{n+1}} d\nu(t) \\
 & \leq \int_{B^n} (1-|t|^2)^m |Df(t)| \int_{B^n} \frac{\omega(1-|\zeta|) d\nu(\zeta)}{|1-\langle t, \zeta \rangle|^{n+m} (1-|\zeta|^2)^{n+1}} d\nu(t) \\
 & \leq \int_{B^n} (1-|t|^2)^m |Df(t)| \frac{\omega(1-|t|) d\nu(t)}{(1-|t|)^{m-1+n+1}} \\
 & = \int_{B^n} |Df(t)| \frac{\omega(1-|t|) d\nu(t)}{(1-|t|)^n} = \|f\|_{B_p(\omega)}.
 \end{aligned}$$

Thus, we proved the following

Theorem 5. *If $1 \leq p < \infty$ and $\alpha + \gamma > -1$, then R_α is a bounded operator from $B_p(\omega)$ to $L_p(\omega)$.*

For the case $0 < p < 1$, we consider the harmonic subspace $b_p(\omega)$ of $L_p(\omega)$. Repeating the argument of [3, Lemma 2.3] one can prove the following lemma.

Lemma 9. *Let $0 < p < \infty$ and let g be harmonic on the ball $B_r^n = B_r^n(z_0)$. Then there is a constant C_p depending only on p and such that*

$$|u(z_0)|^p \leq \frac{C_p}{\pi r^{n+1}} \int_{B_r} |u(z)|^p d\nu(z).$$

As corollaries of the above lemma, one can prove also the following two statements.

Lemma 10. *Let $0 < p < \infty$ and let $g \in b_p(\omega)$. Then*

$$|g(z)| \leq \frac{\|g\|_{b_p(\omega)}}{\omega^{1/p}(1-|z|)}, \quad z \in B^n.$$

Lemma 11. *Let $0 < p < \infty$ and let $g \in b_p(\omega)$. Then*

$$\int_{B^n} |g(z)| \frac{\omega^{1/p}(1-|z|)}{(1-|z|^2)^{n+1}} d\nu(z) \leq \int_{B^n} |g(z)|^p \frac{\omega(1-|z|)}{(1-|z|)^{n+1}} d\nu(z).$$

The last lemma gives a possibility to consider the operator $P_\alpha(f)$ on $b_p(\omega)$ in the case when $0 < p < 1$. Assuming that $f \in b_p(\omega)$, we shall show that $P_\alpha(f) \in B_p(\omega)$. To this end, we use Lemma 11 and obtain

$$\begin{aligned} & \int_{B^n} |P_\alpha f(z)|^p \frac{\omega(1-|z|)}{(1-|z|)^{n+1}} d\nu(z) \\ & \leq \int_{B^n} |f(\zeta)|^p \frac{(1-|\zeta|^2)^{(\alpha+n+1)p}}{(1-|\zeta|^2)^{n+1}} \int_{B^n} \frac{\omega(1-|z|)(1-|z|^2)^{-n-1}}{|1-\langle z, \zeta \rangle|^{(n+1+\alpha)p}} d\nu(z) d\nu(\zeta) \\ & \preceq \int_{B^n} \frac{\omega(1-|\zeta|)(1-|\zeta|^2)^{\alpha p}}{(1-|\zeta|^2)^{n+1+\alpha p}} |f(\zeta)|^p d\nu(\zeta) = \|f\|_{b_p(\omega)}. \end{aligned}$$

Thus, we proved

Theorem 6. *If $0 < p < 1$, then P_α is a bounded operator $b_p(\omega) \rightarrow B_p(\omega)$.*

For $0 < p < 1$, the inverse operator R_α defined by (5) maps $B_p(\omega)$ to $L_p(\omega)$. Indeed, by Lemma 11

$$\begin{aligned} & \int_{B^n} (1-|z|^2)^{\gamma p} \frac{\omega(1-|z|)}{(1-|z|^2)^{n+1}} \int_{B^n} \frac{(1-|\zeta|^2)^{\alpha p+(n+1)p} |f(\zeta)|^p d\nu(\zeta) d\nu(z)}{|1-\langle z, \zeta \rangle|^{(n+1+\alpha+\gamma)p} (1-|\zeta|^2)^{n+1}} \\ & = \int_{B^n} \frac{(1-|\zeta|^2)^{\alpha p+(n+1)p}}{(1-|\zeta|^2)^{n+1}} |f(\zeta)|^p \int_{B^n} \frac{(1-|z|^2)^{\gamma p} \omega(1-|z|) d\nu(\zeta) d\nu(z)}{(1-|z|^2)^{n+1} |1-\langle z, \zeta \rangle|^{(n+1+\alpha+\gamma)p}} \\ & \preceq \int_{B^n} \frac{(1-|\zeta|^2)^{\alpha p+(n+1)p} (1-|\zeta|^2)^{\gamma p} \omega(1-|\zeta|)}{(1-|\zeta|^2)^{n+1} (1-|\zeta|^2)^{(\alpha+\gamma+n+1)p}} |f(\zeta)|^p d\nu(\zeta) = \|f\|_{L_p(\omega)}. \end{aligned}$$

Summing up, we come to

Theorem 7. *If $0 < p < 1$, then the operator R_α boundedly maps $b_p(\omega)$ to $B_p(\omega)$.*

4. Duals of $B_p(\omega)$ spaces

Theorem 8. *If $1 < p < \infty$ and $\alpha > -(n + \beta_\omega)/p$, then the dual of the space $B_p(\omega)$ under the pairing*

$$(6) \quad \langle f, g \rangle = \int_{B^n} Df(\zeta) \overline{Dg(\zeta)} (1 - |\zeta|^2)^\alpha d\nu(\zeta)$$

is isomorphic to $B_q(\tilde{\omega})$, where $\tilde{\omega}(t) = \omega^{-q/p}(t)t^{(\alpha+n-1)q}$, $1/p + 1/q = 1$.

PROOF: Let $\Phi \in (B_p(\omega))^*$. In virtue of Theorem 2, we can consider $B_p(\omega)$ as a subspace of $L_p(\omega)$. Then, by the Hahn-Banach theorem, we can assume that $\Phi \in L_p(\omega)^*$, and hence there is a function $G \in L_q(\omega)$ ($1/q + 1/p = 1$) such that

$$\Phi(F) = \int_{B^n} F(\zeta) \overline{G(\zeta)} \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|^2)^{n+1}} d\nu(\zeta),$$

and $\|\Phi\| = \|G\|_{L_q(\omega)}$. Taking $F(z) = (1 - |z|^2)Df(z)$, where $f \in B_p(\omega)$, we get

$$\Phi(F) = \int_{B^n} Df(\zeta) \overline{G(\zeta)} \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|^2)^n} d\nu(\zeta).$$

Using the fact that $\alpha > -(n + \beta_\omega)/p$ we can write (2) for Df and get

$$\Phi(F) = \int_{B^n} (1 - |w|^2)^\alpha Df(w) \int_{B^n} \frac{\omega(1 - |\zeta|) \overline{G(\zeta)} d\nu(\zeta)}{(1 - \langle \zeta, w \rangle)^{\alpha+n+1} (1 - |\zeta|^2)^n} d\nu(w).$$

Let $G_1(w)$ be the middle integral:

$$\overline{G_1(w)} = \int_{B^n} \frac{\omega(1 - |\zeta|) \overline{G(\zeta)} d\nu(\zeta)}{(1 - \langle \zeta, w \rangle)^{n+1+\alpha} (1 - |\zeta|^2)^n}.$$

It is clear that $G_1(w)$ is a holomorphic function and there exists a function $g(w)$ such that $\overline{Dg(w)} = \overline{G_1(w)}$ (for example $g(z) = \int_0^1 G_1(rz) dr$). Next we show that $g \in B_q(\omega)$. By Hölder's inequality we get

$$|G_1(w)|^q \leq \frac{\omega^{q/p}(1 - |w|)}{(1 - |w|^2)^{(\alpha+n)q/p}} \int_{B^n} \frac{\omega(1 - |\zeta|) |G(\zeta)|^q d\nu(\zeta)}{|1 - \langle \zeta, w \rangle|^{n+1+\alpha} (1 - |\zeta|^2)^n}.$$

Then

$$\begin{aligned} & \int_{B^n} (1 - |z|^2)^q |Dg(z)|^q \frac{\tilde{\omega}(1 - |z|)}{(1 - |z|^2)^{n+1}} d\nu(z) \\ & \leq \int_{B^n} |G(\zeta)|^q \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|^2)^n} \int_{B^n} \frac{\tilde{\omega}(1 - |z|) |\omega^{q/p}(1 - |z|) (1 - |z|^2)^{q-n-1} d\nu(z)}{|1 - \langle z, \zeta \rangle|^{n+1+\alpha} (1 - |z|^2)^{(\alpha+n)q/p}} d\nu(\zeta) \\ & \leq \int_{B^n} |G(\zeta)|^q \frac{\omega(1 - |\zeta|)}{(1 - |\zeta|^2)^n} \int_{B^n} \frac{(1 - |z|^2)^{\alpha-1} d\nu(z)}{|1 - \langle z, \zeta \rangle|^{n+1+\alpha}} d\nu(\zeta) \end{aligned}$$

$$\leq \int_{B^n} |G(\zeta)|^q \frac{\omega(1-|\zeta|)}{(1-|\zeta|^2)^{n+1}} d\nu(z) = \|G\|_{L_p(\omega)}.$$

Conversely, if $g \in B_q(\tilde{\omega})$, then by Hölder's inequality one can prove that the functional Φ of the form (6) is bounded on $B_p(\omega)$:

$$\begin{aligned} |\Phi(f)| &\leq \int_{B^n} |Df(\zeta)| |Dg(\zeta)| \frac{\omega^{1/p}(1-|\zeta|)(1-|\zeta|^2)^{2+\alpha+n+1}}{(1-|\zeta|^2)^{n+3}\omega^{1/p}(1-|\zeta|)} d\nu(\zeta) \\ &\leq \left(\int_{B^n} (1-|\zeta|^2)^p |Df(\zeta)|^p \frac{\omega(1-|\zeta|)}{(1-|\zeta|^2)^{n+1}} d\nu(\zeta) \right)^{1/p} \\ &\quad \times \left(\int_{B^n} (1-|\zeta|^2)^q |Dg(\zeta)|^q \frac{(1-|\zeta|^2)^{(n+1)q+\alpha q} d\nu(\zeta)}{\omega^{q/p}(1-|\zeta|)(1-|\zeta|^2)^{2q}} \right)^{1/q} = \|f\|_{B_p(\omega)} \|g\|_{B_q(\tilde{\omega})}. \end{aligned}$$

□

The case $0 < p \leq 1$ calls for a different statement connected with the defined below holomorphic Bloch space on the unit ball of C^n (for details see [4]).

Definition 3. Let $\omega \in S$. We say that a function $f \in H(B^n)$ belongs to the Bloch space B_ω if

$$M_f = \sup_{z \in B^n} \frac{(1-|z|^2)|Df(z)|}{\omega(1-|z|)} < \infty.$$

Theorem 9. If $0 < p \leq 1$, then the dual of the space $B_p(\omega)$ under the pairing

$$(7) \quad \langle f, g \rangle = \int_{B^n} Df(\zeta) \overline{Dg(\zeta)} (1-|\zeta|^2)^\alpha d\nu(\zeta) \quad (\alpha > n/p - \beta_\omega/p)$$

is isomorphic to the holomorphic weighted Bloch space $B_{\tilde{\omega}}$ with $\tilde{\omega}(t) = \omega^{1/p}(t)t^{-n-\alpha+1}$.

PROOF: Let $f \in B_1(\omega)$ and let $g \in B_{\tilde{\omega}}$ and Φ be the functional generated by g , i.e. $\Phi(f) = \langle f, g \rangle$. Then using Lemma 4 we obtain

$$\begin{aligned} |\Phi(f)| &\leq \sup_{z \in B^n} \frac{(1-|\zeta|)|Dg(\zeta)|}{\tilde{\omega}(1-|\zeta|)} \int_{B^n} |Df(\zeta)| \tilde{\omega}(1-|\zeta|)(1-|\zeta|^2)^{\alpha-1} d\nu(\zeta) \\ &\leq \|g\|_{B_{\tilde{\omega}}} \int_{B^n} |Df(\zeta)| \frac{\tilde{\omega}^{1/p}(1-|\zeta|)}{(1-|\zeta|^2)^n} d\nu(\zeta) \\ &\leq \|g\|_{B_{\tilde{\omega}}} \left(\int_{B^n} (1-|\zeta|^2)^p |Df(\zeta)|^p \frac{\tilde{\omega}(1-|\zeta|)}{(1-|\zeta|^2)^{n+1}} d\nu(\zeta) \right)^{1/p} \\ &= \|g\|_{B_{\tilde{\omega}}} \|f\|_{B_p(\omega)} \end{aligned}$$

Hence $\|\Phi\| \leq \|g\|_{B_{\tilde{\omega}}}$.

Conversely, if $\Phi \in (B_p(\omega))^*$, $f \in B_p(\omega)$, then $f \in B_1(\omega^*)$ by Corollary 1. Considering $B_1(\omega^*)$ as a subspace of $L_1(\omega^*)$ and using the Hahn-Banach theorem

we get $\Phi \in (L_1(\omega^*))^*$. Therefore, there is a function $G \in L_\infty(B^n)$ such that

$$\Phi(f) = \int_{B^n} F(\zeta) \overline{G(\zeta)} \frac{\omega^{1/p}(1-|\zeta|)}{(1-|\zeta|^2)^{n+1}} d\nu(\zeta)$$

and $\|\Phi\| = \|G\|_{L_\infty}$. Particularly, taking $F(\zeta) = (1-|\zeta|^2)Df(\zeta)$, $f \in B_p(\omega)$ we obtain

$$\Phi(f) = \int_{B^n} F(\zeta) \overline{G(\zeta)} \frac{\omega^{1/p}(1-|\zeta|)}{(1-|\zeta|^2)^{n+1}} d\nu(\zeta).$$

If $\alpha > n/p - \beta_\omega/p$, then $Df \in A^1(\alpha)$. Therefore, by (2)

$$\Phi(f) = \int_{B^n} (1-|w|^2)^\alpha Df(w) \int_{B^n} \frac{\overline{G(\zeta)} \omega^{1/p}(1-|\zeta|)}{(1-\langle w, \zeta \rangle)^{n+\alpha+1}(1-|\zeta|^2)^n} d\nu(\zeta) d\nu(w).$$

As in the case $p > 1$, we consider the inner integral separately, as a function $\overline{G_1(w)} = \overline{Dg(w)}$. Then we show that $g \in B_{\tilde{\omega}}$. To this end, observe that the following estimate is true:

$$|G_1(w)| \leq \|G\|_{L_\infty} \int_{B^n} \frac{\omega^{1/p}(1-|\zeta|) d\nu(\zeta)}{|1-\langle w, \zeta \rangle|^{n+1+\alpha}(1-|\zeta|^2)^n} \preceq \|G\|_{L_\infty} \frac{\omega^{1/p}(1-|w|)}{(1-|w|)^{n+\alpha}}.$$

Hence

$$\sup_{w \in B^n} \frac{(1-|w|^2)^{n+\alpha} |Dg(w)|}{\omega^{1/p}(1-|w|)} = \sup_{w \in B^n} \frac{(1-|w|^2) |Dg(w)|}{\tilde{\omega}(1-|w|)} < \infty,$$

where $\tilde{\omega}(t) = \omega^{1/p}(t)t^{-n-\alpha+1}$. □

REFERENCES

- [1] Arazy J., Fisher S., Peetre J., *Möbius invariant function spaces*, J. Reine Angew. Math. **363** (1985), 110–145.
- [2] Blasco O., *Multipliers on weighted Besov spaces of analytic functions*, Contemp. Math. **144** (1993), 23–33.
- [3] Djrbashian A., Shamoyan F., *Topics in the Theory of A_α^p spaces*, Teubner Texts in Math., 105, Teubner, Leipzig, 1988.
- [4] Harutyunyan A., Lusky W., *Holomorphic Bloch space on the unit ball in C^n* , Comment. Math. Univ. Carolin. **50** (2009), no. 4, 549–562.
- [5] Harutyunyan A., Lusky W., *ω -weighted holomorphic Besov spaces on polydiscs*, J. Funct. Spaces Appl., to appear.
- [6] Harutyunyan A., Lusky W., *Duals of holomorphic Besov spaces on the polydiscs and diagonal mappings*, J. Contemp. Math. Anal. **45** (2010), no. 3, 128–135.
- [7] Harutyunyan A.V., *Bloch spaces of holomorphic functions in the polydisc*, J. Funct. Spaces Appl. **5** (2007), no. 3, 213–230.
- [8] Karapetyants A.N., Kodzoeva F.D., *Analytic weighed Besov spaces on the unit disc*, Proc. A. Razmadze Math.Inst. **139** (2005), 125–127.
- [9] Nowak M., *Bloch and Möbius invariant Besov spaces on the unit ball of C^n* , Complex Variables Theory Appl. **44** (2001), 1–12.
- [10] Li S., Stevic S., *Some characterizations of the Besov space and the α -Bloch space*, J. Math. Anal. Appl. **346** (2008), 262–273.

- [11] Li S., Wulan H., *Besov space on the unit ball of \mathbb{C}^n* , Indian J. Math. **48** (2006), no. 2, 177–186.
- [12] Rudin W., *Function Theory in Unit Ball of C^n* , Springer, New York-Berlin, 1980.
- [13] Seneta E., *Functions of Regular Variation* (Russian), Nauka, Moscow, 1985.
- [14] Stroethoff K., *Besov type characterisations for the Bloch space*, Bull. Australian Math. Soc. **39** (1989), 405–420.
- [15] Zhu K., *Spaces of Holomorphic Functions in the Unit Ball*, Graduate Texts in Mathematics, 226, Springer, New York, 2005.

DEPARTMENT OF APPLIED MATHEMATICS, YEREVAN STATE UNIVERSITY,
1 ALEX MANOOKIAN STR., 375025 YEREVAN, ARMENIA

E-mail: anahit@ysu.am

INSTITUTE OF MATHEMATICS, UNIVERSITY OF PADERBORN, 100 WARBURGER STR.,
33098 PADERBORN, GERMANY

E-mail: lusky@math.uni-paderborn.de

(Received April 23, 2010)