

Kristína Rostás

Minimal and maximal solutions of fourth order iterated differential equations with singular nonlinearity

Archivum Mathematicum, Vol. 47 (2011), No. 1, 23--33

Persistent URL: <http://dml.cz/dmlcz/141507>

Terms of use:

© Masaryk University, 2011

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**MINIMAL AND MAXIMAL SOLUTIONS OF FOURTH ORDER
ITERATED DIFFERENTIAL EQUATIONS
WITH SINGULAR NONLINEARITY**

KRISTÍNA ROSTÁS

ABSTRACT. In this paper we are concerned with sufficient conditions for the existence of minimal and maximal solutions of differential equations of the form

$$L_4y + f(t, y) = 0,$$

where L_4y is the iterated linear differential operator of order 4 and $f: [a, \infty) \times (0, \infty) \rightarrow (0, \infty)$ is a continuous function.

1. INTRODUCTION

The purpose of this paper is to study the existence of positive solutions with specific asymptotic behavior for differential equations of the form

$$L_4y + f(t, y) = 0,$$

where L_4y is the iterated linear differential operator of fourth order defined below and $f: [a, \infty) \times (0, \infty) \rightarrow (0, \infty)$ is a continuous function, nonincreasing in the second variable.

A prototype of such equations is the equation with singular nonlinearity $f(t, y) = Q(t)y^{-\lambda}$, where $\lambda > 0$ and $Q: [a, \infty) \rightarrow (0, \infty)$ is continuous. Such equations of the second order were studied in [3], [4].

Differential equations with iterated linear differential operator were studied, for instance, in [5].

2. ITERATED DIFFERENTIAL EQUATIONS OF THE FOURTH ORDER

If u and v are linearly independent solutions of

$$(A_2) \quad y'' + P(t)y = 0,$$

where $P \in C^2[a, \infty)$, then $u, v \in C^4[a, \infty)$ and the linearly independent functions

$$y_1(t) = u^3(t), \quad y_2(t) = u^2(t)v(t), \quad y_3(t) = u(t)v^2(t), \quad y_4(t) = v^3(t)$$

2010 *Mathematics Subject Classification*: primary 34C10.

Key words and phrases: iterated differential equations, maximal and minimal solutions.

Supported by the VEGA Grant Agency of the Slovak Republic No. 1/0481/08

Received April 15, 2008, revised August 2010. Editor O. Došlý.

satisfy the fourth order linear differential equation

$$(A_4) \quad y^{IV} + 10P(t)y'' + 10P'(t)y' + [3P''(t) + 9P^2(t)]y = 0,$$

(see [1]). Differential equation (A₄) is called *iterated* linear differential equation of the fourth order.

We may suppose without loss of generality that

$$W[u, v](t) \equiv 1 \quad \text{for } t \geq a,$$

where $W[u, v](t)$ denotes Wronskian of functions u and v . An elementary calculation shows that Wronskian of functions

$$y_1(t) = u^3(t), \quad y_2(t) = u^2(t)v(t), \quad y_3(t) = u(t)v^2(t), \quad y_4(t) = v^3(t)$$

satisfies

$$W(u^3, u^2v, uv^2, v^3)(t) \equiv 12 \quad \text{for } t \geq a.$$

We suppose that the equation (A₂) is nonoscillatory and the $u(t)$ (resp. $v(t)$) denote a principal (resp. nonprincipal) solution of (A₂) such that

$$\lim_{t \rightarrow \infty} \frac{u(t)}{v(t)} = 0$$

and

$$\int^{\infty} \frac{dt}{u^2(t)} = \infty \quad (\text{resp. } \int^{\infty} \frac{dt}{v^2(t)} < \infty).$$

We may assume that both $u(t)$ and $v(t)$ are eventually positive. Second, nonprincipal $v(t)$ of (A₂) is given by

$$v(t) = u(t) \int_{t_0}^t \frac{ds}{u^2(s)}, \quad t \geq t_0.$$

In this paper we are concerned with the behavior of solutions of differential equations of the form

$$L_4y + f(t, y) = 0,$$

where L_4y is the iterated linear differential operator of order 4 and $f: [a, \infty) \times (0, \infty) \rightarrow (0, \infty)$ is a continuous function.

From Pólya's factorization theory it follows that the operator L_4y can be written in the form

$$L_4y = a_4(t)(a_3(t)(a_2(t)(a_1(t)(a_0(t)y)')')')',$$

where

$$a_0(t) = \frac{1}{u^3(t)}, \quad a_1(t) = u^2(t), \quad a_2(t) = \frac{u^2(t)}{2}, \quad a_3(t) = \frac{u^2(t)}{3}, \quad a_4(t) = \frac{6}{u^3(t)},$$

see [6].

3. CLASSIFICATION OF POSITIVE SOLUTIONS

Consider the fourth order differential equation

$$(A) \quad L_4y(t) + f(t, y(t)) = 0$$

where L_4y is the iterated linear differential operator of order 4 and $f: [a, \infty) \times (0, \infty) \rightarrow (0, \infty)$ is continuous, and nonincreasing in the second variable.

We assume that the equation (A₂) is nonoscillatory and put

$$L_0y(t) = \frac{y(t)}{u^3(t)},$$

$$L_iy(t) = \frac{u^2(t)}{i} \frac{d}{dt} (L_{i-1}y(t)), \quad 1 \leq i \leq 3,$$

and

$$L_4y = \frac{6}{u^3(t)} \left(\frac{u^2(t)}{3} \left(\frac{u^2(t)}{2} \left(u^2(t) \left(\frac{1}{u^3(t)} y \right)' \right)' \right)' \right)'.$$

The domain $\mathcal{D}(L_4)$ of the operator L_4 is defined to be the set of all continuous functions $y: [T_y, \infty) \rightarrow (0, \infty)$, $T_y \geq a$ such that $L_iy(t)$ for $0 \leq i \leq 3$ exist and are continuously differentiable on $[T_y, \infty)$.

Those functions which vanish in a neighborhood of infinity will be excluded from our consideration.

Our purpose here is to make a detailed analysis of the structure of the set of all possible positive solutions of the equation (A). We use the following lemma which is the special case of generalized Kiguradze's lemma (see [2]).

Lemma 1. *If $y(t)$ is a positive solution of (A), then either*

$$(1) \quad L_0y(t) > 0, \quad L_1y(t) > 0, \quad L_2y(t) < 0, \quad L_3y(t) > 0, \quad L_4y(t) < 0,$$

or

$$(2) \quad L_0y(t) > 0, \quad L_1y(t) > 0, \quad L_2y(t) > 0, \quad L_3y(t) > 0, \quad L_4y(t) < 0,$$

for all sufficiently large t .

Solutions satisfying (1) and (2) are called solutions of *Kiguradze's degree 1* and 3, respectively. If we denote by P the set of all positive solution of (A) and by P_l the set of all solution of degree l , then we have:

$$P = P_1 \cup P_3.$$

Consider P_l for $l = \{1, 3\}$. For any $y \in P_l$ the limits

$$\lim_{t \rightarrow \infty} L_ly(t) = c_l \quad (\text{finite}),$$

$$\lim_{t \rightarrow \infty} L_{l-1}y(t) = c_{l-1} \quad (\text{finite or infinite but not zero})$$

both exist.

Solution $y \in P_l$ is called a *maximal* in P_l , if c_l is nonzero and a *minimal* in P_l , if c_{l-1} is finite. The set of all maximal solutions in P_l denote $P_l[\text{max}]$ and the set of all minimal solutions in P_l denote $P_l[\text{min}]$.

If $c_l = 0$ for a solutions $y \in P_l$ for $l \in \{1, 3\}$, then y is called *intermediate in P_l* . The set of all intermediate solutions in P_l denote $P_l[\text{int}]$. Then

$$P = P_1[\text{min}] \cup P_1[\text{int}] \cup P_1[\text{max}] \cup P_3[\text{min}] \cup P_3[\text{int}] \cup P_3[\text{max}].$$

Our objective is to give sufficient conditions for the existence of maximal and minimal solutions in P_i for $i = 1, 3$.

Crucial role will be played by integral representations for those fourth types of solutions of (A) as derived below.

First we define: $I_0 = 1$ and

$$I_i(t, s; u) = \int_s^t \frac{1}{u^2(r)} I_{i-1}(r, s; u) dr, \quad 1 \leq i \leq 3.$$

If the second, linearly independent solution $v(t)$ of (A₂) is given by

$$v(t) = u(t) \int_{t_0}^t \frac{ds}{u^2(s)} \quad \text{for } t \geq t_0, \quad \text{then the set of positive functions}$$

$$x_0(t) = u^3(t),$$

$$x_1(t) = u^3(t) \int_{t_0}^t \frac{1}{u^2(s)} ds = u^2(t) v(t),$$

$$x_2(t) = u^3(t) \int_{t_0}^t \frac{1}{u^2(s)} \int_{t_0}^s \frac{2}{u^2(r)} dr ds = u(t) v^2(t),$$

$$x_3(t) = u^3(t) \int_{t_0}^t \frac{1}{u^2(s)} \int_{t_0}^s \frac{2}{u^2(r)} \int_{t_0}^r \frac{3}{u^2(\xi)} d\xi dr ds = v^3(t)$$

defined on $[t_0, \infty)$ form fundamental set of positive solutions for $L_4x = 0$ (i.e. (A₄), which are asymptotically ordered in the sense that

$$\lim_{t \rightarrow \infty} \frac{x_i(t)}{x_j(t)} = 0$$

for $0 \leq i < j \leq 3$, see [2]. It is useful to note that

$$I_i(t, a; u) = \frac{1}{i!} \left(\frac{v(t)}{u(t)} \right)^i \quad \text{for } i = 1, 2, 3.$$

The solutions from the classes $P_3[\text{max}]$, $P_3[\text{min}]$, $P_1[\text{max}]$ and $P_1[\text{min}]$ satisfy the properties

$$\lim_{t \rightarrow \infty} \frac{y(t)}{v^3(t)} = \lambda_3, \quad \lim_{t \rightarrow \infty} \frac{y(t)}{u(t)v^2(t)} = \lambda_2, \quad \lim_{t \rightarrow \infty} \frac{y(t)}{u^2(t)v(t)} = \lambda_1$$

$$\text{and } \lim_{t \rightarrow \infty} \frac{y(t)}{u^3(t)} = \lambda_0, \quad \text{respectively, where } 0 < \lambda_i < \infty, \quad i = 1, 2, 3.$$

4. INTEGRAL REPRESENTATIONS FOR SOLUTIONS

Now we can derive integral representations for types $P_3[\max]$, $P_3[\min]$, $P_1[\max]$ and $P_1[\min]$.

Let y be a solution of (A) such that $y(t) > 0$ for $t \geq T \geq a$. Integrating (A) from t to ∞ gives

$$(3) \quad L_3 y(t) = c_3 + \int_t^\infty \frac{u^3(s)}{6} f(s, y(s)) ds, \quad t \geq T,$$

where $c_3 = \lim_{t \rightarrow \infty} L_3 y(t) \geq 0$.

If $y \in P_3[\max]$, then we integrate (3) three times over $[T, t]$ to obtain

$$\begin{aligned} y(t) &= k_0 u^3(t) + k_1 u^3(t) \int_T^t \frac{1}{u^2(s)} ds + 2k_2 u^3(t) \int_T^t \frac{1}{u^2(s_1)} \int_T^{s_1} \frac{1}{u^2(s_2)} ds_2 ds_1 \\ &\quad + 6c_3 u^3(t) \int_T^t \frac{1}{u^2(s_1)} \int_T^{s_1} \frac{1}{u^2(s_2)} \int_T^{s_2} \frac{1}{u^2(s_3)} ds_3 ds_2 ds_1 \\ &\quad + u^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} \int_s^\infty u^3(r) f(r, y(r)) dr ds, \end{aligned}$$

for $t \geq T$, where $k_i = L_i y(T)$ for $i = 0, 1, 2$ and we used Fubini theorem.

If y is a solution of type $P_3[\min]$, then integrating (3) with $c_3 = 0$ from t to ∞ and then integrating the resulting equation twice from T to t , we have

$$\begin{aligned} y(t) &= k_0 u^3(t) + k_1 u^3(t) \int_T^t \frac{1}{u^2(s_1)} ds_1 + 2c_2 u^3(t) \int_T^t \frac{1}{u^2(s_1)} \int_T^{s_1} \frac{1}{u^2(s_2)} ds_2 ds_1 \\ &\quad - u^3(t) \int_T^t I_1(t, s; u) \frac{1}{u^2(s)} \int_s^\infty I_1(r, s; u) u^3(r) f(r, y(r)) dr ds, \end{aligned}$$

for $t \geq T$, where $c_2 = \lim_{t \rightarrow \infty} L_2 y(t)$.

An integral representation for a solution y of type $P_1[\max]$ is derived by integrating (3) with $c_3 = c_2 = 0$ twice from t to ∞ and once on $[T, t]$

$$\begin{aligned} y(t) &= k_0 u^3(t) + c_1 u^3(t) \int_T^t \frac{1}{u^2(s_1)} ds_1 \\ &\quad + u^3(t) \int_T^t \frac{1}{u^2(s)} \int_s^\infty I_2(r, s; u) u^3(r) f(r, y(r)) dr ds, \end{aligned}$$

for $t \geq T$, where $c_1 = \lim_{t \rightarrow \infty} L_1 y(t)$ and we used Fubini theorem.

If $y \in P_1[\min]$, then integrations of (3) with $c_3 = c_2 = c_1 = 0$ three times on (t, ∞) yield

$$y(t) = c_0 u^3(t) - u^3(t) \int_t^\infty I_3(s, t; u) u^3(s) f(s, y(s)) ds,$$

for $t \geq T$, where $c_0 = \lim_{t \rightarrow \infty} L_0 y(t)$.

5. EXISTENCE THEOREMS

We are now prepared to discuss the existence of maximal and minimal solutions of equation (A) of type P_1 and P_3 .

Theorem 1. *The equation (A) has a positive solution of types $P_3[\max]$ if*

$$(4) \quad \int^{\infty} u^3(t) f(t, cv^3(t)) dt < \infty,$$

for some $c > 0$.

Proof. We assume that (4) holds. Then there is $T \geq a$ such that

$$\int_T^{\infty} u^3(t) f(t, cv^3(t)) dt < c.$$

Let C denote locally convex space of all continuous functions $y: [T, \infty) \rightarrow R$ with the topology of uniform convergence on compact subintervals of $[T, \infty)$.

Define the subset Y_3 of $C[T, \infty)$ and mapping $\Phi_3: Y_3 \rightarrow C[T, \infty)$ by

$$Y_3 = \{y \in C[T, \infty) : cv^3(t) \leq y(t) \leq 2cv^3(t), t \geq T\}$$

and

$$\Phi_3 y(t) = cv^3(t) + u^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} \int_s^{\infty} u^3(r) f(r, y(r)) dr ds.$$

We will show that (i): Φ_3 maps Y_3 into Y_3 , (ii): Φ_3 is continuous on Y_3 , (iii): $\Phi_3(Y_3)$ is relatively compact.

(i) Since

$$\begin{aligned} 0 &\leq u^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} \int_s^{\infty} u^3(r) f(r, y(r)) dr ds \\ &\leq u^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} \int_s^{\infty} u^3(r) f(r, cv^3(r)) dr ds, \end{aligned}$$

then

$$\Phi_3 y(t) \geq cv^3(t)$$

and

$$\begin{aligned} \Phi_3 y(t) &\leq cv^3(t) + u^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} \int_s^{\infty} u^3(r) f(r, cv^3(r)) dr ds \\ &\leq cv^3(t) + cu^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} ds \\ &\leq cv^3(t) + cv^3(t) = 2cv^3(t). \end{aligned}$$

And so $\Phi_3 y \in Y_3$.

(ii) Suppose that $\{y_n\} \subset Y_3$ and $y \in Y_3$, and that $\lim_{n \rightarrow \infty} y_n = y$ in the topology of $C[T, \infty)$. We have

$$\begin{aligned} |\Phi_3 y_n(t) - \Phi_3 y(t)| &\leq u^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} \\ &\quad \times \int_s^\infty u^3(r) |f(r, y_n(r)) - f(r, y(r))| dr ds \\ &\leq 6v^3(t) \int_T^\infty u^3(s) |f(s, y_n(s)) - f(s, y(s))| ds \end{aligned}$$

Because

$$|f(s, y_n(s)) - f(s, y(s))| \leq 2f(s, cv^3(s))$$

and $\lim_{t \rightarrow \infty} |f(s, y_n(s)) - f(s, y(s))| = 0$ for $s \geq T$, then applying the Lebesgue convergence theorem, we have $|\Phi_3 y_n(t) - \Phi_3 y(t)| \rightarrow 0$ for $n \rightarrow \infty$ on every compact subinterval of $[T, \infty)$, which implies that $\Phi_3 y$ is continuous on Y_3 .

(iii) If $y \in Y_3$, then we have for $t \in (T, \infty)$

$$\left| \frac{d}{dt} \left(\frac{1}{u^3(t)} \Phi_3 y(t) \right) \right| \leq 2c \frac{1}{u^2(t)} I_2(t, a; u).$$

This shows that the function $\frac{d}{dt} \left(\frac{1}{u^3(t)} \Phi_3 y(t) \right)$ is uniformly bounded on any compact subinterval of $[T, \infty)$, and so function $\frac{1}{u^3(t)} \Phi_3 y(t)$ is equicontinuous on (T, ∞) .

Now for $t_1, t_2 \in [T, \infty)$ we see that

$$\begin{aligned} |\Phi_3 y(t_2) - \Phi_3 y(t_1)| &\leq |u^3(t_2) - u^3(t_1)| \left| \frac{1}{u^3(t_2)} \Phi_3 y(t_2) \right| \\ &\quad + u^3(t_1) \left| \frac{1}{u^3(t_2)} \Phi_3 y(t_2) - \frac{1}{u^3(t_1)} \Phi_3 y(t_1) \right| \\ &\leq 2c \frac{1}{u^3(t_2)} v^3(t_2) |u^3(t_2) - u^3(t_1)| \\ &\quad + u^3(t_1) \left| \frac{1}{u^3(t_2)} \Phi_3 y(t_2) - \frac{1}{u^3(t_1)} \Phi_3 y(t_1) \right|, \end{aligned}$$

and hence $\Phi_3(Y_3)$ is equicontinuous at every point of $[T, \infty)$. Since $\Phi_3(Y_3)$ is clearly uniformly bounded on $[T, \infty)$, it follow from Ascoli-Arzelà theorem that $\Phi_3(Y_3)$ is relatively compact.

Therefore, by the Schauder-Tychonoff fixed point theorem, there exists a fixed element $y \in Y_3$ of Φ_3 , i.e. $\Phi_3 y = y$, which satisfies the integral equation

$$y(t) = cv^3(t) + u^3(t) \int_T^t I_2(t, s; u) \frac{1}{u^2(s)} \int_s^\infty u^3(r) f(r, y(r)) dr ds.$$

A simple computation shows that this fixed point is a solution of (A) of type $P_3[\max]$. The proof of Theorem 1 is complete. \square

Theorem 2. *The equation (A) has a positive solution of type $P_3[\min]$ if*

$$(5) \quad \int_a^\infty u^2(t)v(t) f(t, cu(t)v^2(t)) dt < \infty,$$

for some $c > 0$.

Proof. Suppose that (5) holds. Choose $T \geq a$ so that

$$(6) \quad \int_T^\infty u^2(t)v(t) f(t, cu(t)v^2(t)) dt < c.$$

Consider the set Y_2 functions $y \in C[T, \infty)$ and mapping $\Psi_3: Y_3 \rightarrow C[T, \infty)$ defined by

$$Y_2 = \{y \in C[T, \infty) : cu(t)v^2(t) \leq y(t) \leq 2cu(t)v^2(t), t \geq T\}$$

and

$$\begin{aligned} \Psi_3 y(t) &= 2cu(t)v^2(t) - u^3(t)e \int_T^t I_1(t, s; u) \frac{1}{u^2(s)} \\ &\quad \times \int_s^\infty I_1(r, s; u) u^3(r) f(r, y(r)) dr ds. \end{aligned}$$

That $\Psi_3(Y_2) \subset Y_2$ is an immediate consequence of (6). Since the continuity of Ψ_3 and the relative compactness of $\Psi_3(Y_2)$ can be proved as in the proof of Theorem 1, there exists an element $y \in Y_2$ such that $\Psi_3 y = y$, which satisfies

$$\begin{aligned} y(t) &= 2cu(t)v^2(t) - u^3(t) \int_T^t I_1(t, s; u) \frac{1}{u^2(s)} \\ &\quad \times \int_s^\infty I_1(r, s; u) u^3(r) f(r, y(r)) dr ds \end{aligned}$$

for $t \geq T$. It is easy to verify that this fixed point is a solution of degree 3 of (A) such that $\lim_{t \rightarrow \infty} L_2 y(t) = c_2$ exists and is finite and nonzero. This completes the proof. \square

Theorem 3. *The equation (A) has a positive solution of type $P_1[\max]$ if*

$$(7) \quad \int_a^\infty u(t)v^2(t) f(t, cu^2(t)v(t)) dt < \infty,$$

for some $c > 0$.

Proof. Suppose that (7) holds. Take $T \geq a$ so large that

$$\int_T^\infty u(t)v^2(t) f(t, cu^2(t)v(t)) dt < c.$$

Consider a closed convex subset Y_1 of $C[T, \infty)$ defined by a

$$Y_1 = \{y \in C[T, \infty) : cu^2(t)v(t) \leq y(t) \leq 2cu^2(t)v(t), t \geq T\}.$$

Define the operator $\Phi_1: Y_1 \rightarrow C[T, \infty)$ by the following formula

$$\Phi_1 y(t) = cu^2(t)v(t) + u^3(t) \int_T^t \frac{1}{u^2(s)} \int_s^\infty I_2(r, s; u) u^3(r) f(r, y(r)) dr ds.$$

Again we can show that (i) $\Phi_1(Y_1) \subset Y_1$, (ii) Φ_1 is a continuous operator and (iii) $\Phi_1(Y_1)$ is relatively compact.

Therefore, Φ_1 has a fixed point $y \in Y_1$, which gives rise to a type $P_1[\max]$ solution of (A) since it satisfies

$$y(t) = cu^2(t)v(t) + u^3(t) \int_T^t \frac{1}{u^2(s)} \int_s^\infty I_2(r, s; u) u^3(r) f(r, y(r)) dr ds$$

for $t \geq T$. Note that $\lim_{t \rightarrow \infty} L_1 y(t) = c$. The proof is thus complete. \square

Theorem 4. *The equation (A) has positive solution of type $P_1[\min]$ if*

$$(8) \quad \int_a^\infty v^3(t) f(t, cu^3(t)) dt < \infty,$$

for some $c > 0$.

Proof. Suppose now that (8) holds. There exists a constant $T \geq a$ such that

$$\int_T^\infty v^3(t) f(t, cu^3(t)) dt < c.$$

Define the mapping Ψ_1 by

$$\Psi_1 y(t) = 2cu^3(t) - u^3(t) \int_t^\infty I_3(s, t; u) u^3(s) f(s, y(s)) ds.$$

Then, it can be verified without difficulty that Ψ_1 has a fixed element y in the set

$$Y_0 = \{y \in C[T, \infty) : cu^3(t) \leq y(t) \leq 2cu^3(t), t \geq T\}.$$

This fixed point gives rise to a required positive solution of (A), since it satisfies

$$y(t) = 2cu^3(t) - u^3(t) \int_t^\infty I_3(s, t; u) u^3(s) f(s, y(s)) ds.$$

Note that $\lim_{t \rightarrow \infty} L_0 y(t) = 2c$. This completes the proof. \square

6. SPECIAL CASE AND EXAMPLE

We consider equation (A) with special function $f(t, y) = Q(t)y^{-\lambda}$

$$(B) \quad L_4 y(t) + Q(t)y^{-\lambda} = 0,$$

where $\lambda > 0$ and $Q: [a, \infty) \rightarrow (0, \infty)$ is continuous. The objective of this section is

to use above theorems to establish sufficient conditions for equation (B) to have solutions $x_i(t)$, $i = 1, 2, 3, 4$ defined in some neighborhood of infinity with the same asymptotic behavior as

$$x_i(t) = u^{4-i}(t)v^{i-1}(t), \quad 1 \leq i \leq 4,$$

respectively, as $t \rightarrow \infty$. We write them as corollaries, where the symbol \sim is used to denote the asymptotic equivalence

$$f(t) \sim g(t) \quad \text{as } t \rightarrow \infty \quad \Leftrightarrow \quad \lim_{t \rightarrow \infty} \frac{f(t)}{g(t)} = 1.$$

Corollary 1. *A sufficient condition for (B) to have a positive solution $y_4(t)$ which satisfies*

$$y_4(t) \sim mv^3(t)$$

for some $m > 0$ is that

$$\int_a^\infty u^3(t) v^{-3\lambda}(t) Q(t) dt < \infty.$$

Corollary 2. *A sufficient condition for (B) to have a positive solution $y_3(t)$ which satisfies*

$$y_3(t) \sim mu(t)v^2(t)$$

for some $m > 0$ is that

$$\int_a^\infty u^{2-\lambda}(t) v^{1-2\lambda}(t) Q(t) dt < \infty.$$

Corollary 3. *A sufficient condition for (B) to have a positive solution $y_2(t)$ which satisfies*

$$y_2(t) \sim mu^2(t)v(t)$$

for some $m > 0$ is that

$$\int_a^\infty u^{1-2\lambda}(t) v^{2-\lambda}(t) Q(t) dt < \infty.$$

Corollary 4. *A sufficient condition for (B) to have a positive solution $y_1(t)$ which satisfies*

$$y_1(t) \sim 2mu^3(t)$$

for some $m > 0$ is that

$$\int_a^\infty u^{-3\lambda}(t) v^3(t) Q(t) dt < \infty.$$

We present here an example which illustrates theorems proved above and the corollaries.

Example. Consider the nonoscillatory linear differential equation of the second order

$$x'' + \frac{1}{4t^2}x = 0, \quad t \geq 1.$$

We know, that this equation has principal solution

$$u(t) = t^{\frac{1}{2}}$$

and nonprincipal solution

$$v(t) = t^{\frac{1}{2}} \ln t$$

and that the iterated equation

$$x^{IV} + \frac{5}{2t^2}x'' - \frac{5}{t^3}x' + \frac{81}{16t^4}x = 0,$$

has independent solutions in the form

$$x_1(t) = t^{\frac{3}{2}}, \quad x_2(t) = t^{\frac{3}{2}} \ln t, \quad x_3(t) = t^{\frac{3}{2}} \ln^2 t, \quad x_4(t) = t^{\frac{3}{2}} \ln^3 t.$$

Then equation

$$y^{IV} + \frac{5}{2t^2}y'' - \frac{5}{t^3}y' + \frac{81}{16t^4}y + Q(t)y^{-\lambda} = 0, \quad t \geq 1,$$

where $\lambda > 0$ and $Q: [1, \infty) \rightarrow (0, \infty)$ is continuous, has positive regular solution

- a) $y_1(t)$ satisfying $y_1(t) \sim mt^{\frac{3}{2}}$ if $\int_a^\infty t^{-\frac{3}{2}(\lambda-1)}(\ln t)^3 Q(t) dt < \infty$,
- b) $y_2(t)$ satisfying $y_2(t) \sim mt^{\frac{3}{2}} \ln t$ if $\int_a^\infty t^{-\frac{3}{2}(\lambda-1)}(\ln t)^{2-\lambda} Q(t) dt < \infty$,
- c) $y_3(t)$ satisfying $y_3(t) \sim mt^{\frac{3}{2}} \ln^2 t$ if $\int_a^\infty t^{-\frac{3}{2}(\lambda-1)}(\ln t)^{1-2\lambda} Q(t) dt < \infty$,
- d) $y_4(t)$ satisfying $y_4(t) \sim mt^{\frac{3}{2}} \ln^3 t$ if $\int_a^\infty t^{-\frac{3}{2}(\lambda-1)}(\ln t)^{-3\lambda} Q(t) dt < \infty$

for some $m > 0$.

Acknowledgement. The author is grateful to the referee for his useful suggestions.

REFERENCES

- [1] Barret, J. H., *Oscillation theory of ordinary linear differential equations*, Adv. Math. **3** (1969), 415–509.
- [2] Fink, A. M., Kusano, T., *Nonoscillation theorems for differential equations with general deviating arguments*, Lecture Notes in Math. **1032** (1983), 224–239.
- [3] Kusano, T., Swanson, C. A., *Asymptotic properties of semilinear elliptic equations*, Funkcial. Ekvac. **26** (1983), 115–129.
- [4] Kusano, T., Swanson, C. A., *Asymptotic theory of singular semilinear elliptic equations*, Canad. Math. Bull. **27** (1984), 223–232.
- [5] Neuman, F., *Oscillatory behavior of iterative linear ordinary differential equations depends on their order*, Arch. Math. (Brno) **22** (4) (1986), 187–192.
- [6] Pólya, G., *On the mean-value theorem corresponding to a given linear homogeneous differential equations*, Trans. Amer. Math. Soc. **24** (1922), 312–324.

DEPARTMENT OF MATHEMATICS, COMENIUS UNIVERSITY,
 MLYNSKÁ DOLINA, BRATISLAVA, SLOVAKIA
 E-mail: kristina.rostas@fmph.uniba.sk