

Ivan Chajda
Pseudocomplemented and Stone Posets

Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica, Vol. 51 (2012),
No. 1, 29--34

Persistent URL: <http://dml.cz/dmlcz/142871>

Terms of use:

© Palacký University Olomouc, Faculty of Science, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Pseudocomplemented and Stone Posets^{*}

Ivan CHAJDA

*Department of Algebra and Geometry, Faculty of Science, Palacký University
17. listopadu 12, 771 46 Olomouc, Czech Republic
e-mail: ivan.chajda@upol.cz*

(Received September 25, 2011)

Abstract

We show that every pseudocomplemented poset can be equivalently expressed as a certain algebra where the operation of pseudocomplementation can be characterized by means of remaining two operations which are binary and nullary. Similar characterization is presented for Stone posets.

Key words: pseudocomplement, pseudocomplemented poset, Stone poset

2000 Mathematics Subject Classification: 06A06, 06A11, 06D15

The concept of pseudocomplement was introduced by O. Frink [2] for meet-semilattices, Stone lattices were studied by R. Balbes and A. Horn [1]. S. K. Nimbhokar and A. Rahemani [3] modified the approach developed for posets by P. V. Venkatarasimhan [4] and use it for characterization of Stone join-semilattices.

The aim of this paper is to get another approach which goes in a sense conversely. We will show that every pseudocomplemented poset can be organized in a certain algebra. This can be analogously done for Stone posets.

Let us recall that the concept of pseudocomplement in a poset with the least element 0 was introduced in [4] by means of order-ideals. However, it can be easily paraphrased as follows.

Definition 1 Let $\mathcal{P} = (P; \leq, 0)$ be a poset with the least element 0, let $a \in P$. We say that $a^* \in P$ is a *pseudocomplement* of a if

- (i) there exists the infimum $a \wedge a^*$ of $\{a, a^*\}$ and is equal to 0;
- (ii) if $b \in P$ and $a \wedge b$ exists and equals 0, then $b \leq a^*$.

^{*}Supported by the project Algebraic Methods in Quantum Logic, CZ.1.07/2.3.00/20.0051.

A poset $\mathcal{P} = (P; \leq, 0)$ is called *pseudocomplemented* if there exists a pseudocomplement a^* for each $a \in P$. This fact will be expressed by notation $\mathcal{P} = (P; \leq, 0, *)$.

Convention In what follows, the notation $a \wedge b = c$ will be read as “the infimum $a \wedge b$ exists and is equal to c ”.

Example 1 Consider the poset $\mathcal{P} = (\{0, a, b, c, d, 1\}; \leq, 0)$ visualized in Fig. 1:




Fig. 1

Evidently, \mathcal{P} is neither a lattice nor a meet-semilattice. However, \mathcal{P} is pseudocomplemented and the pseudocomplements are determined by Definition 1 as follows

x	0	a	b	c	d	1
x^*	1	b	a	0	0	0

The following is a trivial consequence of the definition.

Lemma 1 Let $\mathcal{P} = (P; \leq, 0)$ be a pseudocomplemented poset. Then

- (a) \mathcal{P} has the greatest element $1 = 0^*$;
- (b) $x \leq x^{**}$, $x^{***} = x^*$ and if $x \leq y$, then $y^* \leq x^*$, for all $x, y \in P$.

We show now that a certain algebra of type $(2, 0)$ can be assigned to every poset $\mathcal{P} = (P; \leq, 0)$.

Definition 2 Let $\mathcal{P} = (P; \leq, 0)$ be a poset with the least element 0. Define a binary operation \sqcap on \mathcal{P} as follows: if $x \wedge y$ exists, then $x \sqcap y = x \wedge y$, and $x \sqcap y = 0$ otherwise. The algebra $\mathcal{A}(P) = (P; \sqcap, 0)$ will be called a \mathcal{P} -algebra.

Example 2 Consider the poset $\mathcal{P} = (\{0, a, b, c, d, 1\}; \leq, 0)$ of Example 1 (visualized in Fig. 1). Then the corresponding \mathcal{P} -algebra $\mathcal{A}(P) = (\{0, a, b, c, d, 1\}; \sqcap, 0)$ is defined uniquely by the operation table

\sqcap	0	a	b	c	d	1
0	0	0	0	0	0	0
a	0	a	0	a	a	a
b	0	0	b	b	b	b
c	0	a	b	c	0	c
d	0	a	b	0	d	d
1	0	a	b	c	d	1

Remark 1 (a) It is obvious that the operation \sqcap is commutative, i.e. $x \sqcap y = y \sqcap x$ for all $x, y \in P$.

(b) If $x \leq y$ then $x \wedge y$ exists and $x \wedge y = x$, i.e. also $x \sqcap y = x$. Conversely, if $x \sqcap y = x$ then either $x \wedge y$ exists, i.e. $x \wedge y = x$ and hence $x \leq y$, or $x \wedge y$ does not exist, i.e. $0 = x \sqcap y = x$ whence $x = 0 \leq y$ again. Thus we have

$$x \leq y \quad \text{if and only if} \quad x \sqcap y = x$$

in every \mathcal{P} -algebra $\mathcal{A}(P) = (P; \sqcap, 0)$.

Now, we prove that also conversely, every poset $\mathcal{P} = (P; \leq, 0)$ can be derived from its assigned \mathcal{P} -algebra $\mathcal{A}(P)$. For this, we characterize the operation \sqcap of $\mathcal{A}(P)$ by several simple axioms.

Lemma 2 *Let $\mathcal{P} = (P; \leq, 0)$ be a poset with 0 and $\mathcal{A}(P) = (P; \sqcap, 0)$ the corresponding \mathcal{P} -algebra. Then the operations \sqcap and 0 satisfy the following conditions:*

(A0) $x \sqcap 0 = 0$

(A1) $x \sqcap x = x$

(A2) $x \sqcap y = y \sqcap x$

(A3) $x \sqcap ((x \sqcap y) \sqcap z) = (x \sqcap y) \sqcap z$

(A4) *if there exists an element t such that (a) $x \sqcap t = t = y \sqcap t$ and (b) for all w , $x \sqcap w = w = y \sqcap w$ implies $w \sqcap t = w$, then $x \sqcap y = t$, and if such an element does not exist, then $x \sqcap y = 0$.*

Proof By Remark 1 we have $x \leq y$ iff $x \sqcap y = x$. Since 0 is the least element of \mathcal{P} , we have $x \sqcap 0 = 0$ which is (A0). The conditions (A1), (A2) follow directly by Definition 2. Further, $x \sqcap y \leq x$ and $(x \sqcap y) \sqcap z \leq x$, thus $x \sqcap ((x \sqcap y) \sqcap z) = x \wedge ((x \sqcap y) \sqcap z) = (x \sqcap y) \sqcap z$ which is (A3). For (A4), assume that such an element t exists in \mathcal{P} . Then, by (a), $t \leq x$, $t \leq y$ and, by (b), it is the greatest element in P of this property, i.e. $t = x \wedge y$ and hence $x \sqcap y = t$. If it does not exist, then $x \sqcap y = 0$, proving (A4). \square

Lemma 3 *Let $\mathcal{A} = (A; \sqcap, 0)$ be an algebra of type $(2, 0)$ satisfying (A0)–(A4). Define $x \leq y$ if $x \sqcap y = x$. Then $\mathcal{P}(A) = (A; \leq, 0)$ is a poset with the least element 0 and $x \sqcap y = x \wedge y$ provided $x \wedge y$ exists, and $x \sqcap y = 0$ otherwise.*

Proof By (A0) and (A2) we have $0 \leq x$ for each $x \in A$. By (A1) we obtain $x \leq x$, reflexivity of \leq . Assume $x \leq y$ and $y \leq z$. Then, by (A2), $x = x \sqcap y = y \sqcap x = y$ proving antisymmetry of \leq . If $x \leq y$ and $y \leq z$, i.e. $x \sqcap y = x$ and $y \sqcap z = y$, then by (A2) and (A3) we derive $x \sqcap z = (x \sqcap y) \sqcap z = (x \sqcap (y \sqcap z)) \sqcap z = x \sqcap (y \sqcap z) = x \sqcap y = x$ whence \leq is also transitive, i.e. it is a partial order on A , thus $(A; \leq, 0)$ is a poset with the least element 0.

Assume now that $a, b \in A$ and $a \wedge b$ exists (with respect to the aforementioned order \leq). Then for $t = a \wedge b$ the assumptions of (A4) are satisfied and hence

$a \sqcap b = t = a \wedge b$. If $a \wedge b$ does not exist, then there is no $t \in A$ satisfying the assumptions of (A4) and hence $a \sqcap b = 0$. \square

Let $\mathcal{A} = (A; \sqcap, 0)$ be an algebra satisfying (A0)–(A4). The poset $\mathcal{P}(A) = (A; \leq, 0)$ derived in Lemma 3 will be called the *induced poset*. We are going to show that posets \mathcal{P} with 0 and the corresponding \mathcal{P} -algebras are in a one-to-one correspondence.

Lemma 4 *Let $\mathcal{P} = (P; \leq, 0)$ be a poset with 0, $\mathcal{A}(P) = (P; \sqcap, 0)$ the \mathcal{P} -algebra and $\mathcal{P}(\mathcal{A}(P)) = (P; \sqsubseteq, 0)$ the induced poset. Then $\mathcal{P} = \mathcal{P}(\mathcal{A}(P))$.*

Let $\mathcal{A} = (A; \sqcap, 0)$ be an algebra satisfying (A0)–(A4), $\mathcal{P}(A) = (A; \leq, 0)$ the induced poset and $\mathcal{A}(\mathcal{P}(A)) = (A; \sqcap, 0)$ its $\mathcal{P}(A)$ -algebra. Then $\mathcal{A} = \mathcal{A}(\mathcal{P}(A))$.

Proof (a) We need to show $\leq = \sqsubseteq$. Assume $x \leq y$ in \mathcal{P} . By Remark 1, this is equivalent to $x \sqcap y = x$ in $\mathcal{A}(P)$ which is equivalent by definition to $x \sqsubseteq y$. Hence $\mathcal{P} = \mathcal{P}(\mathcal{A}(P))$.

(b) Assume $a \wedge b$ exists in $\mathcal{P}(A)$. Then $a \sqcap b = a \wedge b$ in $\mathcal{A}(\mathcal{P}(A))$ but also $a \sqcap b = a \wedge b$ in \mathcal{A} by Lemma 3. In both cases, we obtain $a \sqcap b = a \sqcap b$ and hence $\mathcal{A} = \mathcal{A}(\mathcal{P}(A))$. \square

Now, we are ready to characterize pseudocomplementation in posets by means of the corresponding \mathcal{P} -algebra.

Theorem 1 *Let $\mathcal{P} = (P; \leq, 0)$ be a poset with the least element 0, let $\mathcal{A}(P) = (P; \sqcap, 0)$ be its \mathcal{P} -algebra. Let $*$ be a unary operation on P . Then $\mathcal{P} = (P; \leq, 0, *)$ is a pseudocomplemented poset if and only if $(P; \sqcap, *, 0)$ satisfies the following conditions:*

$$(P1) \quad x \sqcap 0^* = x$$

$$(P2) \quad x \sqcap (x^* \sqcap y) = 0$$

$$(P3) \quad \text{if } x \sqcap (y \sqcap z) = 0 \text{ for all } z \in P, \text{ then } y \sqcap x^* = y$$

Proof Assume that $\mathcal{P} = (P; \leq, 0, *)$ is a pseudocomplemented poset. Then for each $x, y \in P$ we have $x^* \sqcap y \leq x^*$. Since $x \wedge x^*$ exists and is equal to 0, we conclude that also $x \wedge (x^* \sqcap y)$ exists and is equal to 0, i.e. $x \sqcap (x^* \sqcap y) = x \wedge (x^* \sqcap y)$ proving (P2). Assume $x \sqcap (y \sqcap z) = 0$ for each $z \in P$. If there exists $c \in P$ such that $c \neq 0$ and $x \sqcap y = c$ then, by (A2), (A3) and the assumption, $0 = x \sqcap (y \sqcap c) = x \sqcap (y \sqcap (x \sqcap y)) = y \sqcap (x \sqcap y) = x \sqcap y = c \neq 0$, a contradiction. Therefore $x \wedge y = 0$ whence $y \leq x^*$ and $y \sqcap x^* = y$ proving (P3). The condition (P1) is evident.

Conversely, let $(P; \sqcap, *, 0)$ satisfy (P1), (P2) and (P3). By (P1), 0^* is the greatest element of \mathcal{P} . If $y \leq x$ and $y \leq x^*$ then, according to (P2), we obtain $y = x \sqcap y = x \sqcap (x^* \sqcap y) = 0$. Hence $x \wedge x^* = 0$. Assume now $x \wedge z = 0$. Then $x \sqcap (z \sqcap c) \leq x$ and $x \sqcap (z \sqcap c) \leq z \sqcap c \leq z$ for each c , thus $x \sqcap (z \sqcap c) \leq x \wedge z = 0$. By (P3) we conclude $z \leq x^*$, i.e. x^* is the greatest element of P satisfying $x \wedge z = 0$, i.e. it is the pseudocomplement of x . \square

We focus our attention on Stone posets in the rest of the paper. As in the previous case, the definition of [4] can be paraphrased as follows.

Definition 3 Let $\mathcal{P} = (P; \leq, 0, *)$ be a pseudocomplemented poset. Then \mathcal{P} is called a *Stone poset* if for each $x \in P$ the supremum $x^* \vee x^{**}$ exists and equals 1 (where $1 = 0^*$).

Example 3 The poset from Example 1 is pseudocomplemented, but it is not a Stone one because, e.g., $a^* \vee a^{**} = b \vee a$ does not exist.

Example 4 Consider the poset $\mathcal{P} = (\{0, a, b, c, d, p, q, 1\}; \leq, 0)$ depicted in Fig. 2.




Fig. 2

Then \mathcal{P} is pseudocomplemented, pseudocomplements are given by the table:

x	0	p	q	d	a	b	c	r	1
x^*	1	c	c	a	c	0	a	c	0
x^{**}	0	a	a	c	a	1	c	a	1

Since $a \vee c = 1$ and $0 \vee 1 = 1$, we have $x^* \vee x^{**} = 1$ for each $x \in P$, thus \mathcal{P} is a Stone poset.

We proceed analogously as in the previous case. Consider a bounded poset $\mathcal{P} = (P; \leq, 0, 1)$. The operation \sqcap on P is defined by Definition 2. Now we define \sqcup on P dually: if $x \vee y$ exists, then $x \sqcup y = x \vee y$, and $x \sqcup y = 1$ otherwise. The algebra $\mathcal{B}(P) = (P; \sqcup, \sqcap, 0, 1)$ will be called the \mathcal{P}_1 -algebra assigned to \mathcal{P} .

Remark 2 Analogously as in the previous case, one can easily check that \sqcup has the properties:

- (B0) $x \sqcup 1 = 1$
- (B1) $x \sqcup x = x$
- (B2) $x \sqcup y = y \sqcup x$
- (B3) $x \sqcup ((x \sqcup y) \sqcup z) = (x \sqcup y) \sqcup z$
- (B4) if there exists an element s such that (a) $x \sqcup s = s = y \sqcup s$ and (b) for all u , $x \sqcup u = u = y \sqcup u$ implies $u \sqcup s = u$, then $x \sqcup y = s$, and if such an element does not exist, then $x \sqcup y = 1$.