

Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium.
Mathematica

Maslina Darus; Imran Faisal

Problems Concerning Subclasses of Analytic Functions

Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica, Vol. 51 (2012),
No. 1, 35-42

Persistent URL: <http://dml.cz/dmlcz/142872>

Terms of use:

© Palacký University Olomouc, Faculty of Science, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to
digitized documents strictly for personal use. Each copy of any part of this document must
contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped
with digital signature within the project *DML-CZ: The Czech Digital
Mathematics Library* <http://project.dml.cz>

Problems Concerning Subclasses of Analytic Functions *

Maslina DARUS¹, Imran FAISAL²

*School of Mathematical Sciences, Faculty of Science and Technology
Universiti Kebangsaan Malaysia, Bangi 43600 Selangor D. Ehsan, Malaysia*

¹e-mail: maslina@ukm.my

²e-mail: faisalmath@gmail.com

(Received March 27, 2011)

Abstract

In the present paper, we establish some interesting results concerning the quasi-Hadamard product for certain subclasses of analytic functions.

Key words: analytic, Starlike, Convex, Quasi-Hadamard product

2000 Mathematics Subject Classification: 30C45

1 Introduction and preliminaries

Throughout this paper, let the of functions of the form

$$f(z) = a_1 z + \sum_{n=2}^{\infty} a_n z^n, \quad a_1 > 0, \quad a_n \geq 0, \quad (1)$$

$$f_i(z) = a_{1,i} z + \sum_{n=2}^{\infty} a_{n,i} z^n, \quad a_{1,i} > 0, \quad a_{n,i} \geq 0, \quad (2)$$

$$g(z) = b_1 z + \sum_{n=2}^{\infty} b_n z^n, \quad b_1 > 0, \quad b_n \geq 0, \quad (3)$$

$$g_j(z) = b_{1,j} z + \sum_{n=2}^{\infty} b_{n,j} z^n, \quad b_{1,i} > 0, \quad b_{n,j} \geq 0, \quad (4)$$

be regular and univalent in the unit disc $\mathbb{U} = \{z: |z| < 1\}$.

*Supported by UKM-ST-06-FRGS0244-2010.

For a function $f(z)$ defined by (1) (with $a_1 = 1$) we define

$$\begin{aligned} D^0 f(z) &= f(z), \\ D_\lambda^1(\alpha, \beta, \mu) f(z) &= (\frac{\alpha-\mu+\beta-\lambda}{\alpha+\beta})f(z) + (\frac{\mu+\lambda}{\alpha+\beta})zf'(z), \\ D_\lambda^2(\alpha, \beta, \mu) f(z) &= D(D_\lambda^1(\alpha, \beta, \mu)f(z)), \\ &\vdots \\ D_\lambda^k(\alpha, \beta, \mu) f(z) &= D(D_\lambda^{k-1}(\alpha, \beta, \mu)f(z)). \end{aligned} \quad (5)$$

If f is given by (1) then from (5) we have

$$D_\lambda^k(\alpha, \beta, \mu) f(z) = z + \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^k a_n z^n, \quad (6)$$

where $f \in A$, $\alpha, \beta, \mu, \lambda \geq 0$, $\alpha + \beta \neq 0$, $n \in N_o$. The operator $D_\lambda^k(\alpha, \beta, \mu)f(z)$ was introduced by authors, Darus and Faisal in [7].

By specializing the parameters of $D_\lambda^n(\alpha, \beta, \mu)f(z)$ we get the following differential operators. If we substitute

- $\beta = 0$, we get $D_\lambda^k(\alpha, 0, \mu)f(z) = z + \sum_{n=2}^{\infty} (\frac{\alpha+(\mu+\lambda)(n-1)}{\alpha})^k a_n z^n$ of differential operator given by Darus and Faisal [8].
- $\beta = 1$, $\mu = 0$, we get $D_\lambda^k(\alpha, 1, 0)f(z) = z + \sum_{n=2}^{\infty} (\frac{\alpha+\lambda(n-1)+1}{\alpha+1})^k a_n z^n$ of differential operator given by Aouf et. al [3].
- $\alpha = 1$, $\beta = o$, and $\mu = 0$, we get $D_\lambda^k(1, 0, 0)f(z) = z + \sum_{n=2}^{\infty} (1+\lambda(n-1))^k a_n z^n$ of differential operator given by Al-Oboudi [1].
- $\alpha = 1$, $\beta = o$, $\mu = 0$ and $\lambda = 1$, we get $D_1^k(1, 0, 0)f(z) = z + \sum_{n=2}^{\infty} (n)^k a_n z^n$ of differential operator given by Sălăgean's [12].
- $\alpha = 1$, $\beta = 1$, $\lambda = 1$ and $\mu = 0$, we get $D_1^k(1, 1, 0)f(z) = z + \sum_{n=2}^{\infty} (\frac{n+1}{2})^k a_n z^n$ of differential operator given by Uralegaddi and Somanatha [16].
- $\beta = 1$, $\lambda = 1$ and $\mu = 0$, we get $D_1^k(\alpha, 1, 0)f(z) = z + \sum_{n=2}^{\infty} (\frac{n+\alpha}{\alpha+1})^k a_n z^n$ of differential operator given by Cho and Srivastava [5, 6].

With the help of the differential operator $D_\lambda^k(\alpha, \beta, \mu)f(z)$, we define the classes $\Psi(k, \rho, \delta, \eta)$, and $R_p(n, \rho, \delta, \eta)$ as follows:

For, $0 \leq \rho < 1$, $0 \leq \delta < 1$ and $\eta \geq 0$, we let $\Psi(k, \rho, \delta, \eta)$ denote the class of functions $f(z)$ defined by (1) and satisfying the analytic criterion

$$\begin{aligned} &\Re \left\{ \frac{z(D_\lambda^k(\alpha, \beta, \mu)f(z))'}{(1-\rho)(D_\lambda^k(\alpha, \beta, \mu)f(z)) + \rho z(D_\lambda^k(\alpha, \beta, \mu)f(z))'} - \delta \right\} \\ &> \eta \left| \frac{z(D_\lambda^k(\alpha, \beta, \mu)f(z))'}{(1-\rho)(D_\lambda^k(\alpha, \beta, \mu)f(z)) + \rho z(D_\lambda^k(\alpha, \beta, \mu)f(z))'} - 1 \right|, \quad z \in \mathbb{U}. \end{aligned} \quad (7)$$

Suitably specializing the parameters of the class $\Psi(k, \rho, \delta, \eta)$ generalize the classes defined by some well known authors see [13]–[15].

Using similar arguments as given in [11], we can easily get the following results for functions in the classes $\Psi(n, \rho, \delta, \eta)$ and $\psi(k+1, \rho, \delta, \eta)$ respectively.

A function $f(z) \in \Psi(k, \rho, \delta, \eta)$ ($0 \leq \rho < 1, 0 \leq \delta < 1, \eta \geq 0$) if and only if

$$\begin{aligned} & \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^k [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] |a_n| \\ & \leq (1-\delta), \end{aligned} \quad (8)$$

Now we introduce the following class of analytic functions which plays an important role in our discussion as follows.

A function $f(z)$ which is analytic in U belongs to the class $R_p(n, \rho, \delta, \eta)$ if and only if

$$\begin{aligned} & \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^p [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] |a_n| \\ & \leq (1-\delta), \end{aligned} \quad (9)$$

where $0 \leq \rho < 1, 0 \leq \delta < 1, \eta \geq 0$ and p is any fixed nonnegative real number.

For $p = k$, it is identical to $\Psi(k, \rho, \delta, \eta)$. Also for $p = \beta = \mu = 0$, it is identical to $S_p(\rho, \delta, \eta)$ defined in [11]. Further, for any positive integer $p > h > h-1 > \dots > k+1 > k$, we have the inclusion relation

$$R_p(n, \rho, \delta, \eta) \subseteq R_h(n, \rho, \delta, \eta) \subseteq \dots \subseteq \psi(k+1, \rho, \delta, \eta) \subseteq \Psi(k, \rho, \delta, \eta).$$

The class $R_p(n, \rho, \delta, \eta)$ is nonempty for any nonnegative real number p as the functions of the form

$$f(z) = a_1 z$$

$$+ \sum_{n=2}^{\infty} \frac{(\alpha + \beta)^p (1-\delta)}{\left(\alpha + (\mu + \lambda)(n-1) + \beta \right)^p [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)]} \lambda_n z^n, \quad (10)$$

where $a_1 > 0, \lambda_n \geq 0$ and $\sum_2^{\infty} \lambda_n \leq 1$; satisfy the inequality (10).

Let us define the quasi-Hadamard product of the functions $f(z)$ and $g(z)$ by

$$f * g(z) = a_1 b_1 z + \sum_{n=2}^{\infty} a_n b_n z^n, \quad a_1 > 0, b_1 > 0, a_n \geq 0. \quad (11)$$

In this work we establish certain results concerning the quasi-Hadamard product of functions for the classes $\Psi(k, \rho, \delta, \eta)$ and $R_p(n, \rho, \delta, \eta)$ analogous to the results due to V. Kumar [9] and [10] as well.

2 The main results

Theorem 1 Let the functions $f_i(z)$ defined by (2) be in the class $\psi(k+1, \rho, \delta, \eta)$ for every $i = 1, 2, \dots, r$ and let the functions $g_j(z)$ defined by (4) be in the class $\Psi(n, \rho, \delta, \eta)$ for every $j = 1, 2, \dots, q$. Then the quasi-Hadamard product $f_1 * f_2 * \dots * f_r * g_1 * g_2 * \dots * g_q(z)$ belongs to the class $R_{r(k+2)+q(k+1)-1}(n, \rho, \delta, \eta)$.

Proof We denote the quasi-Hadamard product $f_1 * f_2 * \dots * f_r * g_1 * g_2 * \dots * g_q(z)$ by the function $G(z)$ for the sake of convenience. Clearly,

$$G(z) = \left[\prod_{i=1}^r |a_{1,i}| \right] \left[\prod_{j=1}^q |b_{1,j}| \right] z + \sum_{n=2}^{\infty} \left[\prod_{i=1}^r |a_{n,i}| \right] \left[\prod_{j=1}^q |b_{n,j}| \right] z^n.$$

Since $f_i(z) \in \psi(k+1, \rho, \delta, \eta)$ implies

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^{k+1} [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] |a_{n,i}| \\ \leq (1-\delta) |a_{1,i}|, \end{aligned} \quad (12)$$

for every $i = 1, 2, \dots, r$. Implies

$$|a_{n,i}| \leq \frac{(\alpha + \beta)^{k+1} (1-\delta)}{\left(\alpha + (\mu + \lambda)(n-1) + \beta \right)^{k+1} [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)]} |a_{1,i}|,$$

for every $i = 1, 2, \dots, r$. Implies

$$|a_{n,i}| \leq \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^{-k-2} |a_{1,i}|, \quad (13)$$

for every $i = 1, 2, \dots, r$. Similarly $g_j(z) \in \Psi(k, \rho, \delta, \eta)$, implies

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^k [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] |b_{n,j}| \\ \leq (1-\delta) |b_{1,j}|, \end{aligned} \quad (14)$$

for every $i = 1, 2, \dots, q$. Implies

$$|b_{n,j}| \leq \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^{-k-1} |b_{1,j}|, \quad (15)$$

for every $j = 1, 2, \dots, q$. To prove the theorem, we need to show that

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^t [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] \\ \times \left[\prod_{i=1}^r |a_{n,i}| \right] \left[\prod_{j=1}^q |b_{n,j}| \right] \leq (1-\delta) \left[\prod_{i=1}^r |a_{1,i}| \right] \left[\prod_{j=1}^q |b_{1,j}| \right], \end{aligned}$$

where $t = r(k+2) + q(k+1) - 1$.

Using (13), (14) and (15) for $i = 1, 2, \dots, r$, $j = q$ and $j = 1, 2, \dots, q-1$ respectively. we have (consider $t = r(k+2) + q(k+1) - 1$)

$$\begin{aligned}
& \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^t [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] \\
& \quad \times \left[\prod_{i=1}^r |a_{n,i}| \right] \left[\prod_{j=1}^q |b_{n,j}| \right] \\
& \leq \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^{r(k+2)+q(k+1)-1} [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] \\
& \quad \times \left[\left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^{-r(k+2)} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^{-(q-1)(k+1)} \right. \\
& \quad \times |b_{n,q}| \left. \prod_{i=1}^r |a_{1,i}| \right] \left[\prod_{j=1}^{q-1} |b_{1,j}| \right] = \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^k \\
& \quad \times [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] |b_{n,q}| \left[\prod_{i=1}^r |a_{1,i}| \right] \left[\prod_{j=1}^{q-1} |b_{1,j}| \right] \\
& \leq (1-\delta) \left[\prod_{i=1}^r |a_{1,i}| \right] \left[\prod_{j=1}^q |b_{1,j}| \right].
\end{aligned}$$

Implies

$$\begin{aligned}
& \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n-1) + \beta}{\alpha + \beta} \right)^t [n(1+\eta) - (\delta + \eta)(1+n\rho - \rho)] \\
& \quad \times \left[\prod_{i=1}^r |a_{n,i}| \right] \left[\prod_{j=1}^q |b_{n,j}| \right] \\
& \leq (1-\delta) \left[\prod_{i=1}^r |a_{1,i}| \right] \left[\prod_{j=1}^q |b_{1,j}| \right], \quad t = r(k+2) + q(k+1) - 1.
\end{aligned}$$

Using (9) we conclude that

$$f_1 * f_2 * \dots * f_r * g_1 * g_2 * \dots * g_q(z) \in R_{r(k+2)+q(k+1)-1}(n, \rho, \delta, \eta). \quad \square$$

Theorem 2 Let the functions $f_i(z)$ defined by (2) be in the class $\psi(k+1, \rho, \delta, \eta)$ for every $i = 1, 2, \dots, r$. Then the Hadamard product $f_1(z) * f_2(z) * \dots * f_r(z)$ belongs to the class $R_{r(k+2)-1}(n, \rho, \delta, \eta)$.

Proof To prove the theorem, we need to show that

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{r(k+2)-1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] \prod_{i=1}^r |a_{n,i}| \\ \leq (1 - \delta) \prod_{i=1}^r |a_{1,i}|. \end{aligned}$$

Since $f_i(z) \in \psi(k + 1, \rho, \delta, \eta)$, implies

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{k+1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] |a_{n,i}| \\ \leq (1 - \delta) |a_{1,i}|, \end{aligned} \quad (16)$$

for every $i = 1, 2, \dots, r$. Implies

$$|a_{n,i}| \leq \frac{(\alpha + \beta)^{k+1}(1 - \delta)}{(\alpha + (\mu + \lambda)(n - 1) + \beta)^{k+1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)]} |a_{1,i}|,$$

for every $i = 1, 2, \dots, r$. Implies

$$|a_{n,i}| \leq \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{-k-2} |a_{1,i}|, \quad (17)$$

for every $i = 1, 2, \dots, r$.

Simultaneously applying (16) as well as (17) for $i = r$ and $i = 1, 2, \dots, r - 1$ respectively, we get

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{r(k+2)-1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] \prod_{i=1}^r |a_{n,i}| \\ \leq \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{r(k+2)-1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] |a_{n,r}| \\ \times \left[\left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{-(r-1)(k+2)} \prod_{i=1}^{r-1} |a_{1,i}| \right] \\ = \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{k+1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] |a_{n,r}| \left[\prod_{i=1}^{r-1} |a_{1,i}| \right] \\ = (1 - \delta) \prod_{i=1}^r |a_{1,i}|. \end{aligned}$$

Hence $f_1(z) * f_2(z) * \dots * f_r(z) \in R_{r(k+2)-1}(k, \rho, \delta, \eta)$. \square

Theorem 3 Let the functions $f_i(z)$ defined by (2) be in the class $\Psi(k, \rho, \delta, \eta)$ for every $i = 1, 2, \dots, r$. Then the Hadamard product $f_1(z) * f_2(z) * \dots * f_r(z)$ belongs to the class $R_{r(k+1)-1}(n, \rho, \delta, \eta)$.

Proof Since $f_i(z) \in \Psi(k, \rho, \delta, \eta)$, implies

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^k [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] |a_{n,i}| \\ \leq (1 - \delta) |a_{1,i}|, \end{aligned} \quad (18)$$

for every $i = 1, 2, \dots, r$. Implies

$$\left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^k [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] |a_{n,i}| \leq (1 - \delta) |a_{1,i}|,$$

or

$$|a_{n,i}| \leq \frac{(1 - \delta)}{\left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^k [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)]} |a_{1,i}|,$$

for every $i = 1, 2, \dots, r$. Implies

$$|a_{n,i}| \leq \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{-k-1} |a_{1,i}|, \quad (19)$$

for every $i = 1, 2, \dots, r$.

To prove that $f_1(z) * f_2(z) * \dots * f_r(z) \in R_{r(k+1)-1}(n, \rho, \delta, \eta)$, it is enough to show that

$$\begin{aligned} \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{r(k+1)-1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] \prod_{i=1}^r |a_{n,i}| \\ \leq (1 - \delta) \prod_{i=1}^r |a_{1,i}|. \end{aligned} \quad (20)$$

Using (18) and (19) for $i = r$ and $i = 1, 2, \dots, r - 1$ respectively, then we have

$$\begin{aligned} & \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{r(k+1)-1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] \prod_{i=1}^r |a_{n,i}| \\ & \leq \sum_{n=2}^{\infty} \left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{r(k+1)-1} [n(1 + \eta) - (\delta + \eta)(1 + n\rho - \rho)] |a_{n,r}| \\ & \quad \times \left[\left(\frac{\alpha + (\mu + \lambda)(n - 1) + \beta}{\alpha + \beta} \right)^{-(r-1)(k+1)} \prod_{i=1}^{r-1} |a_{1,i}| \right] = (1 - \delta) \prod_{i=1}^r |a_{1,i}|. \end{aligned}$$

Hence $f_1(z) * f_2(z) * \dots * f_r(z) \in R_{r(k+1)-1}(n, \rho, \delta, \eta)$. \square

References

- [1] Al-oboudi, F. M.: *On univalent functions defined by a generalized Salagean operator.* Inter. J. Math. Math. Sci. **27** (2004), 1429–1436.
- [2] Altintas, O., Owa, S.: *On subclasses of univalent functions with negative coefficients.* Pusan Kyongnam Math. J. **4** (1998), 41–56.
- [3] Aouf, M. K., El-Ashwah, R. M., El-Deeb, S. M.: *Some inequalities for certain p -valent functions involving extended multiplier transformations.* Proc. Pakistan Acad. Sci. **46** (2009), 217–221.
- [4] Bharati, R., Parvatham, R., Swaminathan, A.: *On subclasses of uniformly convex functions and corresponding class of starlike functions.* Tamkang J. Math. **26**, 1 (1997), 17–32.
- [5] Cho, N. E., Srivastava, H. M.: *Argument estimates of certain analytic functions defined by a class of multiplier transformations.* Math. Comput. Modeling **37** (2002), 39–49.
- [6] Cho, N. E., Kim, T. H.: *Multiplier transformations and strongly close-to-convex functions.* Bull. Korean Math. Soc. **40** (2003), 399–410.
- [7] Darus, M., Faisal, I.: *A different approach to normalized analytic functions through meromorphic functions defined by extended multiplier transformations operator.* Int. J. App. Math. Stat. **23** (2011), 112–121.
- [8] Darus, M., Faisal, I.: *Characterization properties for a class of analytic functions defined by generalized Cho and Srivastava operator.* In: Proc. 2nd Inter. Conf. Math. Sci., Kuala Lumpur, Malaysia, 2010, 1106–1113.
- [9] Kumar, V.: *Hadamard product of certain starlike functions.* J. Math. Anal. Appl. **110** (1985), 425–428.
- [10] Kumar, V.: *Quasi-Hadamard product of certain univalent function.* J. Math. Anal. Appl. **126** (1987), 70–77.
- [11] Murugusundaramoorthy, G., Mageshb, N.: *On certain subclasses of analytic functions associated with hypergeometric functions.* App. Math. Lett. **24** (2011), 494–500.
- [12] Salagean, G. S.: Subclasses of univalent functions. Lecture Notes in Mathematics 1013, Springer-Verlag, Berlin, 1983, 362–372.
- [13] Silverman, H.: *Univalent functions with negative coefficients.* Proc. Amer. Math. Soc. **51**, (1975), 109–116.
- [14] Subramanian, K. G., Sudharsan, T. V., Balasubrahmanyam, P., Silverman, H.: *Classes of uniformly starlike functions.* Publ. Math. Debrecen. **53**, 3-4 (1998), 309–315.
- [15] Subramanian, K. G., Murugusundaramoorthy, G., Balasubrahmanyam, P., Silverman, H.: *Subclasses of uniformly convex and uniformly starlike functions.* Math. Japonica **42**, 3 (1995), 517–522.
- [16] Uralegaddi, B. A., Somanatha, C.: *Certain classes of univalent functions.* In: Srivastava H. M., Owa S. (eds) Current Topics in Analytic Function Theory, World Scientific, Singapore, 1992, 371–374.