

# Applications of Mathematics

---

Abdallah El Farissi; Benharrat Belaidi

On the growth of solutions of some higher order linear differential equations

*Applications of Mathematics*, Vol. 57 (2012), No. 4, 377–390

Persistent URL: <http://dml.cz/dmlcz/142905>

## Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE GROWTH OF SOLUTIONS OF SOME HIGHER ORDER  
LINEAR DIFFERENTIAL EQUATIONS

ABDALLAH EL FARISSI, BENHARRAT BELAÏDI, Mostaganem

(Received April 12, 2010)

*Abstract.* In this paper we discuss the growth of solutions of the higher order nonhomogeneous linear differential equation

$$f^{(k)} + A_{k-1}f^{(k-1)} + \dots + A_2f'' + (D_1(z) + A_1(z)e^{az})f' + (D_0(z) + A_0(z)e^{bz})f = F \quad (k \geq 2),$$

where  $a, b$  are complex constants that satisfy  $ab(a-b) \neq 0$  and  $A_j(z)$  ( $j = 0, 1, \dots, k-1$ ),  $D_j(z)$  ( $j = 0, 1$ ),  $F(z)$  are entire functions with  $\max\{\varrho(A_j) \ (j = 0, 1, \dots, k-1), \varrho(D_j) \ (j = 0, 1)\} < 1$ . We also investigate the relationship between small functions and the solutions of the above equation.

*Keywords:* linear differential equations, entire solutions, order of growth, exponent of convergence of zeros, exponent of convergence of distinct zeros

*MSC 2010:* 34M10, 30D35

## 1. INTRODUCTION AND STATEMENT OF RESULTS

Throughout this paper, we assume that the reader is familiar with the fundamental results and the standard notation of the Nevanlinna value distribution theory and the basic notions of Wiman-Valiron as well (see [11], [12], [15]). In addition, we will use  $\lambda(f)$  ( $\lambda_2(f)$ ) and  $\bar{\lambda}(f)$  ( $\bar{\lambda}_2(f)$ ) to denote respectively the exponents (hyper-exponents) of convergence of the zero-sequence and the sequence of distinct zeros of  $f$ ,  $\varrho(f)$  to denote the order of growth of a meromorphic function  $f$  and  $\varrho_2(f)$  to denote the hyper-order of  $f$ . A meromorphic function  $\varphi(z)$  is called a small function with respect to  $f(z)$  if  $T(r, \varphi) = o(T(r, f))$  as  $r \rightarrow \infty$  except possibly a set of  $r$  of finite linear measure, where  $T(r, f)$  is the Nevanlinna characteristic function of  $f$ . If

$f$  is of infinite order and  $\varphi$  is of finite order, then clearly  $\varphi(z)$  is a small function with respect to  $f(z)$ . We also define

$$\bar{\lambda}(f - \varphi) = \limsup_{r \rightarrow \infty} \frac{\log \bar{N}\left(r, \frac{1}{f - \varphi}\right)}{\log r}$$

and

$$\bar{\lambda}_2(f - \varphi) = \limsup_{r \rightarrow \infty} \frac{\log \log \bar{N}\left(r, \frac{1}{f - \varphi}\right)}{\log r}$$

for any meromorphic function  $\varphi(z)$ .

For the second order linear differential equation

$$(1.1) \quad f'' + e^{-z}f' + B(z)f = 0,$$

where  $B(z)$  is an entire function, it is well known that each solution  $f$  of equation (1.1) is an entire function, and that if  $f_1, f_2$  are two linearly independent solutions of (1.1), then by [7], at least one of  $f_1, f_2$  is of infinite order. Hence, “most” solutions of (1.1) will have infinite order. But equation (1.1) with  $B(z) = -(1 + e^{-z})$  possesses a solution  $f(z) = e^z$  of finite order.

A natural question arises: What conditions on  $B(z)$  will guarantee that every solution  $f \not\equiv 0$  of (1.1) has infinite order? Many authors, Frei [8], Ozawa [16], Amemiya-Ozawa [1], Gundersen [9], and Langley [13] have studied this problem. They proved that when  $B(z)$  is a nonconstant polynomial or  $B(z)$  is a transcendental entire function with order  $\rho(B) \neq 1$ , then every solution  $f \not\equiv 0$  of (1.1) has infinite order.

In 2002, Z. X. Chen [5] considered the question: What conditions on  $B(z)$  when  $\rho(B) = 1$  will guarantee that every nontrivial solution of (1.1) has infinite order? He proved the following results, which improved results of Frei, Amemiya-Ozawa, Ozawa, Langley, and Gundersen.

**Theorem A** ([5]). *Let  $A_j(z) (\not\equiv 0)$  ( $j = 0, 1$ ) and  $D_j(z)$  ( $j = 0, 1$ ) be entire functions with  $\max\{\rho(A_j) (j = 0, 1), \rho(D_j) (j = 0, 1)\} < 1$ , and let  $a, b$  be complex constants that satisfy  $ab \neq 0$  and  $\arg a \neq \arg b$  or  $a = cb$  ( $0 < c < 1$ ). Then every solution  $f \not\equiv 0$  of the equation*

$$(1.2) \quad f'' + (D_1(z) + A_1(z)e^{az})f' + (D_0(z) + A_0(z)e^{bz})f = 0$$

*is of infinite order.*

Setting  $D_j \equiv 0$  ( $j = 0, 1$ ) in Theorem A, we obtain the following result.

**Theorem B.** Let  $A_j(z) (\neq 0) (j = 0, 1)$  be entire functions with  $\max\{\varrho(A_j) : j = 0, 1\} < 1$ , and let  $a, b$  be complex constants that satisfy  $ab \neq 0$  and  $\arg a \neq \arg b$  or  $a = cb (0 < c < 1)$ . Then every solution  $f \neq 0$  of the equation

$$(1.3) \quad f'' + A_1(z)e^{az}f' + A_0(z)e^{bz}f = 0$$

is of infinite order.

**Theorem C** ([5]). Let  $A_j(z) (\neq 0) (j = 0, 1)$  be entire functions with  $\varrho(A_j) < 1 (j = 0, 1)$ , and let  $a, b$  be complex constants that satisfy  $ab \neq 0$  and  $a = cb (c > 1)$ . Then every solution  $f \neq 0$  of equation (1.3) is of infinite order.

Very recently in [18], H. Y. Xu and T. B. Cao have investigated the growth of solutions of some higher order nonhomogeneous linear differential equations and have obtained the following result.

**Theorem D** ([18]). Let  $P(z) = \sum_{i=0}^n a_i z^i$  and  $Q(z) = \sum_{i=0}^n b_i z^i$  be nonconstant polynomials, where  $a_i, b_i (i = 0, 1, \dots, n)$  are complex numbers,  $a_n b_n (a_n - b_n) \neq 0$ . Suppose that  $h_i(z) (2 \leq i \leq k-1)$  are polynomials of degree not greater than  $n-1$  in  $z$ ,  $A_j(z) \neq 0 (j = 0, 1)$  and  $H(z) \neq 0$  are entire functions with  $\max\{\varrho(A_j) (j = 0, 1), \varrho(H)\} < n$ , and  $\varphi$  is an entire function of finite order. Then every nontrivial solution  $f$  of the equation

$$(1.4) \quad f^{(k)} + h_{k-1}f^{(k-1)} + \dots + h_2f'' + A_1(z)e^{P(z)}f' + A_0(z)e^{Q(z)}f = H$$

satisfies  $\varrho(f) = \lambda(f) = \bar{\lambda}(f) = \bar{\lambda}(f - \varphi) = \infty$  and  $\varrho_2(f) = \lambda_2(f) = \bar{\lambda}_2(f) = \bar{\lambda}_2(f - \varphi) \leq n$ .

**Remark 1.1.** In the original statement of Theorem D (see [18]), the condition  $H \neq 0$  must be added. Indeed, if  $H \equiv 0$ , then the conclusions of Theorem D are false. For example, the equation  $f''' - f'' - 2e^z f' - e^{3z} f = 0$  possesses the solution  $f(z) = e^{e^z}$  with  $\varrho(f) = \infty$  and  $\lambda(f) = 0$ .

It is natural to ask whether the polynomials  $h_{k-1}(z), \dots, h_2(z)$  in (1.4) can be replaced by entire functions of orders that are less than 1. The main purpose of this paper is to study the growth and the oscillation of solutions of the linear differential equation

$$(1.5) \quad f^{(k)} + A_{k-1}f^{(k-1)} + \dots + A_2f'' + (D_1(z) + A_1(z)e^{az})f' + (D_0(z) + A_0(z)e^{bz})f = F \quad (k \geq 2).$$

We obtain the following results.

**Theorem 1.1.** *Let  $a$  and  $b$  be complex numbers that satisfy  $ab(a - b) \neq 0$ . Suppose that  $A_j(z)$  ( $j = 0, 1, \dots, k - 1$ ),  $A_j(z) \not\equiv 0$  ( $j = 0, 1$ ),  $D_j(z)$  ( $j = 0, 1$ ) and  $F(z)$  are entire functions with  $\max\{\rho(A_j)$  ( $j = 0, 1, \dots, k - 1$ ),  $\rho(D_j)$  ( $j = 0, 1$ ),  $\rho(F)\} < 1$ , and let  $\varphi(z) \not\equiv 0$  be an entire function of finite order. Then every solution  $f \not\equiv 0$  of equation (1.5) satisfies*

$$(1.6) \quad \bar{\lambda}(f - \varphi) = \rho(f) = \infty, \quad \bar{\lambda}_2(f - \varphi) = \rho_2(f) \leq 1.$$

Furthermore, if  $F \not\equiv 0$ , then every solution  $f$  of equation (1.5) satisfies

$$(1.7) \quad \lambda(f) = \bar{\lambda}(f) = \bar{\lambda}(f - \varphi) = \rho(f) = \infty$$

and

$$(1.8) \quad \lambda_2(f) = \bar{\lambda}_2(f) = \bar{\lambda}_2(f - \varphi) = \rho_2(f) \leq 1.$$

**Remark 1.2.** The proof of Theorem 1.1 in which every solution  $f$  of equation (1.5) has infinite order is quite different from that of Theorem D (see [18]). The main ingredient in the proof is Lemma 2.9.

**Remark 1.3.** In [18], J.H.Y. Xu and T.B. Cao studied equation (1.5) and obtained the same result as in Theorem 1.1 but under the restriction that the complex constants  $a, b$  satisfy  $ab \neq 0$  and  $ab < 0$  and  $A_j(z)$  ( $j = 2, \dots, k - 1$ ) are polynomials of degree not greater than  $n - 1$  in  $z$ .

Setting  $D_j \equiv 0$  ( $j = 0, 1$ ) in Theorem 1.1, we obtain the following corollary.

**Corollary 1.1.** *Let  $a, b$  be complex numbers that satisfy  $ab(a - b) \neq 0$ . Suppose that  $A_j(z)$  ( $j = 0, 1, \dots, k - 1$ ),  $A_j(z) \not\equiv 0$  ( $j = 0, 1$ ) and  $F(z)$  are entire functions with  $\max\{\rho(A_j)$  ( $j = 0, 1, \dots, k - 1$ ),  $\rho(F)\} < 1$ , and let  $\varphi(z) \not\equiv 0$  be an entire function of finite order. Then every solution  $f \not\equiv 0$  of the equation*

$$(1.9) \quad f^{(k)} + A_{k-1}f^{(k-1)} + \dots + A_2f'' + A_1(z)e^{az}f' + A_0(z)e^{bz}f = F \quad (k \geq 2)$$

satisfies (1.6). Furthermore, if  $F \not\equiv 0$ , then every solution  $f$  of equation (1.9) satisfies (1.7) and (1.8).

**Remark 1.4.** If  $\rho(F) \geq 1$ , then equation (1.5) can possess solution of a finite order. For instance the equation

$$f''' - f'' + (e^{-z} - 1)f' + e^z f = e^z$$

satisfies  $\rho(F) = \rho(e^z) = 1$  and has a finite order solution  $f(z) = 1$ .

**Theorem 1.2.** Let  $a, b, A_j(z)$  ( $j = 0, 1, \dots, k-1$ ),  $D_j(z)$  ( $j = 0, 1$ ), and  $\varphi(z)$  satisfy the additional hypotheses of Theorem 1.1, and let  $F(z)$  be an entire function such that  $\varrho(F) \geq 1$ . Then every solution  $f$  of equation (1.5) satisfies (1.7) and (1.8) with at most one finite order solution  $f_0$ . For the exceptional solution  $f_0$ , if  $\varrho(F) > 1$ , then  $\varrho(f_0) = \varrho(F)$  and if  $\varrho(F) = 1$ , then  $\varrho(f_0) \leq 1$ .

**Corollary 1.2.** Let  $a, b, A_j(z)$  ( $j = 0, 1$ ),  $D_j(z)$  ( $j = 0, 1$ ), and  $\varphi(z)$  satisfy the additional hypotheses of Theorem 1.1, and let  $F(z)$  be an entire function. Then the following statements hold:

(i) If  $\varrho(F) < 1$ , then every solution  $f \not\equiv 0$  of the equation

$$(1.10) \quad f'' + (D_1(z) + A_1(z)e^{az})f' + (D_0(z) + A_0(z)e^{bz})f = F \quad (k \geq 2)$$

has infinite order and satisfies (1.6). Furthermore, if  $F \not\equiv 0$ , then every solution  $f$  of equation (1.10) satisfies (1.7) and (1.8).

- (ii) If  $\varrho(F) = 1$ , then every solution  $f$  of equation (1.10) has infinite order and satisfies (1.7) and (1.8), with at most one finite order solution  $f_0$  satisfying  $\varrho(f_0) \leq 1$ .
- (iii) If  $\varrho(F) > 1$ , then every solution  $f$  of equation (1.10) has infinite order and satisfies (1.7) and (1.8), with at most one finite order solution  $f_0$  satisfying  $\varrho(f_0) = \varrho(F)$ .

## 2. PRELIMINARY LEMMAS

Our proofs depend mainly upon the following lemmas. Before starting these lemmas, we recall the concept of the logarithmic density of subsets of  $(1, \infty)$ . For  $E \subset (1, \infty)$ , we define the logarithmic measure of a set  $E$  by  $\text{lm}(E) = \int_1^\infty \chi_E(t)/t \, dt$ , where  $\chi_E$  is the characteristic function of  $E$ . The upper logarithmic density and the lower logarithmic density of  $E$  are defined by

$$\overline{\log \text{dens}}(E) = \limsup_{r \rightarrow \infty} \frac{\text{lm}(E \cap [1, r])}{\log r}$$

and

$$\underline{\log \text{dens}}(E) = \liminf_{r \rightarrow \infty} \frac{\text{lm}(E \cap [1, r])}{\log r}.$$

**Lemma 2.1** ([10]). Let  $f$  be a transcendental meromorphic function of finite order  $\varrho$ , let  $\Gamma = \{(k_1, j_1), (k_2, j_2), \dots, (k_m, j_m)\}$  denote a finite set of distinct pairs of integers that satisfy  $k_i > j_i \geq 0$  for  $i = 1, \dots, m$  and let  $\varepsilon > 0$  be a given constant. Then there exists a set  $E_1 \subset [0, 2\pi)$  that has linear measure zero, such that if  $\psi \in [0, 2\pi) - E_1$ , then there is a constant  $R_1 = R_1(\psi) > 1$  such that for all  $z$  satisfying  $\arg z = \psi$  and  $|z| \geq R_1$  and for all  $(k, j) \in \Gamma$  we have

$$(2.1) \quad \left| \frac{f^{(k)}(z)}{f^{(j)}(z)} \right| \leq |z|^{(k-j)(\varrho-1+\varepsilon)}.$$

The next lemma describing the behavior of  $e^{P(z)}$ , where  $P(z)$  is a linear polynomial, is a special case of a more general result in [14, p. 254].

**Lemma 2.2** ([14]). Let  $P(z) = (\alpha + i\beta)z$ ,  $(\alpha + i\beta \neq 0)$ , and let  $A(z) (\neq 0)$  be a meromorphic function with  $\varrho(A) < 1$ . Set  $f(z) = A(z)e^{P(z)}$ ,  $z = re^{i\theta}$ ,  $\delta(P, \theta) = \alpha \cos \theta - \beta \sin \theta$ . Then for any given  $\varepsilon > 0$  there exists a set  $E_2 \subset [0, 2\pi)$  that has linear measure zero such that if  $\theta \in [0, 2\pi) \setminus (E_2 \cup E_3)$ , where  $E_3 = \{\theta \in [0, 2\pi) : \delta(P, \theta) = 0\}$  is a finite set, then for sufficiently large  $|z| = r$ , we have

(i) If  $\delta(P, \theta) > 0$ , then

$$(2.2) \quad \exp\{(1 - \varepsilon)\delta(P, \theta)r\} \leq |f(z)| \leq \exp\{(1 + \varepsilon)\delta(P, \theta)r\}.$$

(ii) If  $\delta(P, \theta) < 0$ , then

$$(2.3) \quad \exp\{(1 + \varepsilon)\delta(P, \theta)r\} \leq |f(z)| \leq \exp\{(1 - \varepsilon)\delta(P, \theta)r\}.$$

**Lemma 2.3** ([4]). Let  $A_0, A_1, \dots, A_{k-1}, F \neq 0$  be finite order meromorphic functions. If  $f$  is a meromorphic solution with  $\varrho(f) = \infty$  of the equation

$$(2.4) \quad f^{(k)} + A_{k-1}f^{(k-1)} + \dots + A_1f' + A_0f = F,$$

then  $\bar{\lambda}(f) = \lambda(f) = \varrho(f) = \infty$ .

**Lemma 2.4** ([2]). Let  $A_0, A_1, \dots, A_{k-1}, F \neq 0$  be finite order meromorphic functions. If  $f$  is a meromorphic solution with  $\varrho(f) = \infty$  and  $\varrho_2(f) = \varrho$  of equation (2.4), then  $\bar{\lambda}(f) = \lambda(f) = \varrho(f) = \infty$  and  $\bar{\lambda}_2(f) = \lambda_2(f) = \varrho_2(f) = \varrho$ .

**Lemma 2.5** ([3]). *Let  $a$  and  $b$  be complex numbers,  $ab \neq 0$  such that  $\arg a \neq \arg b$  or  $a = cb$  ( $0 < c < 1$ ). We denote index sets by*

$$\begin{aligned}\Lambda_1 &= \{0, a\}, \\ \Lambda_2 &= \{0, a, b, 2a, a + b\}.\end{aligned}$$

- (i) *If  $H_j$  ( $j \in \Lambda_1$ ) and  $H_b \neq 0$  are meromorphic functions of orders that are less than 1, setting  $\Psi_1(z) = \sum_{j \in \Lambda_1} H_j(z)e^{jz}$ , then  $\Psi_1(z) + H_b e^{bz} \neq 0$ .*
- (ii) *If  $H_j$  ( $j \in \Lambda_2$ ) and  $H_{2b} \neq 0$  are meromorphic functions of orders that are less than 1, setting  $\Psi_2(z) = \sum_{j \in \Lambda_2} H_j(z)e^{jz}$ , then  $\Psi_2(z) + H_{2b} e^{2bz} \neq 0$ .*

By interchanging  $a$  and  $b$  in Lemma 2.5, we easily obtain the following lemma.

**Lemma 2.6.** *Let  $a$  and  $b$  be complex numbers,  $ab \neq 0$ , such that  $a = cb$  ( $c > 1$ ). We denote the index set by*

$$\Lambda_3 = \{0, b\}.$$

*If  $H_j$  ( $j \in \Lambda_3$ ) and  $H_a \neq 0$  are meromorphic functions of orders that are less than 1, setting  $\Psi_3(z) = \sum_{j \in \Lambda_3} H_j(z)e^{jz}$ , then  $\Psi_3(z) + H_a e^{az} \neq 0$ .*

**Lemma 2.7** ([6]). *Let  $f(z)$  be a transcendental entire function. Then there is a set  $E_4 \subset (1, \infty)$  that has finite logarithmic measure, such that for all  $z$  with  $|z| = r \notin [0, 1] \cup E_4$  at which  $|f(z)| = M(r, f)$ , we have*

$$(2.5) \quad \left| \frac{f(z)}{f^{(s)}(z)} \right| \leq 2r^s \quad (s \in \mathbb{N}).$$

**Lemma 2.8** ([17]). *Let  $f(z)$  and  $g(z)$  be two nonconstant entire functions with  $\varrho(g) < \varrho(f) < \infty$ . Given  $\varepsilon$  with  $0 < 4\varepsilon < \varrho(f) - \varrho(g)$  and  $0 < \delta < \frac{1}{4}$ , there exists a set  $E_5$  with  $\overline{\log \text{dens}}(E_5) > 0$  such that*

$$(2.6) \quad \left| \frac{g(z)}{f(z)} \right| \leq \exp\{-r^{\varrho(f)-2\varepsilon}\}$$

*for all  $z$  such that  $|z| = r \in E_5$  is sufficiently large and that  $|f(z)| \geq M(r, f)\nu_f(r)^{\delta-\frac{1}{4}}$ .*



**Lemma 2.9.** Let  $a$  and  $b$  be complex numbers that satisfy  $ab(a-b) \neq 0$ . Suppose that  $A_j(z)$  ( $j = 0, 1, \dots, k-1$ ),  $A_j(z) \not\equiv 0$  ( $j = 0, 1$ ) and  $D_j(z)$  ( $j = 0, 1$ ) are entire functions with  $\max\{\varrho(A_j) \ (j = 0, 1, \dots, k-1), \varrho(D_j) \ (j = 0, 1)\} < 1$ . We denote

$$(2.7) \quad L_f = f^{(k)} + A_{k-1}f^{(k-1)} + \dots + A_2f'' \\ + (D_1(z) + A_1(z)e^{az})f' + (D_0(z) + A_0(z)e^{bz})f.$$

If  $f \not\equiv 0$  is a finite order entire function, then we have

$$\varrho(L_f) = \max\{1, \varrho(f)\}.$$

*Proof.* Let  $f \not\equiv 0$  be a finite order entire function. First, if  $f(z) \equiv C \neq 0$ , then

$$L_f = (D_0(z) + A_0(z)e^{bz})C.$$

Hence,  $\varrho(L_f) = 1$  and Lemma 2.9 holds.

We suppose  $f \not\equiv C$ . Then, by (2.7), we have  $\varrho(L_f) \leq \max\{1, \varrho(f)\}$ .

- (i) If  $\varrho(f) = \varrho < 1$ , then  $\varrho(L_f) \leq 1$ . Suppose that  $\varrho(L_f) < 1$ . By (2.7), we easily obtain a contradiction by Lemma 2.5 (i) or Lemma 2.6. Thus  $\varrho(L_f) = 1$ .
- (ii) If  $\varrho(f) = \varrho \geq 1$ , then  $\varrho(L_f) \leq \varrho(f)$ . Suppose that  $\varrho(L_f) < \varrho(f)$ . We can rewrite (2.7) as

$$(2.8) \quad \frac{L_f}{f} = \frac{f^{(k)}}{f} + A_{k-1} \frac{f^{(k-1)}}{f} + \dots + A_2 \frac{f''}{f} \\ + (D_1(z) + A_1(z)e^{az}) \frac{f'}{f} + D_0(z) + A_0(z)e^{bz}.$$

We divide the proof in three cases.

*Case 1.* Suppose first that  $\arg a \neq \arg b$ . Set

$$\max\{\varrho(A_j) \ (j = 0, 1, \dots, k-1), \varrho(D_j) \ (j = 0, 1)\} = \beta < 1.$$

Then, for any given  $\varepsilon$  ( $0 < \varepsilon < \min(1 - \beta, \frac{1}{4}(\varrho(f) - \varrho(L_f)))$ ), we have for sufficiently large  $r$

$$(2.9) \quad |D_j(z)| \leq \exp\{r^{\beta+\varepsilon}\} \quad (j = 0, 1), \\ |A_j(z)| \leq \exp\{r^{\beta+\varepsilon}\} \quad (j = 0, 1, \dots, k-1).$$

By Lemma 2.8, we know that there exists a set  $E_5$  with  $\overline{\log \text{dens}}(E_5) > 0$  such that

$$(2.10) \quad \left| \frac{L_f}{f} \right| \leq \exp\{-r^{\varrho(f)-2\varepsilon}\} \leq 1$$

for all  $z$  such that  $|z| = r \in E_5$  is sufficiently large and that  $|f(z)| \geq M(r, f)\nu_f(r)^{\delta - \frac{1}{4}}$ . Also, by Lemma 2.1, for the above  $\varepsilon$  there exists a set  $E_1 \subset [0, 2\pi)$  that has linear measure zero, such that if  $\theta \in [0, 2\pi) - E_1$ , then there is a constant  $R_1 = R_1(\theta) > 1$  such that for all  $z$  satisfying  $\arg z = \theta$  and  $|z| \geq R_1$ , we have

$$(2.11) \quad \left| \frac{f^{(i)}(z)}{f(z)} \right| \leq |z|^{i(\varrho - 1 + \varepsilon)} \quad (i = 1, \dots, k).$$

By Lemma 2.2, there exists a ray  $\arg z = \theta \in [0, 2\pi) \setminus E_1 \cup E_2 \cup E_3$ ,  $E_3 = \{\theta \in [0, 2\pi) : \delta(az, \theta) = 0 \text{ or } \delta(bz, \theta) = 0\} \subset [0, 2\pi)$ ,  $E_1 \cup E_2$  having linear measure zero,  $E_3$  being a finite set, such that

$$\delta(az, \theta) < 0, \quad \delta(bz, \theta) > 0$$

and for any given  $\varepsilon$  ( $0 < \varepsilon < \min(1 - \beta, \frac{1}{4}(\varrho(f) - \varrho(L_f)))$ ), by (2.9), (2.11) we have for sufficiently large  $|z| = r$

$$(2.12) \quad |A_0 e^{bz}| \geq \exp\{(1 - \varepsilon)\delta(bz, \theta)r\},$$

$$(2.13) \quad \begin{aligned} & \left| \frac{f^{(k)}}{f} A_{k-1} \frac{f^{(k-1)}}{f} + \dots + A_2 \frac{f''}{f} + D_0(z) \right| \\ & \leq \left| \frac{f^{(k)}}{f} \right| + |A_{k-1}| \left| \frac{f^{(k-1)}}{f} \right| + \dots + |A_2| \left| \frac{f''}{f} \right| + |D_0(z)| \\ & \leq r^{k(\varrho - 1 + \varepsilon)} + r^{(k-1)(\varrho - 1 + \varepsilon)} \exp\{r^{\beta + \varepsilon}\} + \dots \\ & \quad + r^{2(\varrho - 1 + \varepsilon)} \exp\{r^{\beta + \varepsilon}\} + \exp\{r^{\beta + \varepsilon}\} \\ & \leq k r^{k(\varrho - 1 + \varepsilon)} \exp\{r^{\beta + \varepsilon}\}, \end{aligned}$$

$$(2.14) \quad \begin{aligned} & \left| (D_1(z) + A_1(z)e^{az}) \frac{f'}{f} \right| \\ & \leq r^{\varrho - 1 + \varepsilon} (\exp\{(1 - \varepsilon)\delta(az, \theta)r\} + \exp\{r^{\beta + \varepsilon}\}) \\ & \leq r^{\varrho - 1 + \varepsilon} (1 + \exp\{r^{\beta + \varepsilon}\}). \end{aligned}$$

By (2.8), (2.10), and (2.12)–(2.14) we have

$$\exp\{(1 - \varepsilon)\delta(bz, \theta)r\} \leq |A_0 e^{bz}| \leq K r^{k(\varrho - 1 + \varepsilon)} \exp\{r^{\beta + \varepsilon}\},$$

where  $K > 0$  is a real constant. This is a contradiction by  $\beta + \varepsilon < 1$ . Hence,  $\varrho(L_f) = \varrho(f)$ .

*Case 2.* Suppose now  $a = cb$  ( $0 < c < 1$ ). Then for any ray  $\arg z = \theta$  we have

$$\delta(az, \theta) = c\delta(bz, \theta).$$

Then, by Lemma 2.2, for any given  $\varepsilon$  ( $0 < \varepsilon < \min((1-c)/(2(1+c)), 1-\beta, \frac{1}{4}(\varrho(f) - \varrho(L_f)))$ ) there exist  $E_j \subset [0, 2\pi)$  ( $j = 1, 2, 3$ ) such that  $E_1, E_2$  have linear measure zero and  $E_3$  is a finite set, where  $E_1, E_2$ , and  $E_3$  are defined as in Case 1, respectively. We take the ray  $\arg z = \theta \in [0, 2\pi) \setminus E_1 \cup E_2 \cup E_3$  such that  $\delta(bz, \theta) > 0$  and for sufficiently large  $|z| = r$ , we have (2.12), (2.13), and

$$(2.15) \quad \left| (D_1(z) + A_1(z)e^{az}) \frac{f'}{f} \right| \leq r^{\varrho-1+\varepsilon} (\exp\{r^{\beta+\varepsilon}\} + \exp\{(1+\varepsilon)c\delta(bz, \theta)r\}).$$

Thus by (2.8), (2.10), (2.12), (2.13), and (2.15) we obtain

$$(2.16) \quad \begin{aligned} \exp\{(1-\varepsilon)\delta(bz, \theta)r\} &\leq |A_0 e^{bz}| \\ &\leq k r^{k(\varrho-1+\varepsilon)} \exp\{r^{\beta+\varepsilon}\} \\ &\quad + r^{\varrho-1+\varepsilon} (\exp\{r^{\beta+\varepsilon}\} + \exp\{(1+\varepsilon)c\delta(bz, \theta)r\}) + 1 \\ &\leq (k+1) r^{k(\varrho-1+\varepsilon)} \exp\{r^{\beta+\varepsilon}\} \\ &\quad + r^{\varrho-1+\varepsilon} \exp\{(1+\varepsilon)c\delta(bz, \theta)r\} + 1. \end{aligned}$$

For  $\varepsilon$  ( $0 < \varepsilon < \min((1-c)/(2(1+c)), 1-\beta, \frac{1}{4}(\varrho(f) - \varrho(L_f)))$ ), we have as  $r \rightarrow \infty$

$$(2.17) \quad \frac{(k+1) r^{k(\varrho-1+\varepsilon)} \exp\{r^{\beta+\varepsilon}\}}{\exp\{(1-\varepsilon)\delta(bz, \theta)r\}} \rightarrow 0,$$

$$(2.18) \quad \frac{r^{\varrho-1+\varepsilon} \exp\{(1+\varepsilon)c\delta(bz, \theta)r\}}{\exp\{(1-\varepsilon)\delta(bz, \theta)r\}} \rightarrow 0,$$

$$(2.19) \quad \frac{1}{\exp\{(1-\varepsilon)\delta(bz, \theta)r\}} \rightarrow 0.$$

By (2.16)–(2.19), we get  $1 \leq 0$ . This is a contradiction. Hence,  $\varrho(L_f) = \varrho(f)$ .

*Case 3.* Finally, we suppose  $a = cb$  ( $c > 1$ ). We can rewrite (2.7) as

$$(2.20) \quad \begin{aligned} \frac{L_f}{f} \frac{f}{f'} &= \frac{f^{(k)}}{f'} + A_{k-1} \frac{f^{(k-1)}}{f'} + \dots + A_2 \frac{f''}{f'} \\ &\quad + (D_0(z) + A_0(z)e^{bz}) \frac{f}{f'} + D_1(z) + A_1(z)e^{az}. \end{aligned}$$

By Lemma 2.7, there is a set  $E_4 \subset (1, \infty)$  that has finite logarithmic measure such that for all  $z$  with  $|z| = r \notin [0, 1] \cup E_4$  at which  $|f(z)| = M(r, f)$  we have

$$(2.21) \quad \left| \frac{f(z)}{f'(z)} \right| \leq 2r.$$

By Lemma 2.8, for  $\varepsilon$  ( $0 < \varepsilon < \min((c-1)/(2(c+1)), 1-\beta, \frac{1}{4}(\varrho(f) - \varrho(L_f)))$ ), we know that there exists a set  $E_5$  with  $\log \text{dens}(E_5) > 0$  such that

$$(2.22) \quad \left| \frac{L_f}{f} \right| \leq \exp\{-r^{\varrho(f)-2\varepsilon}\} \leq 1$$

for all  $z$  such that  $|z| = r \in E_5$  is sufficiently large and that  $|f(z)| \geq M(r, f)\nu_f(r)^{\delta - \frac{1}{4}}$ . Since  $E_4 \subset (1, \infty)$  has finite logarithmic measure and  $E_5$  satisfies  $\overline{\log \text{dens}}(E_5) > 0$ , we have  $\overline{\log \text{dens}}(E_5 - ([0, 1] \cup E_4)) > 0$ . By (2.21) and (2.22) for sufficiently large  $|z| = r$  we get

$$(2.23) \quad \left| \frac{L_f}{f'} \right| = \left| \frac{L_f}{f} \frac{f}{f'} \right| \leq 2r \exp\{-r^{\varrho(f)-2\varepsilon}\} \leq 2r.$$

For any ray  $\arg z = \theta$ , we have

$$\delta(az, \theta) = c\delta(bz, \theta).$$

By Lemma 2.2, there exists a ray  $\arg z = \theta \in [0, 2\pi) \setminus E_1 \cup E_2 \cup E_3$ ,  $E_3 = \{\theta \in [0, 2\pi) : \delta(az, \theta) = 0 \text{ or } \delta(bz, \theta) = 0\} \subset [0, 2\pi)$ ,  $E_1 \cup E_2$  having linear measure zero,  $E_3$  being a finite set, such that

$$\delta(az, \theta) = c\delta(bz, \theta) > 0$$

and by (2.9), (2.11), and (2.21), for sufficiently large  $|z| = r$  we have

$$(2.24) \quad |A_1 e^{az}| \geq \exp\{(1 - \varepsilon)c\delta(bz, \theta)r\},$$

$$(2.25) \quad \left| (D_0(z) + A_0(z)e^{bz}) \frac{f}{f'} \right| \leq 2r \exp\{r^{\beta+\varepsilon}\} + 2r \exp\{(1 + \varepsilon)\delta(bz, \theta)r\},$$

$$(2.26) \quad \left| \frac{f^{(k)}}{f'} + A_{k-1} \frac{f^{(k-1)}}{f'} + \dots + A_2 \frac{f''}{f'} + D_1 \right| \\ \leq \left| \frac{f(z)}{f'(z)} \right| \left( \left| \frac{f^{(k)}}{f} \right| + |A_{k-1}| \left| \frac{f^{(k-1)}}{f} \right| + \dots + |A_2| \left| \frac{f''}{f} \right| \right) + |D_1| \\ \leq 2r(k-1)r^{k(\varrho-1+\varepsilon)} \exp\{r^{\beta+\varepsilon}\} + \exp\{r^{\beta+\varepsilon}\} \\ \leq 2kr^{k(\varrho-1+\varepsilon)+1} \exp\{r^{\beta+\varepsilon}\}.$$

By (2.20), (2.23), and (2.24)–(2.26), we have

$$(2.27) \quad \exp\{(1 - \varepsilon)c\delta(bz, \theta)r\} \leq |A_1 e^{az}| \\ \leq 2kr^{k(\varrho-1+\varepsilon)+1} \exp\{r^{\beta+\varepsilon}\} \\ + 2r \exp\{r^{\beta+\varepsilon}\} + 2r \exp\{(1 + \varepsilon)\delta(bz, \theta)r\} + 2r \\ \leq 2(k+1)r^{k(\varrho-1+\varepsilon)+1} \exp\{r^{\beta+\varepsilon}\} \\ + 2r \exp\{(1 + \varepsilon)\delta(bz, \theta)r\} + 2r.$$

For  $\varepsilon$  ( $0 < \varepsilon < \min((c-1)/(2(c+1)), 1 - \beta, \frac{1}{4}(\varrho(f) - \varrho(L_f)))$ ), we have as  $r \rightarrow \infty$

$$(2.28) \quad \frac{2(k+1)r^{k(\varrho-1+\varepsilon)+1} \exp\{r^{\beta+\varepsilon}\}}{\exp\{(1-\varepsilon)c\delta(bz, \theta)r\}} \rightarrow 0,$$

$$(2.29) \quad \frac{2r \exp\{(1+\varepsilon)\delta(bz, \theta)r\}}{\exp\{(1-\varepsilon)c\delta(bz, \theta)r\}} \rightarrow 0,$$

$$(2.30) \quad \frac{2r}{\exp\{(1-\varepsilon)c\delta(bz, \theta)r\}} \rightarrow 0.$$

By (2.27)–(2.30), we get  $1 \leq 0$ . This is a contradiction. Hence,  $\varrho(L_f) = \varrho(f)$ .  $\square$

By using the Wiman-Valiron theory [12], we easily obtain the following result of which we omit the proof.

**Lemma 2.10.** *Let  $A_0(z), \dots, A_{k-1}(z), F(z)$  be entire functions of finite order. If  $f$  is a solution of the equation*

$$(2.31) \quad f^{(k)} + A_{k-1}(z)f^{(k-1)} + \dots + A_1(z)f' + A_0(z)f = F,$$

then  $\varrho_2(f) \leq \max\{\varrho(A_0), \dots, \varrho(A_{k-1}), \varrho(F)\}$ .

### 3. PROOF of Theorem 1.1

Assume that  $f \not\equiv 0$  is a solution of equation (1.5). We prove that  $f$  is of infinite order. We suppose the contrary:  $\varrho(f) < \infty$ . By Lemma 2.9, we have  $1 \leq \varrho(L_f) = \varrho(F) < 1$  and this is a contradiction. Hence, every solution of equation (1.5) is of infinite order and by Lemma 2.10 we have  $\varrho_2(f) \leq 1$ . Suppose that  $\varphi(z) \not\equiv 0$  is an entire function of finite order. Set  $g = f - \varphi$ , then  $f = g + \varphi$  and due to  $\varrho(\varphi) < \infty$  we have  $\varrho(f) = \varrho(g) = \infty$ ,  $\varrho_2(f) = \varrho_2(g) \leq 1$ . Thus,  $g$  is a solution of the equation

$$g^{(k)} + A_{k-1}g^{(k-1)} + \dots + A_2g'' + (D_1 + A_1e^{az})g' + (D_0 + A_0e^{bz})g = H,$$

where

$$H = F - (\varphi^{(k)} + A_{k-1}\varphi^{(k-1)} + \dots + A_2\varphi'' + (D_1 + A_1e^{az})\varphi' + (D_0 + A_0e^{bz})\varphi).$$

By  $\varphi(z) \not\equiv 0$  and  $\varrho(\varphi) < \infty$  we have  $H \not\equiv 0$ . Since  $\varrho(H) < \infty$ , by Lemma 2.3 and Lemma 2.4 we get

$$\bar{\lambda}(f - \varphi) = \varrho(f - \varphi) = \varrho(f) = \infty, \quad \bar{\lambda}_2(f - \varphi) = \varrho_2(f - \varphi) = \varrho_2(f) \leq 1.$$

Furthermore, if  $F \neq 0$ , then by the fact that  $f$  is an infinite order solution of equation (1.5), and due to Lemma 2.3 and Lemma 2.4 we have

$$\begin{aligned}\lambda(f) &= \bar{\lambda}(f) = \bar{\lambda}(f - \varphi) = \varrho(f) = \infty, \\ \lambda_2(f) &= \bar{\lambda}_2(f) = \bar{\lambda}_2(f - \varphi) = \varrho_2(f) \leq 1.\end{aligned}$$

□

#### 4. PROOF of Theorem 1.2

Assume that  $f_0$  is a solution of (1.5) with  $\varrho(f_0) = \varrho < \infty$ . If  $f_1$  is another finite order solution of (1.5), then  $\varrho(f_1 - f_0) < \infty$ , and  $f_1 - f_0$  is a solution of the corresponding homogeneous equation of (1.5), but  $\varrho(f_1 - f_0) = \infty$  by virtue of Theorem 1.1, which is a contradiction. Hence, (1.5) has at most one finite order solution  $f_0$  and all other solutions  $f_1$  of (1.5) are of infinite order and satisfy (1.7) and (1.8). If  $\varrho(F) > 1$ , suppose there exists a solution  $f_0$  of (1.5) with  $\varrho(f_0) < \infty$ . Then, we have  $\varrho(f_0) > 1$  and by Lemma 2.9 we get  $\varrho(L_f) = \varrho(f_0) = \varrho(F)$ . Suppose that  $\varrho(F) = 1$ . If there exists a solution  $f_0$  of (1.5) with  $\varrho(f_0) < \infty$ , then  $\varrho(f_0) \leq 1$ . Indeed, if we suppose that  $\varrho(f_0) > 1$ , then by Lemma 2.9 we get  $\varrho(L_f) = \varrho(f_0) = \varrho(F) > 1$  and this is a contradiction. □

#### 5. PROOF of Corollary 1.2

By using Theorem 1.1 and Theorem 1.2, we obtain Corollary 1.2. □

**Acknowledgment.** The authors would like to thank the referee for his/her helpful remarks and suggestions improving the paper.

#### References

- [1] *I. Amemiya, M. Ozawa*: Non-existence of finite order solutions of  $w'' + e^{-z}w' + Q(z)w = 0$ . *Hokkaido Math. J.* 10 (1981), 1–17.
- [2] *B. Belaïdi*: Growth and oscillation theory of solutions of some linear differential equations. *Mat. Vesn.* 60 (2008), 233–246.
- [3] *B. Belaïdi, A. El Farissi*: Differential polynomials generated by some complex linear differential equations with meromorphic coefficients. *Glas. Mat., Ser. III* 43 (2008), 363–373.
- [4] *Z. X. Chen*: Zeros of meromorphic solutions of higher order linear differential equations. *Analysis* 14 (1994), 425–438.
- [5] *Z. X. Chen*: The growth of solutions of  $f'' + e^{-z}f' + Q(z)f = 0$  where the order  $(Q) = 1$ . *Sci. China, Ser. A* 45 (2002), 290–300.

- [6] *Z. X. Chen, K. H. Shon*: On the growth of solutions of a class of higher order differential equations. *Acta Math. Sci., Ser. B, Engl. Ed.* *24* (2004), 52–60.
- [7] *M. Frei*: Über die Lösungen linearer Differentialgleichungen mit ganzen Funktionen als Koeffizienten. *Comment. Math. Helv.* *35* (1961), 201–222.
- [8] *M. Frei*: Über die subnormalen Lösungen der Differentialgleichung  $w'' + e^{-z}w' + \text{konst.}w = 0$ . *Comment. Math. Helv.* *36* (1961), 1–8.
- [9] *G. G. Gundersen*: On the question of whether  $f'' + e^{-z}f' + B(z)f = 0$  can admit a solution  $f \not\equiv 0$  of finite order. *Proc. R. Soc. Edinb., Sect. A* *102* (1986), 9–17.
- [10] *G. G. Gundersen*: Estimates for the logarithmic derivative of a meromorphic function, plus similar estimates. *J. Lond. Math. Soc., II. Ser.* *37* (1988), 88–104.
- [11] *W. K. Hayman*: Meromorphic functions. Clarendon Press, Oxford, 1964.
- [12] *G. Jank, L. Volkmann*: Einführung in die Theorie der ganzen und meromorphen Funktionen mit Anwendungen auf Differentialgleichungen. Birkhäuser, Basel, 1985.
- [13] *J. K. Langley*: On complex oscillation and a problem of Ozawa. *Kodai Math. J.* *9* (1986), 430–439.
- [14] *A. I. Markushevich*: Theory of functions of a complex variable, Vol. II. Prentice-Hall, Englewood Cliffs, 1965.
- [15] *R. Nevanlinna*: Eindeutige analytische Funktionen, Zweite Auflage. Reprint. Die Grundlehren der mathematischen Wissenschaften, Band 46. Springer, Berlin-Heidelberg-New York, 1974. (In German.)
- [16] *M. Ozawa*: On a solution of  $w'' + e^{-z}w' + (az + b)w = 0$ . *Kodai Math. J.* *3* (1980), 295–309.
- [17] *J. Wang, I. Laine*: Growth of solutions of second order linear differential equations. *J. Math. Anal. Appl.* *342* (2008), 39–51.
- [18] *H. Y. Xu, T. B. Cao*: Oscillation of solutions of some higher order linear differential equations. *Electron. J. Qual. Theory Differ. Equ.*, paper No. 63, 18 pages (2009).

*Authors' address: A. El Farissi, B. Belaidi, Department of Mathematics, Laboratory of Pure and Applied Mathematics, University of Mostaganem (UMAB), B. P. 227 Mostaganem, Algeria, e-mail: elfarissi.abdallah@yahoo.fr; belaidibenharrat@yahoo.fr, belaidi@univ-mosta.dz.*