

Sulkhan Mukhigulashvili

The Dirichlet boundary value problems for strongly singular higher-order nonlinear functional-differential equations

*Czechoslovak Mathematical Journal*, Vol. 63 (2013), No. 1, 235–263

Persistent URL: <http://dml.cz/dmlcz/143182>

## Terms of use:

© Institute of Mathematics AS CR, 2013

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE DIRICHLET BOUNDARY VALUE PROBLEMS FOR  
STRONGLY SINGULAR HIGHER-ORDER NONLINEAR  
FUNCTIONAL-DIFFERENTIAL EQUATIONS

SULKHAN MUKHIGULASHVILI, Brno

(Received December 19, 2011)

*Abstract.* The a priori boundedness principle is proved for the Dirichlet boundary value problems for strongly singular higher-order nonlinear functional-differential equations. Several sufficient conditions of solvability of the Dirichlet problem under consideration are derived from the a priori boundedness principle. The proof of the a priori boundedness principle is based on the Agarwal-Kiguradze type theorems, which guarantee the existence of the Fredholm property for strongly singular higher-order linear differential equations with argument deviations under the two-point conjugate and right-focal boundary conditions.

*Keywords:* higher order functional-differential equation, Dirichlet boundary value problem, strong singularity, Fredholm property, a priori boundedness principle

*MSC 2010:* 34K06, 34K10

## 1. STATEMENT OF THE MAIN RESULTS

**1.1. Statement of the problem and a survey of the literature.** Consider the functional differential equation

$$(1.1) \quad u^{(n)}(t) = F(u)(t)$$

with the two-point boundary conditions

$$(1.2) \quad u^{(i-1)}(a) = 0 \quad (i = 1, \dots, m), \quad u^{(i-1)}(b) = 0 \quad (i = 1, \dots, n - m).$$

---

This work is supported by the Academy of Sciences of the Czech Republic (Institutional Research Plan # AV0Z10190503) and by the Shota Rustaveli National Science Foundation (Project # GNSF/ST09\_175\_3-101).

Here  $n \geq 2$ ,  $m$  is the integer part of  $n/2$ ,  $-\infty < a < b < +\infty$ , and the operator  $F$  is acting from the set of  $(m-1)$ -th time continuously differentiable on  $]a, b[$  functions to the set  $L_{\text{loc}}(]a, b[)$ . By  $u^{(j-1)}(a)$  ( $u^{(j-1)}(b)$ ) we denote the right (the left) limit of the function  $u^{(j-1)}$  at the point  $a(b)$ .

The problem is singular in the sense that for an arbitrary  $x$  the right-hand side of equation (1.41) may have nonintegrable singularities at the points  $a$  and  $b$ .

Throughout the paper we use the following notation:

- ▷  $\mathbb{R}^+ = [0, +\infty[$ ;
- ▷  $[x]_+$  the positive part of a number  $x$ , that is  $[x]_+ = \frac{1}{2}(x + |x|)$ ;
- ▷  $L_{\text{loc}}(]a, b[)$  ( $L_{\text{loc}}(]a, b[)$ ) is the space of functions  $y: ]a, b[ \rightarrow \mathbb{R}$ , which are integrable on  $[a + \varepsilon, b - \varepsilon]$  for arbitrarily small  $\varepsilon > 0$ ;
- ▷  $L_{\alpha, \beta}(]a, b[)$  ( $L_{\alpha, \beta}^2(]a, b[)$ ) is the space of integrable (square integrable) with the weight  $(t-a)^\alpha(b-t)^\beta$  functions  $y: ]a, b[ \rightarrow \mathbb{R}$ , with the norm

$$\|y\|_{L_{\alpha, \beta}} = \int_a^b (s-a)^\alpha(b-s)^\beta |y(s)| ds$$

$$\left( \|y\|_{L_{\alpha, \beta}^2} = \left( \int_a^b (s-a)^\alpha(b-s)^\beta y^2(s) ds \right)^{1/2} \right);$$

- ▷  $L([a, b]) = L_{0,0}(]a, b[)$ ,  $L^2([a, b]) = L_{0,0}^2(]a, b[)$ ;
- ▷  $M(]a, b[)$  is the set of measurable functions  $\tau: ]a, b[ \rightarrow ]a, b[$ ;
- ▷  $\tilde{L}_{\alpha, \beta}^2(]a, b[)$  ( $\tilde{L}_{\alpha, \beta}^2(]a, b[)$ ) is the Banach space of  $y \in L_{\text{loc}}(]a, b[)$  ( $L_{\text{loc}}(]a, b[)$ ) functions, with the norm

$$\|y\|_{\tilde{L}_{\alpha, \beta}^2} \equiv \max \left\{ \left[ \int_a^t (s-a)^\alpha \left( \int_s^t y(\xi) d\xi \right)^2 ds \right]^{1/2} : a \leq t \leq \frac{a+b}{2} \right\}$$

$$+ \max \left\{ \left[ \int_t^b (b-s)^\beta \left( \int_t^s y(\xi) d\xi \right)^2 ds \right]^{1/2} : \frac{a+b}{2} \leq t \leq b \right\} < +\infty.$$

- ▷  $L_n(]a, b[)$  is the Banach space of  $y \in L_{\text{loc}}(]a, b[)$  functions, with the norm

$$\|y\|_{\tilde{L}_{\alpha, \beta}^2} = \sup \left\{ [(s-a)(b-t)]^{m-1/2} \int_s^t (\xi-a)^{n-2m} |y(\xi)| d\xi : a < s \leq t < b \right\} < +\infty.$$

- ▷  $C_{\text{loc}}^{n-1}(]a, b[)$ , ( $\tilde{C}_{\text{loc}}^{n-1}(]a, b[)$ ) is the space of functions  $y: ]a, b[ \rightarrow \mathbb{R}$  which are continuous (absolutely continuous) together with  $y', y'', \dots, y^{(n-1)}$  on  $[a+\varepsilon, b-\varepsilon]$  for arbitrarily small  $\varepsilon > 0$ .
- ▷  $\tilde{C}^{n-1, m}(]a, b[)$  is the space of functions  $y \in \tilde{C}_{\text{loc}}^{n-1}(]a, b[)$ , such that

$$(1.3) \quad \int_a^b |x^{(m)}(s)|^2 ds < +\infty.$$

▷  $C_1^{m-1}(]a, b[)$  is the Banach space of functions  $y \in C_{\text{loc}}^{m-1}(]a, b[)$ , such that

$$(1.4) \quad \begin{aligned} \limsup_{t \rightarrow a} \frac{|x^{(i-1)}(t)|}{(t-a)^{m-i+1/2}} &< +\infty \quad (i = 1, \dots, m), \\ \limsup_{t \rightarrow b} \frac{|x^{(i-1)}(t)|}{(b-t)^{m-i+1/2}} &< +\infty \quad (i = 1, \dots, n-m), \end{aligned}$$

with the norm:

$$\|x\|_{C_1^{m-1}} = \sum_{i=1}^m \sup \left\{ \frac{|x^{(i-1)}(t)|}{\alpha_i(t)} : a < t < b \right\},$$

where  $\alpha_i(t) = (t-a)^{m-i+1/2}(b-t)^{m-i+1/2}$ .

▷  $\tilde{C}_1^{m-1}(]a, b[)$  is the Banach space of functions  $y \in \tilde{C}_{\text{loc}}^{m-1}(]a, b[)$ , such that conditions (1.3) and (1.4) hold, with the norm:

$$\|x\|_{\tilde{C}_1^{m-1}} = \sum_{i=1}^m \sup \left\{ \frac{|x^{(i-1)}(t)|}{\alpha_i(t)} : a < t < b \right\} + \left( \int_a^b |x^{(m)}(s)|^2 ds \right)^{1/2}.$$

▷  $D_n(]a, b[ \times \mathbb{R}^+)$  is the set of such functions  $\delta: ]a, b[ \times \mathbb{R}^+ \rightarrow L_n(]a, b[)$  that  $\delta(t, \cdot): \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is nondecreasing for every  $t \in ]a, b[$ , and  $\delta(\cdot, \varrho) \in L_n(]a, b[)$  for any  $\varrho \in \mathbb{R}^+$ .

▷  $D_{2n-2m-2, 2m-2}(]a, b[ \times \mathbb{R}^+)$  is the set of such functions  $\delta: ]a, b[ \times \mathbb{R}^+ \rightarrow \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  that  $\delta(t, \cdot): \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is nondecreasing for every  $t \in ]a, b[$ , and  $\delta(\cdot, \varrho) \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  for any  $\varrho \in \mathbb{R}^+$ .

▷ A solution of problem (1.1), (1.2) is sought in the space  $\tilde{C}^{n-1, m}(]a, b[)$ .

The singular ordinary differential and functional-differential equations have been studied with sufficient completeness under different boundary conditions, see for example [1], [3], [4], [5], [6], [7], [8], [9], [11], [12], [13], [14], [16], [21], [22], [23], [24], [25] and the references cited therein. But the equation (1.1), even under the boundary condition (1.2), have not been studied in the case when the operator  $F$  has the form

$$(1.5) \quad F(x)(t) = \sum_{j=1}^m p_j(t)x^{(j-1)}(\tau_j(t)) + f(x)(t),$$

where the singularity of the functions  $p_j: L_{\text{loc}}([a, b])$  is such that the inequalities

$$(1.6) \quad \begin{aligned} \int_a^b (s-a)^{n-1}(b-s)^{2m-1} [(-1)^{n-m} p_1(s)]_+ ds &< +\infty, \\ \int_a^b (s-a)^{n-j}(b-s)^{2m-j} |p_j(s)| ds &< +\infty \quad (j = 2, \dots, m), \end{aligned}$$

are not fulfilled (in this case we say that the linear part of the operator  $F$  is strongly singular), the operator  $f$  is continuously acting from  $C_1^{m-1}(]a, b[)$  to  $L\tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$ , and the inclusion

$$(1.7) \quad \sup\{f(x)(t) : \|x\|_{C_1^{m-1}} \leq \varrho\} \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$$

holds. The first step in studying the differential equations with strong singularities was made by R. P. Agarwal and I. Kiguradze in the article [2], where the linear ordinary differential equations under conditions (1.2), in the case when the functions  $p_j$  have strong singularities at the points  $a$  and  $b$ , are studied. Also the ordinary differential equations with strong singularities under two-point boundary conditions are studied in the articles of I. Kiguradze [10], [19], and N. Partsvania [20]. In the papers [18], [15] these results are generalized to linear differential equations with deviating arguments, i.e., the Agarwal-Kiguradze type theorems, which guarantee Fredholm's property for linear differential equations with deviating arguments are proved.

In this paper, on the bases of articles [2] and [17] we prove the a priori boundedness principle for the problem (1.1), (1.2) in the case when the operator has the form (1.5).

Now we introduce some results from the articles [18], [15], which we need for this work. Consider the equation

$$(1.8) \quad u^{(n)}(t) = \sum_{j=1}^m p_j(t) u^{(j-1)}(\tau_j(t)) + q(t) \quad \text{for } a < t < b.$$

For problem (1.8), (1.2) we assume, that when  $n = 2m$ , then the conditions

$$(1.9) \quad p_j \in L_{\text{loc}}(]a, b[) \quad (j = 1, \dots, m)$$

are fulfilled and when  $n = 2m + 1$ , along with (1.9), the condition

$$(1.10) \quad \limsup_{t \rightarrow b} \left| (b-t)^{2m-1} \int_{t_1}^t p_1(s) ds \right| < +\infty \quad \left( t_1 = \frac{a+b}{2} \right)$$

holds.

By  $h_j : ]a, b[ \times ]a, b[ \rightarrow \mathbb{R}_+$  and  $f_j : [a, b] \times M(]a, b[) \rightarrow C_{\text{loc}}(]a, b[ \times ]a, b[)$  ( $j = 1, \dots, m$ ) we denote the functions and operators, respectively, defined by the equalities

$$(1.11) \quad h_1(t, s) = \left| \int_s^t (\xi - a)^{n-2m} [(-1)^{n-m} p_1(\xi)]_+ d\xi \right|,$$

$$h_j(t, s) = \left| \int_s^t (\xi - a)^{n-2m} p_j(\xi) d\xi \right| \quad (j = 2, \dots, m),$$

and

$$(1.12) \quad f_j(c, \tau_j)(t, s) = \left| \int_s^t (\xi - a)^{n-2m} |p_j(\xi)| \left| \int_\xi^{\tau_j(\xi)} (\xi_1 - c)^{2(m-j)} d\xi_1 \right|^{1/2} d\xi \right|.$$

Let  $k = 2k_1 + 1$  ( $k_1 \in \mathbb{N}$ ), then we denote

$$k!! = \begin{cases} 1 & \text{for } k \leq 0, \\ 1 \cdot 3 \cdot 5 \cdot \dots \cdot k & \text{for } k \geq 1. \end{cases}$$

Now we can introduce the main theorem of the paper [18].

**Theorem 1.1.** *Let there exist numbers  $t^* \in ]a, b[$ ,  $l_{kj} > 0$ ,  $\bar{l}_{kj} \geq 0$ , and  $\gamma_{kj} > 0$  ( $k = 0, 1; j = 1, \dots, m$ ) such that along with*

$$(1.13) \quad B_0 \equiv \sum_{j=1}^m \left( \frac{(2m-j)2^{2m-j+1}l_{0j}}{(2m-1)!!(2m-2j+1)!!} + \frac{2^{2m-j-1}(t^* - a)^{\gamma_{0j}}\bar{l}_{0j}}{(2m-2j-1)!!(2m-3)!!\sqrt{2\gamma_{0j}}} \right) < \frac{1}{2},$$

$$(1.14) \quad B_1 \equiv \sum_{j=1}^m \left( \frac{(2m-j)2^{2m-j+1}l_{1j}}{(2m-1)!!(2m-2j+1)!!} + \frac{2^{2m-j-1}(b - t^*)^{\gamma_{1j}}\bar{l}_{1j}}{(2m-2j-1)!!(2m-3)!!\sqrt{2\gamma_{1j}}} \right) < \frac{1}{2},$$

the conditions

$$(1.15) \quad (t - a)^{2m-j}h_j(t, s) \leq l_{0j}, \quad (t - a)^{m-\gamma_{0j}-1/2}f_j(a, \tau_j)(t, s) \leq \bar{l}_{0j}$$

for  $a < t \leq s \leq t^*$ , and

$$(1.16) \quad (b - t)^{2m-j}h_j(t, s) \leq l_{1j}, \quad (b - t)^{m-\gamma_{1j}-1/2}f_j(b, \tau_j)(t, s) \leq \bar{l}_{1j}$$

for  $t^* \leq s \leq t < b$  hold. Then problem (1.8), (1.2) is uniquely solvable in the space  $\tilde{C}^{n-1,m}(]a, b[)$ .

Also, in [15] the following theorem is proved:

**Theorem 1.2.** *Let all the conditions of Theorem 1.1 be satisfied. Then the unique solution  $u$  of problem (1.8), (1.2) for every  $q \in \tilde{L}_{2n-2m-2, 2m-2}^2(\lceil a, b \rceil)$  admits the estimate*

$$(1.17) \quad \|u^{(m)}\|_{L^2} \leq r \|q\|_{\tilde{L}_{2n-2m-2, 2m-2}^2},$$

with

$$r = \frac{2^m(1+b-a)(2n-2m-1)}{(\nu_n - 2 \max\{B_0, B_1\})(2m-1)!!}, \quad \nu_{2m} = 1, \quad \nu_{2m+1} = \frac{2m+1}{2},$$

and thus the constant  $r > 0$  depends only on the numbers  $l_{kj}, \bar{l}_{kj}, \gamma_{kj}$  ( $k = 1, 2; j = 1, \dots, m$ ), and  $a, b, t^*, n$ .

**Remark 1.1.** Under the conditions of Theorem 1.2, for every

$$q \in \tilde{L}_{2n-2m-2, 2m-2}^2(\lceil a, b \rceil)$$

the unique solution  $u$  of problem (1.8), (1.2) admits the estimate

$$(1.18) \quad \|u^{(m)}\|_{\tilde{C}_1^{m-1}} \leq r_n \|q\|_{\tilde{L}_{2n-2m-2, 2m-2}^2},$$

with

$$r_n = \left( 1 + \sum_{j=1}^m \frac{2^{m-j+1/2}}{(m-j)!(2m-2j+1)^{1/2}(b-a)^{m-j+1/2}} \right) \times \frac{2^m(1+b-a)(2n-2m-1)}{(\nu_n - 2 \max\{B_0, B_1\})(2m-1)!!}.$$

### 1.2. Theorems on solvability of problem (1.1), (1.2).

Define an operator  $P: C_1^{m-1}(\lceil a, b \rceil) \times C_1^{m-1}(\lceil a, b \rceil) \rightarrow L_{\text{loc}}(\lceil a, b \rceil)$  by the equality

$$(1.19) \quad P(x, y)(t) = \sum_{j=1}^m p_j(x)(t)y^{(j-1)}(\tau_j(t)) \quad \text{for } a < t < b$$

where  $p_j: C_1^{m-1}(\lceil a, b \rceil) \rightarrow L_{\text{loc}}(\lceil a, b \rceil)$ , and  $\tau_j \in M(\lceil a, b \rceil)$ . Also, for any  $\gamma > 0$  define a set  $A_\gamma$  by the relation

$$(1.20) \quad A_\gamma = \{x \in \tilde{C}_1^{m-1}(\lceil a, b \rceil): \|x\|_{\tilde{C}_1^{m-1}} \leq \gamma\}.$$

For formulating the a priori boundedness principle we have to introduce

**Definition 1.1.** Let  $\gamma_0$  and  $\gamma$  be positive numbers. We say that the continuous operator  $P: C_1^{m-1}(]a, b[) \times C_1^{m-1}(]a, b[) \rightarrow L_n(]a, b[)$  is  $\gamma_0, \gamma$  consistent with boundary condition (1.2) if:

(i) For any  $x \in A_{\gamma_0}$  and almost all  $t \in ]a, b[$  the inequality

$$(1.21) \quad \sum_{j=1}^m |p_j(x)(t)x^{(j-1)}(\tau_j(t))| \leq \delta(t, \|x\|_{\tilde{C}_1^{m-1}}) \|x\|_{\tilde{C}_1^{m-1}}$$

holds, where  $\delta \in D_n(]a, b[ \times \mathbb{R}^+)$ .

(ii) For any  $x \in A_{\gamma_0}$  and  $q \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  the equation

$$(1.22) \quad y^{(n)}(t) = \sum_{j=1}^m p_j(x)(t)y^{(j-1)}(\tau_j(t)) + q(t)$$

under boundary conditions (1.2) has a unique solution  $y$  in the space  $\tilde{C}^{n-1, m}(]a, b[)$  and

$$(1.23) \quad \|y\|_{\tilde{C}_1^{m-1}} \leq \gamma \|q\|_{\tilde{L}_{2n-2m-2, 2m-2}^2}.$$

**Definition 1.2.** We say that the operator  $P$  is  $\gamma$  consistent with boundary condition (1.2), if the operator  $P$  is  $\gamma_0, \gamma$  consistent with boundary condition (1.2) for any  $\gamma_0 > 0$ .

In the sequel it will always be assumed that the operator  $F_p$  defined by equality

$$F_p(x)(t) = \left| F(x)(t) - \sum_{j=1}^m p_j(x)(t)x^{(j-1)}(\tau_j(t))(t) \right|$$

is continuously acting from  $C_1^{m-1}(]a, b[)$  to  $L_{\tilde{L}_{2n-2m-2, 2m-2}^2}(]a, b[)$ , and

$$(1.24) \quad \tilde{F}_p(t, \varrho) \equiv \sup\{F_p(x)(t) : \|x\|_{C_1^{m-1}} \leq \varrho\} \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$$

for each  $\varrho \in [0, +\infty[$ .

Then the following theorem is valid

**Theorem 1.3.** *Let the operator  $P$  be  $\gamma_0, \gamma$  consistent with boundary condition (1.2), and let there exist a positive number  $\varrho_0 \leq \gamma_0$ , such that*

$$(1.25) \quad \|\tilde{F}_p(\cdot, \min\{2\varrho_0, \gamma_0\})\|_{\tilde{L}_{2n-2m-2, 2m-2}^2} \leq \frac{\gamma_0}{\gamma}.$$

*Let, moreover, for any  $\lambda \in ]0, 1[$  an arbitrary solution  $x \in A_{\gamma_0}$  of the equation*

$$(1.26) \quad x^{(n)}(t) = (1 - \lambda)P(x, x)(t) + \lambda F(x)(t)$$

*under the conditions (1.2) admit the estimate*

$$(1.27) \quad \|x\|_{\tilde{C}_1^{m-1}} \leq \varrho_0.$$

*Then problem (1.1), (1.2) is solvable in the space  $\tilde{C}^{n-1, m}(]a, b[)$ .*

Theorem 1.3 with  $\varrho_0 = \gamma_0$  immediately yields

**Corollary 1.1.** *Let the operator  $P$  be  $\gamma_0, \gamma$  consistent with boundary condition (1.2), and*

$$(1.28) \quad |F(x)(t) - \sum_{j=1}^m p_j(x)(t)x^{(j-1)}(\tau_j(t))(t)| \leq \eta(t, \|x\|_{\tilde{C}_1^{m-1}})$$

*for  $x \in A_{\gamma_0}$  and almost all  $t \in ]a, b[$ , and*

$$(1.29) \quad \|\eta(\cdot, \gamma_0)\|_{\tilde{L}_{2n-2m-2, 2m-2}^2} \leq \frac{\gamma_0}{\gamma},$$

*where  $\eta \in D_{2n-2m-2, 2m-2}(]a, b[ \times \mathbb{R}^+)$ . Then problem (1.1), (1.2) is solvable in the space  $\tilde{C}^{n-1, m}(]a, b[)$ .*

**Corollary 1.2.** *Let the operator  $P$  be  $\gamma$  consistent with boundary condition (1.2), let inequality (1.28) hold for  $x \in \tilde{C}_1^{m-1}(]a, b[)$  and almost all  $t \in ]a, b[$ , where  $\eta(\cdot, \varrho) \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  for any  $\varrho \in \mathbb{R}^+$ , and*

$$(1.30) \quad \limsup_{\varrho \rightarrow +\infty} \frac{1}{\varrho} \|\eta(\cdot, \varrho)\|_{\tilde{L}_{2n-2m-2, 2m-2}^2} < \frac{1}{\gamma}.$$

*Then problem (1.1), (1.2) is solvable in the space  $\tilde{C}^{n-1, m}(]a, b[)$ .*

When we discuss problem (1.41), (1.2), and  $n = 2m + 1$ , we assume that the continuous operator  $p_1: \tilde{C}_1^{m-1}(]a, b[) \rightarrow L_{\text{loc}}(]a, b[)$  is such that

$$(1.31) \quad \limsup_{t \rightarrow b} \left| (b-t)^{2m-1} \int_{t_1}^t p_1(x)(s) \, ds \right| < +\infty \quad \left( t_1 = \frac{a+b}{2} \right)$$

for any  $x \in \tilde{C}_1^{m-1}(]a, b[)$ .

Now define operators  $h_j: C_1^{m-1}(]a, b[) \times ]a, b[ \times ]a, b[ \rightarrow \mathbb{R}_+$ ,  $f_j: C_1^{m-1}(]a, b[) \times ]a, b[ \times M(]a, b[) \rightarrow L_{\text{loc}}(]a, b[ \times ]a, b[)$  ( $j = 1, \dots, m$ ) by the equalities

$$(1.32) \quad \begin{aligned} h_1(x, t, s) &= \left| \int_s^t (\xi - a)^{n-2m} [(-1)^{n-m} p_1(x)(\xi)]_+ \, d\xi \right|, \\ h_j(x, t, s) &= \left| \int_s^t (\xi - a)^{n-2m} p_j(x)(\xi) \, d\xi \right| \quad (j = 2, \dots, m), \end{aligned}$$

and

$$(1.33) \quad f_j(x, c, \tau_j)(t, s) = \left| \int_s^t (\xi - a)^{n-2m} |p_j(x)(\xi)| \left| \int_\xi^{\tau_j(\xi)} (\xi_1 - c)^{2(m-j)} \, d\xi_1 \right|^{1/2} \, d\xi \right|.$$

**Theorem 1.4.** *Let the continuous operator  $P: C_1^{m-1}(]a, b[) \times C_1^{m-1}(]a, b[) \rightarrow L_n(]a, b[)$  fulfil the condition (1.21) where  $\delta \in D_n(]a, b[ \times \mathbb{R}^+)$ ,  $\tau_j \in M(]a, b[)$  and the numbers  $\gamma_0, t^* \in ]a, b[$ ,  $l_{kj} > 0$ ,  $\bar{l}_{kj} > 0$ ,  $\gamma_{kj} > 0$  ( $k = 0, 1; j = 1, \dots, m$ ), are such that the inequalities*

$$(1.34) \quad (t - a)^{2m-j} h_j(x, t, s) \leq l_{0j}, \quad \limsup_{t \rightarrow a} (t - a)^{m-\frac{1}{2}-\gamma_{0j}} f_j(x, a, \tau_j)(t, s) \leq \bar{l}_{0j}$$

for  $a < t \leq s \leq t^*$ ,  $\|x\|_{\tilde{C}_1^{m-1}} \leq \gamma_0$ ,

$$(1.35) \quad (b - t)^{2m-j} h_j(x, t, s) \leq l_{1j}, \quad \limsup_{t \rightarrow b} (b - t)^{m-\frac{1}{2}-\gamma_{1j}} f_j(x, b, \tau_j)(t, s) \leq \bar{l}_{1j}$$

for  $t^* \leq s \leq t < b$ ,  $\|x\|_{\tilde{C}_1^{m-1}} \leq \gamma_0$ , and conditions (1.13), (1.14) hold. Let moreover the operator  $F$  and a function  $\eta \in D_{2n-2m-2, 2m-2}(]a, b[ \times \mathbb{R}^+)$  be such that condition (1.28) and inequality

$$(1.36) \quad \|\eta(\cdot, \gamma_0)\|_{\tilde{L}_{2n-2m-2, 2m-2}^2} < \frac{\gamma_0}{r_n},$$

are fulfilled, where

$$\begin{aligned} r_n &= \left( 1 + \sum_{j=1}^m \frac{2^{m-j+1/2}}{(m-j)!(2m-2j+1)^{1/2}(b-a)^{m-j+1/2}} \right) \\ &\quad \times \frac{2^m(1+b-a)(2n-2m-1)}{(\nu_n - 2 \max\{B_0, B_1\})(2m-1)!!}. \end{aligned}$$

Then problem (1.1), (1.2) is solvable in the space  $\tilde{C}^{n-1, m}(]a, b[)$ .

**Theorem 1.5.** Let the operator  $F$  and the function  $\eta$  be such that conditions (1.28), (1.30) hold and the continuous operator  $P: C_1^{m-1}(]a, b[) \times C_1^{m-1}(]a, b[) \rightarrow L_n(]a, b[) \times \mathbb{R}^+$  fulfils condition (1.21) where  $\delta \in D_n(]a, b[ \times \mathbb{R}^+)$ . Let moreover measurable functions  $\tau_j \in M(]a, b[)$  and numbers  $t^* \in ]a, b[$ ,  $l_{kj} > 0$ ,  $\bar{l}_{kj} > 0$ ,  $\gamma_{kj} > 0$  ( $k = 0, 1$ ;  $j = 1, \dots, m$ ) be such that the inequalities

$$(1.37) \quad (t-a)^{2m-j} h_j(x, t, s) \leq l_{0j}, \quad \limsup_{t \rightarrow a} (t-a)^{m-\frac{1}{2}-\gamma_{0j}} f_j(x, a, \tau_j)(t, s) \leq \bar{l}_{0j}$$

for  $a < t \leq s \leq t^*$ ,  $x \in \tilde{C}_1^{m-1}(]a, b[)$ ,

$$(1.38) \quad (b-t)^{2m-j} h_j(x, t, s) \leq l_{1j}, \quad \limsup_{t \rightarrow b} (b-t)^{m-\frac{1}{2}-\gamma_{1j}} f_j(x, b, \tau_j)(t, s) \leq \bar{l}_{1j}$$

for  $t^* \leq s \leq t < b$ ,  $x \in \tilde{C}_1^{m-1}(]a, b[)$ , and conditions (1.13), (1.14) hold. Then problem (1.1), (1.2) is solvable in the space  $\tilde{C}^{m-1, m}(]a, b[)$ .

**Remark 1.2.** Let  $\gamma_0 > 0$ , let the operators  $\alpha_j(t)p_j(x)(t)$  ( $j = 1, \dots, m$ ) be continuously acting from the space  $C_1^{m-1}(]a, b[)$  to the space  $L_n(]a, b[)$ , let there exist functions  $\delta_j \in D_n(]a, b[)$  such that for any  $x \in A_{\gamma_0}$

$$(1.39) \quad |p_j(x)(t)| \alpha_j(t) \leq \delta_j(t, \|x\|_{\tilde{C}_1^{m-1}}) \quad \text{for } a < t < b,$$

and constants  $\kappa > 0$ ,  $\varepsilon > 0$  such that

$$(1.40) \quad \begin{aligned} |\tau_j(t) - t| &\leq \kappa(t-a) \quad (j = 1, \dots, m) \quad \text{for } a < t < a + \varepsilon, \\ |\tau_j(t) - t| &\leq \kappa(b-t) \quad (j = 1, \dots, m) \quad \text{for } b - \varepsilon < t < b. \end{aligned}$$

Then the operator  $P$  defined by equality (1.19), continuously acting from  $A_{\gamma_0}$  to the space  $L_n(]a, b[)$ , and there exists a function  $\delta \in D_n(]a, b[)$  such that item (ii) of Definition 1.1 holds.

Now consider the equation with deviating arguments

$$(1.41) \quad u^{(n)}(t) = f(t, u(\tau_1(t)), u'(\tau_2(t)), \dots, u^{(m-1)}(\tau_m(t))) \quad \text{for } a < t < b,$$

where  $-\infty < a < b < +\infty$ ,  $f: ]a, b[ \times \mathbb{R}^m \rightarrow \mathbb{R}$  is a function satisfying the local Caratheodory conditions and  $\tau_j \in M(]a, b[)$  ( $j = 0, \dots, n-1$ ) are measurable functions.

**Corollary 1.3.** Let the functions  $\tau_j \in M(]a, b[)$  and the numbers  $t^* \in ]a, b[$ ,  $\kappa \geq 0$ ,  $\varepsilon > 0$ ,  $l_{kj} > 0$ ,  $\bar{l}_{kj} > 0$ ,  $\gamma_{kj} > 0$  ( $k = 1, 2; j = 1, \dots, m$ ) be such that conditions (1.13), (1.14), (1.15), (1.16), (1.40) and the inclusions

$$(1.42) \quad \alpha_j p_j \in L_n(]a, b[) \quad (j = 1, \dots, m)$$

are fulfilled. Let moreover

$$(1.43) \quad \left| f(t, x(\tau_1(t)), x'(\tau_2(t)), \dots, x^{(m-1)}(\tau_m(t))) - \sum_{j=1}^m p_j(t) x^{(j-1)}(\tau_j(t))(t) \right| \leq \eta(t, \|x\|_{\tilde{C}_1^{m-1}})$$

for  $x \in \tilde{C}_1^{m-1}(]a, b[)$  and almost all  $t \in ]a, b[$ , where  $\eta(\cdot, \varrho) \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  for any  $\varrho \in \mathbb{R}^+$ , and let condition (1.30) hold. Then problem (1.41), (1.2) is solvable in the space  $\tilde{C}^{n-1, m}(]a, b[)$ .

**Remark 1.3.** Conditions (1.42) do not imply conditions (1.6).

Now for illustration of our results consider on  $]a, b[$  the second order functional-differential equations

$$(1.44) \quad u''(t) = -\frac{\lambda |u(t)|^k}{[(t-a)(b-t)]^{2+k/2}} u(\tau(t)) + q(x)(t),$$

$$(1.45) \quad u''(t) = -\frac{\lambda |\sin u^k(t)|}{[(t-a)(b-t)]^2} u(\tau(t)) + q(x)(t),$$

where  $\lambda, k \in \mathbb{R}^+$ , the function  $\tau \in M(]a, b[)$ , the operator  $q: C_1^{m-1}(]a, b[) \rightarrow \tilde{L}_{0,0}^2(]a, b[)$  is continuous and

$$\eta(t, \varrho) \equiv \sup\{|q(x)(t)|: \|x\|_{\tilde{C}_1^{m-1}} \leq \varrho\} \in \tilde{L}_{0,0}^2(]a, b[).$$

Then Theorems 1.4 and 1.5 yield

**Corollary 1.4.** Let a continuous operator  $q: C_1^{m-1}(]a, b[) \rightarrow \tilde{L}_{0,0}^2(]a, b[)$ , a function  $\tau \in M(]a, b[)$ , and numbers  $\gamma_0 > 0$ ,  $\lambda \geq 0$ ,  $k > 0$ , be such that

$$(1.46) \quad |\tau(t) - t| \leq \begin{cases} (t-a)^{3/2} & \text{for } a < t \leq (a+b)/2, \\ (b-t)^{3/2} & \text{for } (a+b)/2 \leq t < b, \end{cases}$$

$$(1.47) \quad \|\eta(t, \gamma_0)\|_{\tilde{L}_{0,0}^2} \leq \left(1 + \sqrt{\frac{2}{b-a}}\right)^{-1} \frac{(b-a)^2 - 16\lambda\gamma_0^k(1 + [2(b-a)]^{1/4})}{2(1+b-a)(b-a)^2},$$

and

$$(1.48) \quad \lambda < \frac{(b-a)^2}{32\gamma_0^k(1+[2(b-a)]^{1/4})}.$$

Then problem (1.44), (1.2) is solvable.

**Corollary 1.5.** Let a continuous operator  $q: C_1^{m-1}(]a, b[) \rightarrow \widetilde{L}_{0,0}^2(]a, b[)$ , a function  $\tau \in M(]a, b[)$ , and a number  $\lambda \geq 0$  be such that inequalities (1.30) with  $n = 2$ , (1.46) and

$$(1.49) \quad \lambda < \frac{(b-a)^2}{32(1+[2(b-a)]^{1/4})},$$

hold. Then problem (1.45), (1.2) is solvable.

## 2. AUXILIARY PROPOSITIONS

### 2.1. Lemmas on some properties of the equation $x^{(n)}(t) = \lambda(t)$ .

First, we introduce two lemmas without proofs. The first lemma is proved in [2].

**Lemma 2.1.** Let  $i \in 1, 2$ ,  $x \in \widetilde{C}_{\text{loc}}^{m-1}(]t_0, t_1[)$  and

$$(2.1) \quad x^{(j-1)}(t_i) = 0 \quad (j = 1, \dots, m), \quad \int_{t_0}^{t_1} |x^{(m)}(s)|^2 ds < +\infty.$$

Then

$$(2.2) \quad \left| \int_{t_i}^t \frac{(x^{(j-1)}(s))^2}{(s-t_i)^{2m-2j+2}} ds \right|^{1/2} \leq \frac{2^{m-j+1}}{(2m-2j+1)!!} \left| \int_{t_i}^t |x^{(m)}(s)|^2 ds \right|^{1/2}$$

for  $t_0 \leq t \leq t_1$ .

This second lemma is a particular case of Lemma 4.1 in [11].

**Lemma 2.2.** *If  $x \in C_{\text{loc}}^{n-1}(]a, a_1])$ , then for any  $s, t \in ]a, a_1]$  the equality*

$$(-1)^{n-m} \int_s^t (\xi - a)^{n-2m} x^{(n)}(\xi) x(\xi) \, d\xi = w_n(x)(t) - w_n(x)(s) + \nu_n \int_s^t |x^{(m)}(\xi)|^2 \, d\xi$$

is valid, where  $\nu_{2m} = 1$ ,  $\nu_{2m+1} = \frac{1}{2}(2m + 1)$ ,

$$\begin{aligned} w_{2m}(x)(t) &= \sum_{j=1}^m (-1)^{m+j-1} x^{(2m-j)}(t) x(t), \\ w_{2m+1}(x)(t) &= \sum_{j=1}^m (-1)^{m+j} [(t-a)x^{(2m+1-j)}(t) - jx^{(2m-j)}(t)] x^{(j-1)}(t) \\ &\quad - \frac{t-a}{2} |x^{(m)}(t)|^2. \end{aligned}$$

**Lemma 2.3.** *Let numbers  $a_1 \in ]a, b[$ ,  $t_{0k} \in ]a, a_1[$ , and  $\varepsilon_{ik}, \varepsilon_i, \beta_k, \beta \in \mathbb{R}^+$ ,  $k \in \mathbb{N}$ ,  $i = 1, \dots, n - m$  be such that*

$$(2.3) \quad \lim_{k \rightarrow +\infty} t_{0k} = a, \quad \lim_{k \rightarrow +\infty} \beta_k = \beta, \quad \lim_{k \rightarrow +\infty} \varepsilon_{i,k} = \varepsilon_i.$$

Let, moreover,

$$(2.4) \quad \lambda \in \tilde{L}_{2n-2m-2,0}^2(]a, a_1])$$

be a nonnegative function,  $x_k \in \tilde{C}^{n-1,m}(]a, b])$  a solution of the problem

$$(2.5) \quad x^{(n)}(t) = \beta_k \lambda(t),$$

$$(2.6) \quad x^{(i-1)}(t_{0k}) = 0 \quad (i = 1, \dots, m), \quad x^{(i-1)}(a_1) = \varepsilon_{i,k} \quad (i = 1, \dots, n - m),$$

and  $x \in \tilde{C}^{n-1,m}(]a, b])$  a solution of the problem

$$(2.7) \quad x^{(n)}(t) = \beta \lambda(t),$$

$$(2.8) \quad x^{(i-1)}(a) = 0 \quad (i = 1, \dots, m), \quad x^{(i-1)}(a_1) = \varepsilon_i \quad (i = 1, \dots, n - m).$$

Then

$$(2.9) \quad \lim_{k \rightarrow +\infty} x_k^{(j-1)}(t) = x^{(j-1)}(t) \quad (j = 1, \dots, n) \quad \text{uniformly in } ]a, a_1].$$

**Proof.** First, let us prove our lemma under the assumption that there exists a number  $r_1 > 0$  such that the estimates

$$(2.10) \quad \int_{t_{0k}}^{a_1} |x_k^{(m)}(s)|^2 \, ds \leq r_1, \quad k \in \mathbb{N}$$

hold. Now, suppose that  $t_1, \dots, t_n$  are such numbers that  $t_{0k} < t_1 < \dots < t_n < a_1$  ( $k \in \mathbb{N}$ ), and  $g_i(t)$  are polynomials of  $(n-1)$ -th degree, satisfying the conditions  $g_j(t_j) = 1, g_j(t_i) = 0$  ( $i \neq j; i, j = 1, \dots, n$ ). Then if  $x_k$  is a solution of problem (2.5), (2.6), and  $x$  is a solution of problem (2.7), (2.8). For the solution  $x - x_k$  of the equation  $d^n(x(t) - x_k(t))/dt^n = (\beta - \beta_k)\lambda(t)$ , the representation

$$(2.11) \quad x(t) - x_k(t) = \sum_{j=1}^n \left( (x(t_j) - x_k(t_j)) - \frac{\beta - \beta_k}{(n-1)!} \int_{t_1}^{t_j} (t_j - s)^{n-1} \lambda(s) ds \right) g_j(t) + \frac{\beta - \beta_k}{(n-1)!} \int_{t_1}^t (t - s)^{n-1} \lambda(s) ds \quad k \in \mathbb{N} \quad \text{for } t_{0k} \leq t \leq a_1$$

is valid. On the other hand, in view of inequality (2.10), the identities

$$x_k^{(i-1)}(t) = \frac{1}{(m-i)!} \int_{t_{0k}}^t (t-s)^{m-i} x_k^{(m)}(s) ds \quad (i = 1, 2, k \in \mathbb{N})$$

by Schwartz inequality yield

$$(2.12) \quad |x_k^{(i-1)}(t)| \leq r_2 (t - a)^{m-i-1/2} \quad \text{for } t_{0k} \leq t \leq a_1 \quad (i = 1, 2, k \in \mathbb{N}),$$

where  $r_2 = r_1 / (m-i)! \sqrt{2m-2i+1}$ . By virtue of the Arzela-Ascoli lemma and (2.3), (2.12) the sequence  $\{x_k\}_{k=1}^{+\infty}$  contains a subsequence  $\{x_{k_l}\}_{l=1}^{+\infty}$  which is uniformly convergent in  $]a, a_1]$ . Suppose  $\lim_{l \rightarrow +\infty} x_{k_l}(t) = x_0(t)$ . Thus (2.11) by (2.3) yields the existence of such  $r_3 > 0$  that

$$|x_{k_l}^{(j-1)}(t)| \leq r_3 + |x^{(j-1)}(t)| \quad (j = 1, \dots, n) \quad \text{for } t_{0k_l} \leq t \leq a_1,$$

and then without loss of generality we can assume that

$$(2.13) \quad \lim_{l \rightarrow +\infty} x_{k_l}^{(j-1)}(t) = x_0^{(j-1)}(t) \quad (j = 1, \dots, n) \quad \text{uniformly in } ]a, a_1].$$

Then in virtue of (2.3), (2.11), and (2.13) we have

$$x(t) - x_0(t) = \sum_{j=1}^n ((x(t_j) - x_0(t_j))) g_j(t) \quad \text{for } a \leq t \leq a_1.$$

From the last two relations by (2.10) it is clear that  $x^{(n)} = x_0^{(n)}$  and  $x_0 \in \tilde{C}^{n-1,m}(]a, b])$ . So, the function  $x_0 \in \tilde{C}^{n-1,m}(]a, b])$  is a solution of problem (2.7), (2.8). In view of (2.4) all the conditions of Theorem 1.1 are fulfilled, thus problem

(2.7), (2.8) is uniquely solvable in the space  $\widetilde{C}^{n-1,m}(]a, b[)$  and  $x = x_0$ . Therefore (2.13) implies

$$(2.14) \quad \lim_{l \rightarrow +\infty} x_{k_l}^{(j-1)}(t) = x^{(j-1)}(t) \quad (j = 1, \dots, n) \quad \text{uniformly in } ]a, a_1].$$

Now suppose that relations (2.9) are not fulfilled. Then there exist  $\delta \in ]0, \frac{1}{2}(a_1 - a)[$ ,  $\varepsilon > 0$ , and an increasing sequence of natural numbers  $\{k_l\}_{l=1}^{+\infty}$  such that

$$(2.15) \quad \max \left\{ \sum_{j=1}^n |x_{k_l}^{(j-1)}(t) - x^{(j-1)}(t)| : a + \delta \leq t \leq a_1 \right\} > \varepsilon \quad (l \in \mathbb{N}).$$

By virtue of the Arzela-Ascoli lemma and condition (2.10) the sequence  $\{x_{k_l}^{(j-1)}\}_{l=1}^{+\infty}$  ( $j = 1, \dots, m$ ), without loss of generality, can be assumed to be uniformly converging in  $]a + \delta, a_1]$ . Then, in view of what we have shown above, equality (2.14) holds. But this contradicts condition (2.15). Thus (2.9) holds if the conditions (2.10) are fulfilled.

Let now the conditions (2.10) be not fulfilled. Then there exists a subsequence  $\{t_{0k_l}\}_{l=1}^{+\infty}$  of the sequence  $\{t_{0k}\}_{k=1}^{+\infty}$ , such that

$$(2.16) \quad \int_{t_{0k}}^{a_1} |x_{k_l}^{(m)}(s)|^2 ds \geq l \quad (l \in \mathbb{N}).$$

Suppose that  $\beta_l = \left( \int_{t_{0k}}^{a_1} |x_{k_l}^{(m)}(s)|^2 ds \right)^{-1}$  and  $v_l(t) = u_{k_l}(t)\beta_l$ . Thus in view of (2.16) and our notation

$$(2.17) \quad \int_{t_{0k_l}}^{a_1} |v_{k_l}^{(m)}(s)|^2 ds = 1 \quad (l \in \mathbb{N}), \quad \lim_{l \rightarrow +\infty} \beta_l = 0,$$

$$(2.18) \quad v_l^{(n)}(t) = \beta_l \lambda(t),$$

$$(2.19) \quad v_l^{(i-1)}(t_{0k_l}) = 0 \quad (i = 1, \dots, m),$$

$$v_l^{(i-1)}(a_1) = \varepsilon_{i,k_l} \beta_l \quad (i = 1, \dots, n - m, l \in \mathbb{N}).$$

From the first part of our lemma it follows by (2.17) that the limit  $\lim_{l \rightarrow +\infty} v_l(t) \equiv v_0(t)$  exists, and  $v_0$  is a solution of the homogeneous problem corresponding to (2.18), (2.19). Thus  $v_0 \equiv 0$ . On the other hand, from (2.17) it is clear that  $\int_{t_{0k_l}}^{a_1} |v_0^{(m)}(s)|^2 ds = 1$ , which contradicts  $v_0 \equiv 0$ . Thus our assumption is invalid and (2.10) holds.  $\square$

Analogously one can prove

**Lemma 2.4.** Let numbers  $b_1 \in ]a, b[$ ,  $t_{0k} \in ]b_1, b[$ , and  $\varepsilon_{ik}$ ,  $\varepsilon_i$ ,  $\beta_k$ ,  $\beta \in \mathbb{R}^+$ ,  $k \in \mathbb{N}$ ,  $i = 1, \dots, n - m$  be such that

$$\lim_{k \rightarrow +\infty} t_{0k} = b, \quad \lim_{k \rightarrow +\infty} \beta_k = \beta, \quad \lim_{k \rightarrow +\infty} \varepsilon_{i,k} = \varepsilon_i.$$

Let, moreover,  $\lambda \in \tilde{L}_{0,2m-2}^2(]b_1, b[)$  be a nonnegative function,  $x_k \in \tilde{C}^{n-1,m}(]a, b[)$  a solution of the problem (2.5) under the conditions

$$x^{(i-1)}(b_1) = \varepsilon_{i,k} \quad (i = 1, \dots, m), \quad x^{(i-1)}(t_{0k}) = 0 \quad (i = 1, \dots, n - m),$$

and  $x \in \tilde{C}^{n-1,m}(]a, b[)$  a solution of the equation (2.7) under the conditions

$$(2.20) \quad x^{(i-1)}(b_1) = \varepsilon_i \quad (i = 1, \dots, m), \quad x^{(i-1)}(b) = 0 \quad (i = 1, \dots, n - m).$$

Then the equalities (2.9) hold.

**Lemma 2.5.** Let  $a < a_1 < b_1 < b$ ,  $\varepsilon_i \in \mathbb{R}^+$  and let

$$\lambda \in \tilde{L}_{2n-2m-2,0}^2(]a, a_1[) \quad (\lambda \in \tilde{L}_{0,2m-2}^2(]b_1, b[))$$

be a nonnegative function. Then for the solution  $x \in \tilde{C}^{n-1,m}(]a, b[)$  of the problem (2.7), (2.8) ((2.7), (2.20)) with  $\beta = 1$ , the estimate

$$(2.21) \quad \int_a^{a_1} |x^{(m)}(s)|^2 ds \leq \Theta_1(x, a_1, \lambda) \\ \left( \int_{b_1}^b |x^{(m)}(s)|^2 ds \leq \Theta_2(x, b_1, \lambda) \right) \quad (k \in \mathbb{N})$$

is valid, where

$$(2.22) \quad \Theta_1(x, a_1, \lambda) = 2|w_n(x)(a_1)| + \gamma_1 \|\lambda\|_{\tilde{L}_{2n-2m-2,0}^2(]a, a_1[)}^2, \\ (\Theta_2(x, b_1, \lambda) = 2|w_n(x)(b_1)| + \gamma_2 \|\lambda\|_{\tilde{L}_{0,2m-2}^2(]b_1, b[)}^2),$$

and

$$\gamma_1 = \left( \frac{2^{m-1}(2m+1)}{(2m-1)!!} \right)^2, \quad \gamma_2 = \left( \frac{2^{m-1}(2m+1)(b-a+1)}{(2m-1)!!} \right)^2.$$

**Proof.** Suppose that  $x_k$  is a solution of problem (2.5), (2.6) with  $\beta_k = 1$ ,  $\varepsilon_{ik} = \varepsilon_i$ . Then in view of Lemma 2.3, relations (2.9) hold. On the other hand, by Lemma 2.2 we get

$$(2.23) \quad \nu_n \int_{t_{0k}}^{a_1} |x_k^{(m)}(s)|^2 ds \leq -w_n(x_k)(a_1) + \int_{t_{0k}}^{a_1} (s-a)^{n-2m} \lambda(s) |x_k(s)| ds.$$

Now, on the basis of Lemma 2.1, Schwartz's and Young's inequalities we get

$$\begin{aligned}
& \left| \int_{t_{0k}}^{a_1} (s-a)^{n-2m} \lambda(s) x_k(s) ds \right| \\
&= \left| \int_{t_{0k}}^{a_1} [(n-2m)x_k(s) + (s-a)^{n-2m} x_k'(s)] \left( \int_s^{a_1} \lambda(\xi) d\xi \right) ds \right| \\
&\leq \left[ (n-2m) \left( \int_{t_{0k}}^{a_1} \frac{x_k^2(s)}{(s-a)^{2m}} ds \right)^{1/2} \right. \\
&\quad \left. + \left( \int_{t_{0k}}^{a_1} \frac{x_k'^2(s)}{(s-a)^{2m-2}} ds \right)^{1/2} \right] \|\lambda\|_{\tilde{L}_{2n-2m-2,0}(\lceil a, a_1 \rceil)} \\
&\leq \frac{2^{m-1}(2m+1)}{(2m-1)!!} \left( \int_{t_{0k}}^{a_1} |x_k^{(m)}(s)|^2 ds \right)^{1/2} \|\lambda\|_{\tilde{L}_{2n-2m-2,0}(\lceil a, a_1 \rceil)} \\
&\leq \frac{1}{2} \int_{t_{0k}}^{a_1} |x_k^{(m)}(s)|^2 ds + \frac{1}{2} \left( \frac{2^{m-1}(2m+1)}{(2m-1)!!} \right)^2 \|\lambda\|_{\tilde{L}_{2n-2m-2,0}(\lceil a, a_1 \rceil)}^2.
\end{aligned}$$

Thus from (2.23) by the definition of the numbers  $\nu_n$  we immediately obtain the estimate

$$\int_{t_{0k}}^{a_1} |x_k^{(m)}(s)| ds \leq 2|w_n(x_k)(a_1)| + \left( \frac{2^{m-1}(2m+1)}{(2m-1)!!} \right)^2 \|\lambda\|_{\tilde{L}_{2n-2m-2,0}(\lceil a, a_1 \rceil)}^2 \quad (k \in \mathbb{N}).$$

By (2.9) from the last inequality (2.21) and (2.22) follow. Thus the lemma is proved for problem (2.7), (2.8).

Analogously, by using Lemma 2.4 one can prove the case of problem (2.7), (2.20).  $\square$

## 2.2. Lemmas on Banach space $\tilde{C}_1^{m-1}(\lceil a, b \rceil)$ .

**Definition 2.1.** Let  $\varrho \in \mathbb{R}^+$  and let the function  $\eta \in L_{\text{loc}}(\lceil a, b \rceil)$  be nonnegative. Then  $S(\varrho, \eta)$  is a set of such  $y \in C_{\text{loc}}^{n-1}(\lceil a, b \rceil)$  that

$$(2.24) \quad \left| y^{(i-1)} \left( \frac{a+b}{2} \right) \right| \leq \varrho \quad (i = 1, \dots, n),$$

$$(2.25) \quad |y^{(n-1)}(t) - y^{(n-1)}(s)| \leq \int_s^t \eta(\xi) d\xi \quad \text{for } a < s \leq t < b,$$

and

$$(2.26) \quad y^{(i-1)}(a) = 0 \quad (i = 1, \dots, m), \quad y^{(i-1)}(b) = 0 \quad (i = 1, \dots, n-m).$$

**Lemma 2.6.** Let for a function  $y \in \widetilde{C}^{n-1,m}([a, b])$ , conditions (2.26) be satisfied. Then  $y \in \widetilde{C}_1^{m-1}([a, b])$  and the estimates

$$(2.27) \quad |y^{(i-1)}(t)| \leq \frac{|t - c_k|^{m-i+1/2}}{(m-i)!(2m-2i+1)^{1/2}} \left| \int_{c_k}^t |y^{(m)}(s)|^2 ds \right|^{1/2} \quad \text{for } a < t < b,$$

$i = 1, \dots, m$ , hold for  $k = 1, 2$ , where  $c_1 = a$ ,  $c_2 = b$ .

**Proof.** First note that in view of inclusion  $y \in \widetilde{C}^{n-1,m}([a, b])$ , the equality

$$(2.28) \quad y^{(i-1)}(t) = \sum_{j=i}^l \frac{(t-c)^{j-i}}{(j-i)!} y^{(j-1)}(c) + \frac{1}{(l-i)!} \int_c^t (t-s)^{l-i} y^{(l)}(s) ds \quad \text{for } a < t < b$$

for  $i = 1, \dots, l$ ,  $l = 1, \dots, n$ , holds, where

- (1)  $c \in [a, b]$  if  $l \leq m$ ;
- (2)  $c \in ]a, b]$  if  $l = m + 1$  and  $n = 2m + 1$ ;
- (3)  $c \in ]a, b[$  if  $l > m$ ,

and there exists  $r > 0$  such that

$$(2.29) \quad \int_a^b |y^{(m)}(s)|^2 ds \leq r.$$

Equality (2.28) with  $l = m$ ,  $c = a$  and with  $l = m$ ,  $c = b$  by conditions (2.26), (2.29) and the Schwartz inequality yields (2.27). From (2.27) and (2.29) it is clear that  $y \in \widetilde{C}_1^m([a, b])$ .  $\square$

**Lemma 2.7.** Let  $\varrho \in \mathbb{R}^+$ , and let  $\eta \in \widetilde{L}_{2n-2m-2, 2m-2}^2([a, b])$  be a nonnegative function. Then  $S(\varrho, \eta)$  is a compact subset of the space  $\widetilde{C}_1^{m-1}([a, b])$ .

**Proof.** Condition (2.25) yields the inequality  $|y^{(n)}(t)| \leq \eta(t)$ . Thus there exists such a function  $\eta_1 \in \widetilde{L}_{2n-2m-2, 2m-2}^2([a, b])$  that

$$(2.30) \quad y^{(n)}(t) = \eta_1(t), \quad \text{for } a < t < b,$$

$$(2.31) \quad |\eta_1(t)| \leq \eta(t) \quad \text{for } a < t < b.$$

From Theorem 1.1 it follows that problem (2.30), (2.26) has a unique solution  $y \in C^{n-1,m}([a, b])$ , i.e. there exists  $r > 0$  such that the inequality (2.29) holds.

For any  $y \in S(\varrho, \eta)$ , from equality (2.28) with  $l = n$ , by (2.24), (2.30) and (2.31) we get

$$(2.32) \quad |y^{(i-1)}(t)| \leq \gamma_i(t) \quad \text{for } a < t < b \quad (i = 1, \dots, n),$$

where

$$\gamma_i(t) = \varrho_i + \frac{1}{(n-i)!} \left| \int_c^t (t-s)^{n-i} \eta(s) ds \right| \quad (i = 1, \dots, n).$$

Let now  $y_k \in S(\varrho, \eta)$  ( $k \in \mathbb{N}$ ). By virtue of the Arzela-Ascoli lemma and conditions (2.25), (2.32) the sequence  $\{y_k\}_{k=1}^{+\infty}$  contains a subsequence  $\{y_{k_l}\}_{l=1}^{+\infty}$  such that  $\{y_{k_l}^{(i-1)}\}_{l=1}^{+\infty}$  ( $i = 1, \dots, n$ ) are uniformly convergent on  $]a, b[$ . Thus without loss of generality we can assume that  $\{y_k^{(i-1)}\}_{k=1}^{+\infty}$  ( $i = 1, \dots, n-1$ ) are uniformly convergent on  $]a, b[$ . Let  $\lim_{k \rightarrow +\infty} y_k(t) = y_0(t)$ , then  $y_0 \in \tilde{C}_{\text{loc}}^{n-1}(]a, b[)$  and

$$(2.33) \quad \lim_{k \rightarrow +\infty} y_k^{(i-1)}(t) = y_0^{(i-1)}(t) \quad (i = 1, \dots, n) \quad \text{uniformly on } ]a, b[.$$

From (2.33) in view of the inclusions  $y_k \in S(\varrho, \eta)$  it immediately follows that

$$(2.34) \quad \left| y_0^{(i-1)} \left( \frac{a+b}{2} \right) \right| \leq \varrho \quad (i = 1, \dots, n),$$

$$(2.35) \quad y_0^{(i-1)}(a) = 0 \quad (j = 1, \dots, m), \quad y_0^{(i-1)}(b) = 0 \quad (j = 1, \dots, n-m),$$

and

$$(2.36) \quad |y_0^{(n-1)}(t) - y_0^{(n-1)}(s)| \leq \int_s^t \eta(\xi) d\xi \quad \text{for } a < s \leq t < b.$$

From (2.34), (2.35), (2.36) it is clear that  $y_0 \in S(\varrho, \eta)$ . To complete the proof we must show that

$$(2.37) \quad \lim_{k \rightarrow +\infty} \|y_k(t) - y_0(t)\|_{\tilde{C}_1^{m-1}} = 0$$

and

$$(2.38) \quad S(\varrho, \eta) \subset \tilde{C}_1^{m-1}(]a, b[).$$

Let,  $x_k = y_0 - y_k$  and  $a_1 \in ]a, b[$ ,  $b_1 \in ]a_1, b[$ . Then it is clear that  $x_k \in S(\varrho', \eta')$  where  $\varrho' = 2\varrho$ ,  $\eta' = 2\eta$ . Thus for any  $x_k$  there exists  $\eta_k \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  such that

$$(2.39) \quad x_k^{(n)}(t) = \eta_k(t),$$

$$(2.40) \quad x_k^{(i-1)}(a) = 0 \quad (i = 1, \dots, n), \quad x_k^{(i-1)}(b) = 0 \quad (i = 1, \dots, n-m)$$

where

$$(2.41) \quad |\eta_k(t)| \leq 2\eta(t) \quad \text{for } a < t < b \quad (k \in \mathbb{N}).$$

On the other hand, from (2.27) with  $y = x_k$ , in view of (2.40) we get

$$(2.42) \quad \begin{aligned} |x_k^{(i-1)}(t)| &\leq \left( \int_a^t |x_k^{(m)}(s)|^2 ds \right)^{1/2} (t-a)^{m-i+1/2} \quad \text{for } a < t < a_1, \\ |x_k^{(i-1)}(t)| &\leq \left( \int_t^b |x_k^{(m)}(s)|^2 ds \right)^{1/2} (b-t)^{m-i+1/2} \quad \text{for } b_1 < t < b, \end{aligned}$$

for  $i = 1, \dots, m$ .

Let now  $w_n$  be the operator defined in Lemma 2.2 and  $\Theta_1, \Theta_2$  the functions defined by (2.22) with  $\lambda = \eta_k$ . Then conditions (2.33) yield

$$(2.43) \quad \lim_{k \rightarrow +\infty} w_n(x_k)(a_1) = 0, \quad \lim_{k \rightarrow +\infty} w_n(x_k)(b_1) = 0 \quad (k \in \mathbb{N}),$$

and from the definition of the norm  $\|\cdot\|_{\tilde{L}_{\alpha,\beta}^2}$ , (2.41) and (2.43) it follows that for any  $\varepsilon > 0$  we can choose  $a_1 \in ]a, \min\{a+1, b\}[$ ,  $b_1 \in ]\max\{b-1, b\}, a_1[$  and  $k_0 \in \mathbb{N}$  such that

$$(2.44) \quad \begin{aligned} \Theta_1(x_k, a_1, 2\eta) &\leq \frac{\varepsilon}{6} (b-b_1)^{m-1/2} \quad (k \geq k_0), \\ \Theta_2(x_k, b_1, 2\eta) &\leq \frac{\varepsilon}{6} (a_1-a)^{m-1/2} \quad (k \geq k_0). \end{aligned}$$

By using Lemma 2.5 for  $x_k$ , in view of (2.42) and (2.44) we get

$$(2.45) \quad \int_a^{a_1} |x_k^{(m)}(s)|^2 ds \leq \frac{\varepsilon}{6}, \quad \int_{b_1}^b |x_k^{(m)}(s)|^2 ds \leq \frac{\varepsilon}{6} \quad (k \geq k_0),$$

$$(2.46) \quad \frac{|x_k^{(i-1)}(t)|}{\alpha_i(t)} \leq \frac{\varepsilon}{2m} \quad \text{for } t \in ]a, a_1] \cup [b_1, b[ \quad (1 \leq i \leq m, k \geq k_0).$$

Also, in view of (2.33) without loss of generality we can assume that

$$(2.47) \quad \frac{|x_k^{(i-1)}(t)|}{\alpha_i(t)} \leq \frac{\varepsilon}{2m} \quad \text{for } a_1 \leq t \leq b_1 \quad (1 \leq i \leq m, k \geq k_0),$$

and

$$(2.48) \quad \int_{a_1}^{b_1} |x_k^{(m)}(s)|^2 ds \leq \frac{\varepsilon}{6} \quad (k \geq k_0).$$

From (2.45), (2.46), (2.47), (2.48), equality (2.37) immediately follows.

Let now  $y \in S(\varrho, \eta)$  and  $y_k = \delta_k y$ , where  $\lim_{k \rightarrow +\infty} \delta_k = 0$ . Then by (2.33) it is clear that  $y_0 \equiv 0$  and then (2.37) implies  $y \in \tilde{C}_1^{m-1}(]a, b[)$ , i.e. the inclusion (2.38) holds.  $\square$

**Lemma 2.8.** Let  $\tau_j \in M(]a, b[)$ ,  $\alpha \geq 0$ ,  $\beta \geq 0$  and let there exist  $\delta \in ]0, b - a[$  such that

$$(2.49) \quad |\tau_j(t) - t| \leq k_1(t - a)^\beta \quad \text{for } a < t \leq a + \delta.$$

Then

$$\left| \int_t^{\tau(t)} (s - a)^\alpha ds \right| \leq \begin{cases} k_1[1 + k_1\delta^{\beta-1}]^\alpha (t - a)^{\alpha+\beta} & \text{for } \beta \geq 1, \\ k_1[\delta^{1-\beta} + k_1]^\alpha (t - a)^{\alpha+\beta} & \text{for } 0 \leq \beta < 1 \end{cases}$$

for  $a < t \leq a + \delta$ .

*Proof.* First note that

$$\left| \int_t^{\tau(t)} (s - a)^\alpha ds \right| \leq (\max\{\tau(t), t\} - a)^\alpha |\tau(t) - t| \quad \text{for } a \leq t \leq a + \delta,$$

and  $\max\{\tau(t), t\} \leq t + |\tau(t) - t|$  for  $a \leq t \leq a + \delta$ . Then in view of condition (2.49) we get

$$\left| \int_t^{\tau(t)} (s - a)^\alpha ds \right| \leq k_1[(t - a) + k_1(t - a)^\beta]^\alpha (t - a)^\beta \quad \text{for } a \leq t \leq a + \delta.$$

The last inequality yields the validity of our lemma. □

Analogously one can prove

**Lemma 2.9.** Let  $\tau_j \in M(]a, b[)$ ,  $\alpha \geq 0$ ,  $\beta \geq 0$  and let there exist  $\delta \in ]0, b - a[$  such that

$$(2.50) \quad |\tau_j(t) - t| \leq k_1(b - t)^\beta \quad \text{for } b - \delta \leq t < b.$$

Then

$$\left| \int_t^{\tau(t)} (b - t)^\alpha ds \right| \leq \begin{cases} k_1[1 + k_1\delta^{\beta-1}]^\alpha (b - t)^{\alpha+\beta} & \text{for } \beta \geq 1, \\ k_1[\delta^{1-\beta} + k_1]^\alpha (b - t)^{\alpha+\beta} & \text{for } 0 \leq \beta < 1 \end{cases}$$

for  $b - \delta \leq t < b$ .

### 2.3. Lemmas on the solutions of auxiliary problems.

Throughout this section we assume that the operator

$$P: C_1^{m-1}(]a, b[) \times C_1^{m-1}(]a, b[) \rightarrow L_n(]a, b[)$$

is  $\gamma_0, \gamma$  consistent with boundary condition (1.2), and the operator  $q: C_1^{m-1}(]a, b[) \rightarrow \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  is continuous.

Consider for any  $x \in \tilde{C}_1^{m-1}(]a, b[) \subset C_1^{m-1}(]a, b[)$  the nonhomogeneous equation

$$(2.51) \quad y^{(n)}(t) = \sum_{i=1}^m p_i(x)(t)y^{(i-1)}(\tau_i(t)) + q(x)(t),$$

and the corresponding homogeneous equation

$$(2.52) \quad y^{(n)}(t) = \sum_{i=1}^m p_i(x)(t)y^{(i-1)}(\tau_i(t)),$$

and let  $E^n$  be the set of solutions of problem (2.51), (2.26).

From inequality (1.23) of item (ii) of Definition 1.1 it follows that the boundary problem (2.51), (2.26) has a unique solution  $y$  in the space  $\tilde{C}^{n-1, m}(]a, b[)$ . But in view of Lemma 2.6 it is clear that  $y \in \tilde{C}_1^{m-1}(]a, b[)$ . Thus  $E^n \cap \tilde{C}_1^{m-1}(]a, b[) \neq \emptyset$ , and there exists the operator  $U: \tilde{C}_1^{m-1}(]a, b[) \rightarrow E^n \cap \tilde{C}_1^{m-1}(]a, b[)$  defined by the equality

$$U(x)(t) = y(t).$$

**Lemma 2.10.**  $U: \tilde{C}_1^{m-1}(]a, b[) \rightarrow E^n \cap \tilde{C}_1^{m-1}(]a, b[)$  is a continuous operator.

**Proof.** Let  $x_k \in \tilde{C}_1^{m-1}(]a, b[)$  and  $y_k(t) = U(x_k)(t)$  ( $k = 1, 2$ ),  $y = y_2 - y_1$ , and let the operator  $P$  be defined by (1.19). Then

$$y^{(n)}(t) = P(x_2, y)(t) + q_0(x_1, x_2)(t)$$

where  $q_0(x_1, x_2)(t) = P(x_2, y_1)(t) - P(x_1, y_1)(t) + q(x_2)(t) - q(x_1)(t)$ . Hence, by item (ii) of Definition 1.1 we have

$$\|U(x_2) - U(x_1)\|_{\tilde{C}_1^{m-1}} \leq \gamma \|q_0(x_1, x_2)\|_{\tilde{L}_{2n-2m-2, 2m-2}^2}.$$

Since the operators  $P$  and  $q$  are continuous, this estimate implies the continuity of the operator  $U$ . □

### 3. PROOFS

**P r o o f** of Remark 1.1. Let  $x$  be a solution of problem (1.8), (1.2), then inequalities (2.27) imply the estimate

$$(3.1) \quad |x^{(i-1)}(t)| \leq \frac{[(b-t)(t-a)]^{m-i+1/2}}{(m-i)!(2m-2i+1)^{1/2}} \left(\frac{2}{b-a}\right)^{m-i+1/2} \|x^{(m)}\|_{L^2}$$

for  $a \leq t \leq b$ . This estimate, by the definition of the norm in the space  $\tilde{C}^{m-1}(]a, b[)$  and estimate (1.17) immediately yields (1.18).  $\square$

**P r o o f** of Theorem 1.3. Let  $\delta$  and  $\lambda$  be the functions and numbers appearing in Definition 1.1. We set

$$(3.2) \quad \eta(t) = \delta(t, \gamma_0)\gamma_0 + \tilde{F}_p(t, \min\{2\varrho_0, \gamma_0\}),$$

$$(3.3) \quad \chi(s) = \begin{cases} 1 & \text{for } 0 \leq s \leq \varrho_0, \\ 2 - s/\varrho_0 & \text{for } \varrho_0 < s < 2\varrho_0, \\ 0 & \text{for } s \geq 2\varrho_0, \end{cases}$$

$$(3.4) \quad q(x)(t) = \chi(\|x\|_{\tilde{C}_1^{m-1}})F_p(x)(t).$$

From (1.24) it is clear that the nonnegative functions  $\tilde{F}_p, \eta$ , admit the inclusion

$$(3.5) \quad \tilde{F}_p(\cdot, \min\{2\varrho_0, \gamma_0\}), \eta \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[),$$

and for every  $x \in A_{\gamma_0} \subset \tilde{C}_1^{m-1}(]a, b[)$  and almost all  $t \in ]a, b[$  the inequality

$$(3.6) \quad |q(x)(t)| \leq \tilde{F}_p(t, \min\{2\varrho_0, \gamma_0\}) \quad \text{for } a < t < b$$

holds.

Let  $U: A_{\gamma_0} \rightarrow E^n \cap \tilde{C}_1^{m-1}(]a, b[)$  be the operator appearing in Lemma 2.10, from which it follows that  $U$  is a continuous operator. On the other hand, from items (i) and (ii) of Definition 1.1, (1.25) and (3.6) it is clear that for each  $x \in A_{\gamma_0}$ , the conditions

$$\|y\|_{\tilde{C}_1^{m-1}} \leq \gamma_0, \quad |y^{(n-1)}(t) - y^{(n-1)}(s)| \leq \int_s^t \eta(\xi) \, d\xi \quad \text{for } a < t < b$$

hold. Thus in view of Definition 2.1 the operator  $U$  maps the ball  $A_{\gamma_0}$  into its own subset  $S(\varrho_1, \eta)$ . From Lemma 2.2 it follows that  $S(\varrho_1, \eta)$  is a compact subset of the ball  $A_{\gamma_0} \subset \tilde{C}_1^{m-1}(]a, b[)$ , i.e. the operator  $u$  maps the ball  $A_{\gamma_0}$  into its own compact

subset. Therefore, owing to Schauders's principle, there exists  $x \in S(\varrho_1, \eta) \subset A_{\gamma_0}$  such that

$$x(t) = U(x)(t) \quad \text{for } a < t < b.$$

Thus by (2.51) and notation (3.4), the function  $x$  ( $x \in A_{\gamma_0}$ ) is a solution of problem (1.26), (1.2), where

$$(3.7) \quad \lambda = \chi(\|x\|_{\tilde{C}_1^{m-1}}).$$

If  $\gamma_0 = \varrho_0$  then in view of condition  $x \in A_{\gamma_0}$ , by (3.3) we have that  $\lambda = 1$ , and then in view of (2.51) and (3.4) the function  $x$  is a solution of problem (1.1), (1.2) which admits the estimate (1.27).

Let us show now that  $x$  admits estimate (1.27) in the case when  $\varrho_0 < \gamma_0$ . Assume the contrary. Then either

$$(3.8) \quad \varrho_0 < \|x\|_{\tilde{C}_1^{m-1}} < 2\varrho_0,$$

or

$$(3.9) \quad \|x\|_{\tilde{C}_1^{m-1}} \geq 2\varrho_0.$$

If condition (3.8) holds, then by virtue of (3.3) and (3.7) we have that  $\lambda \in ]0, 1[$ , which by the conditions of our theorem guarantees the validity of estimate (1.27). But this contradicts (3.8).

Assume now that (3.9) is fulfilled. Then by virtue of (3.3) and (3.7) we have that  $\lambda = 0$ . Therefore  $x \in A_{\gamma_0}$  is a solution of problem (2.52), (1.2). Thus from item (ii) of Definition 1.1 it is obvious that  $x \equiv 0$ , because problem (2.52), (1.2) has only the trivial solution. But this contradicts condition (3.9), i.e. estimate (1.27) is valid. From estimate (1.27) and (3.3) we have that  $\lambda = 1$ , and then in view of (2.51) and (3.4) the function  $x$  is a solution of problem (1.1), (1.2) which admits the estimate (1.27).  $\square$

**P r o o f** of Corollary 1.2. First note that in view of condition (1.30) there exists such  $\gamma_0 > 2\varrho_0$  that condition (1.25) holds, and in view of Definition 1.2 the operator  $P$  is  $\gamma_0, \gamma$  consistent.

On the other hand, (1.30) implies the existence of a number  $\varrho_0$  such that

$$(3.10) \quad \gamma\|\eta(\cdot, \varrho)\|_{\tilde{L}_{2m-2m-2, 2m-2}^2} < \varrho \quad \text{for } \varrho > \varrho_0.$$

Let  $x$  be a solution of problem (1.26), (1.2) for some  $\lambda \in ]0, 1[$ . Then  $y = x$  is also a solution of problem (1.22), (1.2) where  $q(t) = \lambda(F(x)(t) - P(x, x)(t))$ . Let now

$\varrho = \|x\|_{\tilde{C}_1^{m-1}}$  and assume that

$$(3.11) \quad \varrho > \varrho_0$$

holds. Then in view of the  $\gamma$ -consistency of the operator  $p$  with boundary conditions (1.2), inequality (1.23) holds and thus by condition (1.28) we have

$$\varrho = \|x\|_{\tilde{C}_1^{m-1}} \leq \gamma \|q(x)\|_{\tilde{L}_{2n-2m-2, 2m-2}^2} \leq \gamma \|\eta(\cdot, \varrho)\|_{\tilde{L}_{2n-2m-2, 2m-2}^2}.$$

But the last inequality contradicts (3.10). Thus assumption (3.11) is not valid and  $\varrho \leq \varrho_0$ . Therefore for any  $\lambda \in ]0, 1[$  an arbitrary solution of problem (1.26), (1.2) admits the estimate (1.27). Therefore all the conditions of Theorem 1.3 are fulfilled, from which the solvability of problem (1.1), (1.2) follows.  $\square$

**P r o o f** of Theorem 1.4. Let  $r_n$  be the constant defined in Remark 1.1. First we prove that the operator  $P$  is  $\gamma_0, r_n$  consistent with boundary conditions (1.2). From the conditions of our theorem it is obvious that the item (i) of Definition 1.1 is satisfied. Let now  $x$  be an arbitrary fixed function from the set  $A_{\gamma_0}$  and let  $p_j(t) \equiv p_j(x)(t)$ . Thus in view of (1.34), (1.35) all the assumptions of Theorem 1.1 are satisfied, and then for any  $q \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  problem (1.22), (1.2) has a unique solution  $y$ . Also in view of Remark 1.1 there exists a constant  $r_n > 0$  (which depends only on the numbers  $l_{kj}, \bar{l}_{kj}, \gamma_{kj}$  ( $k = 0, 1; j = 1, \dots, m$ ) and  $a, b, t^*, n$ ) such that estimate (1.23) holds with  $\gamma = r_n$ . So, the operator  $P$  is  $\gamma_0, r_n$  consistent with boundary conditions (1.2). Therefore all the assumptions of Corollary 1.1 are fulfilled, from which the solvability of problem (1.1), (1.2) follows.  $\square$

**P r o o f** of Theorem 1.5. Let  $r_n$  be the constant defined in Remark 1.1. First we prove that the operator  $P$  is  $r_n$  consistent with boundary conditions (1.2). From the conditions of our theorem it is obvious that the item (i) of Definition 1.1 is satisfied. Let now  $\gamma_0$  be an arbitrary nonnegative number,  $x$  an arbitrary fixed function from the space  $A_{\gamma_0}$  and let  $p_j(t) \equiv p_j(x)(t)$ . Then in view of (1.37), (1.38) all the assumptions of Theorem 1.1 are satisfied and then for any  $q \in \tilde{L}_{2n-2m-2, 2m-2}^2(]a, b[)$  problem (1.22), (1.2) has a unique solution  $y$ . Also in view of Remark 1.1 there exists a constant  $r_n > 0$  (which depends only on the numbers  $l_{kj}, \bar{l}_{kj}, \gamma_{kj}$  ( $k = 0, 1; j = 1, \dots, m$ ) and  $a, b, t^*, n$ ) such that estimate (1.23) holds with  $\gamma = r_n$ . So, the operator  $P$  is  $\gamma_0, r_n$  consistent with boundary conditions (1.2) for arbitrary  $\gamma_0 > 0$ . Thus by Definition 1.1, the operator  $P$  is  $r_n$  consistent with boundary conditions (1.2). Therefore all the assumptions of Corollary 1.2 are fulfilled, from which the solvability of problem (1.1), (1.2) follows.  $\square$

Proof of Remark 1.2. By Schwartz's inequality, the definition of the norm  $\|y\|_{\tilde{C}_1^{m-1}}$  and inequalities (1.39), (2.2) for any  $x, y \in A_{\gamma_0}$  and  $z = y - x$  we have

$$(3.12) \quad |p_j(y)(t)z^{(j-1)}(\tau_j(t))| = |p_j(y)(t)z^{(j-1)}(t)| + |p_j(y)(t)| \left| \int_t^{\tau_j(t)} z^{(j)}(\psi) d\psi \right| \\ \leq \|z\|_{\tilde{C}_1^{m-1}} |p_j(y)(t)| \alpha_j(t) \left( 1 + \frac{1}{\alpha_j(t)} \left( \int_t^{\tau_j(t)} (\psi - a)^{2m-2j} d\psi \right)^{1/2} \right)$$

for  $a < t < b$ . On the other hand, from the conditions (1.40) by Lemmas 2.8 and 2.9 it is clear that

$$\alpha_j^{-1}(s) \left( \int_s^{\tau_j(s)} (\xi - a)^{2m-2j} d\xi \right)^{1/2} \leq \frac{\sqrt{\kappa(1+\kappa)}}{\varepsilon^{m-j+1/2}} \quad \text{for } s \in ]a, a + \varepsilon] \cup [b - \varepsilon, b[, \\ \alpha_j^{-1}(s) \left( \int_s^{\tau_j(s)} (\xi - a)^{2m-2j} d\xi \right)^{1/2} \leq \varepsilon^{-2m+2j-1} \left( \int_a^b (\xi - a)^{2m-2j} d\xi \right)^{1/2} \\ = \frac{(b-a)^{m-j+1/2}}{\sqrt{2m-2j+1} \varepsilon^{2m-2j+1}} \quad \text{for } s \in ]a + \varepsilon, b - \varepsilon[.$$

Then if we put

$$(3.13) \quad \kappa_0 = \max_{1 \leq j \leq m} \left\{ \frac{\sqrt{\kappa(1+\kappa)}}{\varepsilon^{m-j+1/2}}, \frac{(b-a)^{m-j+1/2}}{\sqrt{2m-2j+1} \varepsilon^{2m-2j+1}} \right\},$$

from (3.12) by the last estimates we get the inequality

$$(3.14) \quad |p_j(y)(t)z^{(j-1)}(\tau_j(t))| \leq \|z\|_{\tilde{C}_1^{m-1}} (1 + \kappa_0) |p_j(y)(t)| \alpha_j(t) \\ \leq \|z\|_{\tilde{C}_1^{m-1}} (1 + \kappa_0) \delta_j(t, \|y\|_{\tilde{C}_1^{m-1}})$$

for  $a < t < b$ . Analogously we get that

$$|(p_j(y)(t) - p_j(x)(t))x^{(j-1)}(\tau_j(t))| \leq \|x\|_{\tilde{C}_1^{m-1}} (1 + \kappa_0) |p_j(y)(t) - p_j(x)(t)| \alpha_j(t)$$

for  $a < t < b$ . From (3.14) and the last inequality it is obvious that the operator  $P$  defined by equality (1.19) is continuously acting from  $A_{\gamma_0}$  to the space  $L_n(]a, b[)$ , and the item (ii) of Definition 1.1 holds with  $\delta(t, \varrho) = (1 + \kappa_0) \sum_{j=1}^m \delta_j(t, \varrho)$ .  $\square$

Proof of Corollary 1.3. From conditions (1.42) and (1.40) by Remark 1.2 we obtain that the operator  $P$  defined by equality (1.19) with  $p_j(x)(t) = p_j(t)$  is continuously acting from  $A_{\gamma_0}$  to the space  $L_n(]a, b[)$  for any  $\gamma_0 > 0$ , i.e., it is continuously acting from  $\tilde{C}_1^{m-1}(]a, b[)$  to the space  $L_n(]a, b[)$ .

Therefore it is clear that all the conditions of Theorem 1.5 are satisfied with

$$F(x)(t) = f(t, x(\tau_1(t)), x'(\tau_2(t)), \dots, x^{(m-1)}(\tau_m(t))), \quad \delta(t, \varrho) = (1 + \kappa_0) \sum_{j=1}^m |p_j(t)|,$$

where the constant  $\kappa_0$  is defined by equality (3.13). Thus problem (1.41), (1.2) is solvable.  $\square$

**P r o o f** of Corollary 1.4. Let the operators  $F, p_1: C^{m-1}(]a, b[) \rightarrow L_{\text{loc}}(]a, b[)$ , and the function  $\eta: ]a, b[ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be defined by equalities

$$F(x)(t) = -\frac{\lambda|x(t)|^k}{[(t-a)(b-t)]^{2+k/2}}x(\tau(t))+q(x)(t), \quad p_1(x)(t) = -\frac{\lambda|x(t)|^k}{[(t-a)(b-t)]^{2+k/2}}.$$

Then it is easy to verify that in view of (1.46), (1.47), (1.48), conditions (1.13), (1.14), (1.28), (1.34), (1.35), (1.36), (1.37), (1.38), (1.39), (1.40), (1.41), (1.42), (1.43) are satisfied with

$$(3.15) \quad \delta(t, \varrho) = \frac{\varrho^k \lambda}{[(t-a)(b-t)]^2}, \quad l_{01} = l_{11} = \frac{4\gamma_0^k \lambda}{(b-a)^2}, \quad \bar{l}_{01} = \bar{l}_{11} = \frac{16\gamma_0^k \lambda}{(b-a)^2},$$

$$r_2 = \left(1 + \sqrt{\frac{2}{b-a}}\right) \frac{2(1+b-a)(b-a)^2}{(b-a)^2 - 16\lambda\gamma_0^k(1 + [2(b-a)]^{1/4})},$$

$$B_0 = B_1 = \frac{16\lambda\gamma_0^k}{(b-a)^2}(1 + [2(b-a)]^{1/4}), \quad t^* = \frac{a+b}{2}, \quad \gamma_{01} = \gamma_{11} = \frac{1}{4}.$$

Thus all the condition of Theorem 1.4 are satisfied, from which solvability of problem (1.44), (1.2) follows.  $\square$

**P r o o f** of Corollary 1.5. Let the operators  $F, p_1: C^{m-1}(]a, b[) \rightarrow L_{\text{loc}}(]a, b[)$ , and the function  $\eta: ]a, b[ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be defined by equalities

$$F(x)(t) = -\frac{\lambda|\sin x^k(t)|}{[(t-a)(b-t)]^2}x(\tau(t)) + q(x)(t), \quad p_1(x)(t) = -\frac{\lambda|\sin x^k(t)|}{[(t-a)(b-t)]^2}.$$

Then it is easy to verify that in view of (1.30), (1.46), and (1.49), all the conditions of Theorem 1.5 are fulfilled, where  $\delta, l_{01}, \bar{l}_{01}, r_2, B_0, B_1, t^*, \gamma_{01}, \gamma_{11}$ , are defined by (3.15) with  $\varrho = 1, \gamma_0 = 1$ , which implies solvability of problem (1.44), (1.2).  $\square$

## References

- [1] *R. P. Agarwal*: Focal Boundary Value Problems for Differential and Difference Equations. Mathematics and Its Applications, Dordrecht: Kluwer Academic Publishers, 1998.
- [2] *R. P. Agarwal, I. Kiguradze*: Two-point boundary value problems for higher-order linear differential equations with strong singularities. Bound. Value Probl., Article ID 83910 (2006), pp. 32.
- [3] *R. P. Agarwal, D. O'Regan*: Singular Differential and Integral Equations with Applications. Dordrecht: Kluwer Academic Publishers, 2003.
- [4] *E. Bravyi*: A note on the Fredholm property of boundary value problems for linear functional-differential equations. Mem. Differ. Equ. Math. Phys. 20 (2000), 133–135.
- [5] *T. I. Kiguradze*: On conditions for linear singular boundary value problems to be well posed. Differ. Equ. 46 (2010), 187–194; translation from Differ. Uravn. 46 (2010), 183–190.
- [6] *I. Kiguradze*: On two-point boundary value problems for higher order singular ordinary differential equations. Mem. Differ. Equ. Math. Phys. 31 (2004), 101–107.
- [7] *I. T. Kiguradze*: On a singular multi-point boundary value problem. Ann. Mat. Pura Appl., IV. Ser. 86 (1970), 367–399.
- [8] *I. Kiguradze*: Some optimal conditions for the solvability of two-point singular boundary value problems. Funct. Differ. Equ. 10 (2003), 259–281.
- [9] *I. T. Kiguradze*: Some Singular Boundary Value Problems for Ordinary Differential Equations. Tbilisi University Press, 1975. (In Russian.)
- [10] *I. Kiguradze*: The Dirichlet and focal boundary value problems for higher order quasi-halflinear singular differential equations. Mem. Differential Equ. Math. Phys. 54 (2011), 126–133.
- [11] *I. T. Kiguradze, T. A. Chanturiya*: Asymptotic Properties of Solutions of Nonautonomous Ordinary Differential Equations. Translation from the Russian. Mathematics and Its Applications. Soviet Series. 89, Dordrecht, Kluwer Academic Publishers, 1993.
- [12] *I. Kiguradze, B. Pūža, I. P. Stavroulakis*: On singular boundary value problems for functional-differential equations of higher order. Georgian Math. J. 8 (2001), 791–814.
- [13] *I. Kiguradze, G. Tskhovrebadze*: On two-point boundary value problems for systems of higher order ordinary differential equations with singularities. Georgian Math. J. 1 (1994), 31–45.
- [14] *A. G. Lomtadze*: A boundary value problem for nonlinear second order ordinary differential equations with singularities. Differ. Equations 22 (1986), 301–310; translation from Differ. Uravn. 22 (1986), 416–426.
- [15] *S. Mukhigulashvili*: On one estimate for solutions of two-point boundary value problems for strongly singular higher-order linear differential equations. Mem. Differential Equ. Math. Phys. 58 (2013).
- [16] *S. Mukhigulashvili*: Two-point boundary value problems for second order functional differential equations. Mem. Differ. Equ. Math. Phys. 20 (2000), 1–112.
- [17] *S. Mukhigulashvili, N. Partsvania*: On two-point boundary value problems for higher order functional differential equations with strong singularities. Mem. Differ. Equ. Math. Phys. 54 (2011), 134–138.
- [18] *S. Mukhigulashvili, N. Partsvania*: Two-point boundary value problems for strongly singular higher-order linear differential equations with deviating arguments. Electron. J. Qual. Theory Differ. Equ. 38 (2012), 1–34.
- [19] *N. Partsvania*: On extremal solutions of two-point boundary value problems for second order nonlinear singular differential equations. Bull. Georgian Natl. Acad. Sci. (N.S.) 5 (2011), 31–36.

- [20] *N. Partsvania*: On solvability and well-posedness of two-point weighted singular boundary value problems. *Mem. Differ. Equ. Math. Phys.* 54 (2011), 139–146.
- [21] *B. Půža*: On a singular two-point boundary value problem for a nonlinear  $m$ th-order differential equation with deviating arguments. *Georgian Math. J.* 4 (1997), 557–566.
- [22] *B. Půža, A. Rabbimov*: On a weighted boundary value problem for a system of singular functional-differential equations. *Mem. Differ. Equ. Math. Phys.* 21 (2000), 125–130.
- [23] *I. Rachůnková, S. Staněk, M. Tvrđý*: Singularities and Laplacians in Boundary Value Problems for Nonlinear Ordinary Differential Equations. *Handbook of differential equations: ordinary differential equations*. Vol. III, *Handb. Differ. Equ.*, Elsevier/North-Holland, Amsterdam, 2006.
- [24] *I. Rachůnková, S. Staněk, M. Tvrđý*: Solvability of Nonlinear Singular Problems for Ordinary Differential Equations. *Contemporary Mathematics and Its Applications*, 5, Hindawi Publishing Corporation, New York, 2008.
- [25] *Š. Schwabik, M. Tvrđý, O. Vejvoda*: *Differential and Integral Equations. Boundary Value Problems and Adjoints*. D. Reidel Publishing Co., Dordrecht, Boston, London, 1979.

*Author's addresses:* Sul Khan Mukhigulashvili, Mathematical Institute, Academy of Sciences of the Czech Republic, Žitkova 22, 616 62 Brno, Czech Republic; I. Chavchavadze State University, Faculty of physics and mathematics, I. Chavchavadze St. No. 32, Tbilisi 0179, Georgia, e-mail: mukhig@ipm.cz.